

EQUATIONS AND CONSTANTS FOR GRADE 12 PHYSICS

These equations are meant to make doing homework and exams a bit easier. They are **not** an excuse for not learning the course material. If you don't know what these equations mean and how to use them, they will not help you at all. Vector quantities are expressed in **bold fonts**.

KINEMATICS (constant acceleration):

$$\begin{array}{llll} \mathbf{v} = \frac{\Delta \mathbf{d}}{\Delta t} & \Delta \mathbf{d} = \mathbf{v}_1 \Delta t + \frac{1}{2} \mathbf{a} \Delta t^2 & \Delta \mathbf{d} = \frac{\mathbf{v}_1 + \mathbf{v}_2}{2} \Delta t & v_2^2 = v_1^2 + 2a\Delta d \\ \mathbf{a} = \frac{\Delta \mathbf{v}}{\Delta t} & \Delta \mathbf{d} = \mathbf{v}_2 \Delta t - \frac{1}{2} \mathbf{a} \Delta t^2 & \mathbf{v}_2 = \mathbf{v}_1 + \mathbf{a} \Delta t & \end{array}$$

SYMMETRIC PROJECTILE MOTION:

$$\text{Time of flight: } t_{\text{tot}} = \frac{2v_i \sin(\theta)}{g} \quad \text{Range: } R = \frac{v_i^2 \sin(2\theta)}{g} \quad \text{Max height: } h_{\text{max}} = \frac{v_i^2 \sin^2(\theta)}{2g}$$

UNIFORM CIRCULAR MOTION:

$$\mathbf{a}_c \perp \mathbf{v} \quad a_c = \frac{v^2}{r} \quad F_c = ma_c \quad T = \frac{2\pi r}{v} \quad f = \frac{1}{T} \quad \text{Bank angle, no friction: } \tan \theta = \frac{v^2}{rg}$$

NEWTON'S LAWS, MOMENTUM, IMPULSE, KINETIC ENERGY:

$$\mathbf{F}_{\text{net}} = m\mathbf{a} = \frac{\Delta \mathbf{p}}{\Delta t} \quad \mathbf{F}_{A \text{ on } B} = -\mathbf{F}_{B \text{ on } A} \quad \mathbf{p} = m\mathbf{v} \quad \mathbf{J} = \Delta \mathbf{p} = \mathbf{F}_{\text{ave}} \Delta t \quad W = Fd \cos \theta \quad K = \frac{1}{2}mv^2$$

COLLISIONS:

$$\begin{array}{l} \mathbf{p}_A + \mathbf{p}_B = \mathbf{p}'_A + \mathbf{p}'_B \\ K_A + K_B = K'_A + K'_B \text{ (elastic)} \end{array}$$

FRICTION:

$$\begin{array}{ll} \text{Static:} & \max F_s = \mu_s F_N \\ \text{Kinetic:} & F_k = \mu_k F_N \end{array}$$

SPRING:

$$\begin{array}{ll} \text{Hooke's Law:} & \mathbf{F} = k\mathbf{x} \\ \text{Elastic Potential:} & U_e = \frac{1}{2}kx^2 \end{array}$$

VELOCITIES AND ENERGIES IN ORBIT (m orbiting M):

$$\begin{array}{lll} v_{\text{escape}} = \sqrt{\frac{2GM}{r}} & K_{\text{orbit}} = \frac{GMm}{2r} = \frac{1}{2}mv_{\text{orbit}}^2 & E_{\text{tot}} = K_{\text{orbit}} + U_g = -\frac{GMm}{2r} = -K_{\text{orbit}} \\ v_{\text{orbit}} = \sqrt{\frac{GM}{r}} & U_{\text{orbit}} = -\frac{GMm}{r} = -2K_{\text{orbit}} & \frac{T^2}{r^3} = \text{constant} \end{array}$$

FORCE FIELDS:

$$\begin{array}{llll} F_g = \frac{Gm_1m_2}{r^2} & U_g = -\frac{Gm_1m_2}{r} & g = \frac{Gm_s}{r^2} & \mathbf{F}_g = m\mathbf{g} \\ F_q = \frac{kq_1q_2}{r^2} & U_q = \frac{kq_1q_2}{r} & E = \frac{kq_s}{r^2} & \mathbf{F}_q = q\mathbf{E} \\ F_M = qvB \sin \theta & F_M = IlB \sin \theta & V = \frac{kq_s}{r} & E = \frac{\Delta V}{d} \text{ (Parallel plate)} \end{array}$$

SINGLE-SLIT DIFFRACTION:

$$\begin{array}{ll} \text{Bright fringes:} & \pm \left(m + \frac{1}{2}\right) \lambda = W \sin \theta \quad y_m = \left(m + \frac{1}{2}\right) \frac{\lambda L}{W} \\ \text{Dark fringes:} & \pm m \lambda = W \sin \theta \quad y_m = \frac{m \lambda L}{W} \quad \text{for } m = 1, 2, 3, \dots \end{array}$$

DOUBLE-SLIT INTERFERENCE & DIFFRACTION GRATING:

Bright fringe: $\pm n\lambda = d \sin \theta$ Dark: $\pm \left(n + \frac{1}{2}\right) \lambda = d \sin \theta$ $\lambda \approx \frac{\Delta y d}{x}$

REFRACTION:

$$n_1 \sin \theta_1 = n_2 \sin \theta_2 \quad n = \frac{c}{v}$$

OPTICAL RESOLUTION: (θ in radians)

Rectangular aperture: $\theta_{\min} = \frac{\lambda}{W}$

Circular aperture: $\theta_{\min} = \frac{1.22\lambda}{D}$

UNIVERSAL WAVE EQUATION:

$$v = f\lambda$$

SPECIAL RELATIVITY:

$$\gamma = \frac{1}{\sqrt{1 - \left(\frac{v}{c}\right)^2}} \quad \Delta t = \gamma \Delta t_0 \quad L = \frac{L_0}{\gamma} \quad m = \gamma m_0 \quad K = mc^2 - m_0 c^2$$

QUANTUM MECHANICS:

$$E = hf \quad K_{\max} = \begin{cases} hf - \phi & \text{if } hf > \phi \\ 0 & \text{otherwise} \end{cases} \quad p = \frac{E}{c} = \frac{hf}{c} = \frac{h}{\lambda} \quad \lambda = \frac{h}{mv} \quad \sigma_p \sigma_x \geq \frac{1}{2} \hbar \quad \hbar = \frac{h}{2\pi}$$

USEFUL CONSTANTS:

Acceleration due to gravity	$g = 9.81 \text{ m/s}^2$ (near surface of Earth)
Gravitational constant	$G = 6.674 \times 10^{-11} \text{ N m}^2/\text{kg}^2$
Coulomb's constant	$k = 8.988 \times 10^9 \text{ N m}^2/\text{C}^2$
Electron rest mass	$m_e = 9.11 \times 10^{-31} \text{ kg}$
Proton rest mass	$m_p = 1.673 \times 10^{-27} \text{ kg}$
Elementary charge	$e = 1.602 \times 10^{-19} \text{ C}$
Speed of light	$c = 2.998 \times 10^8 \text{ m/s}$
Planck's Constant	$h = 6.626 \times 10^{-34} \text{ J s}$
Mass of Earth	$5.972 \times 10^{24} \text{ kg}$
Radius of Earth	$6.371 \times 10^6 \text{ m}$
Mass of Sun	$1.989 \times 10^{30} \text{ kg}$
Radius of sun	$6.957 \times 10^8 \text{ m}$
Mass of Moon	$7.348 \times 10^{22} \text{ kg}$
Radius of Moon	$1.737 \times 10^6 \text{ m}$
Earth-to-moon distance	$3.844 \times 10^8 \text{ m}$ (centre to centre)

CONVERSION TO SI UNITS:

Electron volt	$1 \text{ eV} = 1.602 \times 10^{-19} \text{ J}$
Kilowatt-hour	$1 \text{ kWh} = 3.6 \times 10^6 \text{ J}$
Kilometres per hour	$1 \text{ km/h} = 0.278 \text{ m/s}$, $1 \text{ m/s} = 3.6 \text{ km/h}$
Light year	$1 \text{ ly} = 9.461 \times 10^{15} \text{ m}$

UNIT PREFIXES:

tera	10^{12}	T
giga	10^9	G
mega	10^6	M
kilo	10^3	k
centi	10^{-2}	c
milli	10^{-3}	m
micro	10^{-6}	μ
nano	10^{-9}	n

MATHEMATICAL FORMULAS:

Circumference of a circle	$C = 2\pi r$
Area of a circle	$A = \pi r^2$
Volume of a sphere	$V = \frac{4}{3}\pi r^3$
Density	$\rho = m/V$
Small-angle approximation	$\tan \theta \approx \sin \theta \approx \theta$ (θ in radians)