

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e1f38f83c45700544e3d4b

Fecha de creación: 2016-03-10T22:22:13.918Z

annualCycles

annualCycleUnstructured:

habitats

habitatUnstructured:

[commonNamesAtomized](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:

dataType:

identifier:

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Español

name: Avispa

creation\_date: 2016-03-10T22:22:13.918Z

usesManagementAndConservation

## ancillaryData

managementAndConservationUnstructured:

## managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataFormat:

identifier:

fullDescriptionUnstructured: Húmero no se proyecta a la altura de la tégula; prominencia anterolateral del pronoto completa o casi totalmente pálida. Especie negra con marcas amarillas.

briefDescription:

## references

keywords: *Agelaia angulata angulata*, *Agelaia angulicollis*, *Agelaia brevistigma*, *Agelaia cajennensis*, *Agelaia fulvofasciata*, *Agelaia hamiltoni*, *Agelaia testacea*, *Angiopolybia obidensis*, *Angiopolybia pallens*, *Angiopolybia paraensis*, *Angiopolybia zischkai*, *Apoica thoracica*, *Asteloeca trilli*, *Brachygastra albula*, *Brachygastra augusti*, *Brachygastra lecheguana*, *Brachygastra propodealis*, *Brachygastra scutellaris*, *Brachymenes wagnerianus*, *Chartergellus amazonicus*, *Chartergellus punctatior*, *Chartergus chartarius*, *Chartergus globiventris*, *Chartergus metanotalis*, *Clypearia sulcata*, *Clypearia weyrauchi*, *Epipona guerini*, *Epipona tatua*, *Eumenes versicolor filiformis*, *Leipomeles dorsata*, *Metapolybia acincta*, *Metapolybia docilis*, *Metapolybia nigra*, *Metapolybia suffusa*, *Mischocyttarus adolphi*, *Mischocyttarus alfkeni*, *Mischocyttarus alfkenii excrucians*, *Mischocyttarus alfkenii trinitatis*, *Mischocyttarus carbonarius carbonarius*, *Mischocyttarus flavicans*, *Mischocyttarus foveatus*, *Mischocyttarus injucundus tingomariae*, *Mischocyttarus insolitus*, *Mischocyttarus labiatus*, *Mischocyttarus latissimus*, *Mischocyttarus metoecus*, *Mischocyttarus petiolatus*, *Mischocyttarus pseudomimeticus*, *Mischocyttarus punctatus*, *Mischocyttarus tomentosus*, *Mischocyttarus vaqueroi*, *Omicron aequale nigrum*, *Omicron conclamatum*, *Omicron criticum pulchripictum*, *Omicron foxi*, *Omicron globicolle*, *Omicron graculum atripes*, *Omicron lustratum*, *Omicron nanum incarum*, *Omicron paranympus*, *Omicron ypsilon*, *Pachymenes ghilianii olivaceus*, *Pachymenes orellanae orellanae*, *Pachymenes orellanae vardei*, *Parachartergus amazonensis*, *Parachartergus colobopterus*, *Parachartergus fasciipennis*, *Parachartergus fraternus*, *Parachartergus fulgidipennis*, *Parachartergus griseus*, *Parachartergus richardsi*, *Parachartergus smithii*, *Parachartergus weyrauchi*, *Pararhaphidoglossa clypeolaris*, *Pararhaphidoglossa confluenta confluenta*, *Pararhaphidoglossa cressoniana*, *Pararhaphidoglossa proxima*, *Polybia bistriata*, *Polybia catillifex*, *Polybia eberhardae*, *Polybia emaciata*, *Polybia fastidiosuscula*, *Polybia jurinei*, *Polybia liliacea*, *Polybia micans*, *Polybia occidentalis*, *Polybia platycephala sylvestris*, *Polybia quadricincta*, *Polybia raii raii*, *Polybia rejecta*, *Polybia scrobalis pronotalis*, *Polybia scrobalis scrobalis*, *Polybia sericea*, *Polybia striata*, *Polybia velutina*, *Protopolybia acutiscutis*, *Protopolybia amarella*, *Protopolybia chartergoides chartergoides*, *Protopolybia*

exigua binominata,Protopolybia exigua exigua,Protopolybia fuscatus,Protopolybia rugulosa,Protopolybia sedula,Pseudopolybia compressa,Pseudopolybia vespiceps,Santamenes peregrinus,Synoeca septentrionalis,Synoeca surinama,Synoeca virginea,Synoecoides depressus,Zethus brasiliensis fuscatus,Zethus mexicanus lugubris,Zethus mexicanus mexicanus

## editors

institution: Universidad Nacional de Colombia

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 1997

## authors

authors: Sarmiento, C. E.

title: Véspidos de Colombia (Hymenoptera: Vespidae)

source:

type: thesis

tags: Agelaia angulata angulata,Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Asteloea trilli,Brachygastra albula,Brachygastra augusti,Brachygastra lecheguana,Brachygastra propodealis,Brachygastra scutellaris,Brachymenes wagnerianus,Chartergellus amazonicus,Chartergellus punctatior,Chartergus chartarius,Chartergus globiventris,Chartergus metanotalis,Clypearia sulcata,Clypearia weyrauchi,Epipona guerini,Epipona tatua,Eumenes versicolor filiformis,Leipomeles dorsata,Metapolybia acincta,Metapolybia docilis,Metapolybia nigra,Metapolybia suffusa,Mischocyttarus adolphi,Mischocyttarus alfkeni,Mischocyttarus alfkenii excrucians,Mischocyttarus alfkenii trinitatis,Mischocyttarus carbonarius carbonarius,Mischocyttarus flavicans,Mischocyttarus foveatus,Mischocyttarus injucundus tingomariae,Mischocyttarus insolitus,Mischocyttarus labiatus,Mischocyttarus latissimus,Mischocyttarus metoecus,Mischocyttarus petiolatus,Mischocyttarus pseudomimeticus,Mischocyttarus punctatus,Mischocyttarus tomentosus,Mischocyttarus vaqueroi,Omicron aequale nigrum,Omicron conclamatum,Omicron criticum pulchripictum,Omicron foxi,Omicron globicolle,Omicron graculum atripes,Omicron lustratum,Omicron nanum incarum,Omicron paranympus,Omicron ypsilon,Pachymenes ghilianii olivaceus,Pachymenes orellanae orellanae,Pachymenes orellanae vardyi,Parachartergus amazonensis,Parachartergus colobopteris,Parachartergus fasciipennis,Parachartergus fraternus,Parachartergus fulgidipennis,Parachartergus griseus,Parachartergus richardsi,Parachartergus smithii,Parachartergus weyrauchi,Pararhaphidoglossa clypeolaris,Pararhaphidoglossa confluenta confluenta,Pararhaphidoglossa cressoniana,Pararhaphidoglossa proxima,Polybia bistrata,Polybia catillifex,Polybia eberhardae,Polybia emaciata,Polybia fastidiosuscula,Polybia jurinei,Polybia liliacea,Polybia micans,Polybia occidentalis,Polybia platycephala sylvestris,Polybia quadricincta,Polybia raii raii,Polybia rejecta,Polybia scrobalis pronotalis,Polybia scrobalis scrobalis,Polybia sericea,Polybia striata,Polybia velutina,Protopolybia acutiscutis,Protopolybia amarella,Protopolybia chartergoides chartergoides,Protopolybia exigua binominata,Protopolybia exigua exigua,Protopolybia fuscatus,Protopolybia rugulosa,Protopolybia sedula,Pseudopolybia compressa,Pseudopolybia vespiceps,Santamenes peregrinus,Synoeca septentrionalis,Synoeca surinama,Synoeca virginea,Synoecoides depressus,Zethus brasiliensis fuscatus,Zethus mexicanus lugubris,Zethus mexicanus mexicanus

abstract:

## identifiers

last\_modified:

created:

group\_id:

profile\_id:

id:  
taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

conservedAgainst

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:

publishedIn

simple:  
source:

datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: Richards, 1978  
canonicalName  
infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: Metapolybia docilis  
rank: SPECIES  
simple: Metapolybia docilis Richards, 1978

attributes

nomenclaturalCode:  
isAnamorphic:  
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:  
additionalInformation:

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

## audience

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-10T22:21:53.903Z

created: 2016-03-10T22:21:53.903Z

## agent

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

data Type:

identifier:

parentTaxon: Metapolybia

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Metapolybia

family: Vespidae

order: Hymenoptera

classHierarchy: Insecta

phylum: Arthropoda

kingdom: Animalia

recommended:

classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
reproductionUnstructured: En esta especie puede haber variación en la coloración de los individuos.

0

dataObject:  
additionalInformation:

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder:  
rights:  
license:  
modified:  
created:

agent

mimeType:  
dataType:  
identifier:  
relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedBy: TraitBank  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue: Sexual  
measurementType: Tipo de Reproducción  
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:



lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

feedingUnstructured:

0

0

strategy: Depredadores

type:

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
behaviorUnstructured: Metapolybia docilis es una especie muy poco agresiva y presenta comportamiento eusocial.

#### ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

#### measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

#### endemicAtomized

#### distribution

#### ancillaryData

dataObject:  
additionalInformation:

#### reference

geoPoint:  
location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:

dataType:

identifier:

distributionUnstructured: Se distribuye en Centroamérica y Sudamérica (México, Colombia, Venezuela, Trinidad y Tobago, Guyana, Guyana Francesa, Surinam, Ecuador y Perú) en zonas que alcanzan hasta los 1600 msnm de altitud. En Colombia se ha reportado en las regiones de la Amazonia, Orinoquía, Pacífica.

[distributionAtomized](#)

stateProvince: Valle del Cauca

[ancillaryData](#)

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

[threatStatus](#)

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Orificio propodeal más largo que ancho; valvas estrechas anteriormente y con la parte distal más ancha; gena, en perfil, más estrecha que el ojo; segmento abdominal I muy aplanado y estrecho, casi tan largo como el tórax, tubérculo espiracular prominente, especies esbeltas; margen lateral del proepisterno o pronoto con una estría crenada; ala posterior con la primera vena cubital 1/2 más larga que la cubito-anal. Puede haber variación en la coloración de los ejemplares.

[associatedParty](#)

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomez@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

## ancillaryData

dataObject:

additionalInformation: Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

Registros biológicos: Colección Entomológica, Instituto de Ciencias Naturales, Universidad Nacional de Colombia. Bogotá, Colombia <http://www.biovirtual.unal.edu.co/colecciones/datos/animalia/consultasMammalia.jsp?clase=4>  
Información de tipos: Holotipo hembra proveniente Colombia y depositado en el United States National Museum, Washington, EEUU (USNM).

Etimología del nombre científico: El epíteto específico hace referencia a su comportamiento poco agresivo.

## reference

geoPoint:

location:

thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:

rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

## agent

contentType:

dataType:

identifier:

\_id

\_bsontype: ObjectID

id: Váo•fÄW DãÔ°