Reporte de Fichas para el usuario: edith gómez Ficha ID: 56d5f6f83c16479905cba906 Fecha de creación: 2016-03-01T20:09:29.770Z annualCycles annualCycleUnstructured: habitats 0 dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Equipo Coordinador del SiB Colombia license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: habitatUnstructured: Residente de invierno poco común en bordes de selva húmeda, claros con matorral y monte secundario, ocasionalmente en el dosel de selva no perturbada (Hilty & Brown 1986). commonNamesAtomized ancillaryData usedBy: usedIn distributionUnstructured: endDate: startDate: ancillaryData type: language: Inglés name: Great Crested Flycatcher creation date: 2016-03-01T20:09:29.770Z

usesManagementAndConservation

# ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

# fullDescription

fullDescriptionUnstructured: briefDescription:

#### references

#### taxonRecordName

# ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

# replacementNameFor

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

sanctioned

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

#### laterHomonymOf

microReference:

source: datatype: identifier: source:

datatype: identifier: note: ruleConsidered:
conservedAgainst
basedOn
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
basionym
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
spellingCorrentionOf
typeNameEntity
source: datatype: identifier:

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

 ${\bf lectotype MicroReference Voucher}$ 

lectotypePublicationVoucher

# voucherReference

simple:

microReference:

year:

# publishedIn simple: source: datatype: identifier: specialAuthorship authors year simple: canonicalAuthorship authors year simple: simple: (Linnaeus, 1758) canonicalName infraspecificEpithet: specificEpithet: infragenericEpithet: linkType: ref: uninomial: simple: Myiarchus crinitus rank: SPECIES simple: Myiarchus crinitus (Linnaeus, 1758) attributes nomenclaturalCode: isAnamorphic: synonymsAtomized ancillaryData synonymStatus: synonymName anyTwo: anyOne providerLink: microReference: publishedIn source:

datatype: identifier: reletedName source: datatype: identifier: note: ruleConsidered: microReference: publishedIn

source: datatype: identifier:

#### reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

#### publishedIn

source: datatype: identifier:

#### reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

#### publishedIn

source: datatype: identifier:

#### reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

#### publishedIn

source: datatype: identifier:

#### reletedName

source: datatype: identifier:

note:

ruleConsidered: microReference:

# publishedIn

source: datatype: identifier:

#### reletedName

source: datatype: identifier: note:

ruleConsidered:

#### typeNameEntity

source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:

# typeVoucherEntity

typeOfType: simple:

microReference:

year: source: datatype: identifier:

# combinationAuthorship

# basionymAuthorship

simple:

authorship

simple:

simple: Bangs, 1898

epithet

infraspecificEpithet: specificEpithet: infragenericEpithet:

#### genus

linkType:

ref: uninomial:

simple: Myiarchus crinitus boreus

rank: SUBSPECIES

simple: Myiarchus crinitus boreus Bangs, 1898

nomenclaturalCode: isAnamorphic:

id.

# hierarchy

# ancillaryData

dataObject:

additionalInformation:

#### reference

geoPoint: location:

thumbnailURL:

### mediaURL

description:

# subject

source: The Catalogue of Life, 3rd January 2011

#### audience

bibliographicCitation:

rigthsHolder:

rights: license: title:

modified: 2016-03-01T20:09:09.769Z created: 2016-03-01T20:09:09.769Z

# agent

homepage:

role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:

identifier:

parentTaxon: Myiarchus higherClassification:

infraspecificEpithet: specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Myiarchus family: Tyrannidae order: Passeriformes classHierarchy: Aves phylum: Chordata kingdom: Animalia recommended: classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

feedingUnstructured:

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type: purpose:

behavior

dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights Equipo Coordinador del SiB Colombia license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: behaviorUnstructured: Se puede observar con frecuencia en árboles y arbustos con frutos (Hilty & Brown 1986). Vocaliza un juiiip con inflexión ascendente, también una serie de notas prrriip guturales (Hilty & Brown 1986).
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID:
endemicAtomized
distribution

# ancillaryData

dataObject:

# additionalInformation: reference geoPoint: location: thumbnailURL: mediaURL description: subject source: audience bibliographicCitation: rightsHolder: rights: Equipo Coordinador del SiB Colombia license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: agent mimeType: dataType: identifier: distributionUnstructured: En bases de datos se encontró que: Esta especie se distribuye en: Canadá, Colombia, Costa Rica, El Salvador, Estados Unidos, Honduras, México (Arctos 2012). En bases de datos y monitoreos de fauna se encontró que: Esta especie se distribuye en: La Guajira (Cerrejón 2009-2010); Antioquia, Bogotá, Boyacá, Cauca, Chocó, Cundinamarca, La Guajira, Meta, Norte de Santander, Santander, Sucre (ICN 2004); Antioquia, Bolívar, Cesar, Chocó, Córdoba, Magdalena, Norte de Santander, Valle del Cauca (Collections MNH SI 2012). En todas las regiones del país hasta los 110 m (Hilty & Brown 1986). distributionAtomized stateProvince: Cauca ancillaryData locality: municipality: country: Colombia temporalCoverage endDate: startDate: distributionScope type: threatStatus

# ancillaryData

threatStatusUnstructured:

threatStatusAtomized
apendiceCITES: locality: municipality: county: stateProvince: country: Global 0: IUCN measurementValue: Preocupación menor (LC) relatedTo: measurementRemarks: measurementMethod:
measurementDeterminedBy
measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementType: IUCN measurementID:
territory
territoryUnstructured: areaOfOccupancy: extentOfOccurrence:
populationBiology
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Equipo Coordinador del SiB Colombia license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: populationBiologyUnstructured: La especie tiene un rango de distribución muy amplio y no se acerca al umbral de vulnerabilidad; la tendencia de la población es estable y su tamaño es extremadamente grande (BirdLife International 2012).
legislation
legislationUnstructured:
directThreats
directThreatsUnstructured:

### ancillaryData

dataObject:

additionalInformation:

geoPoint: location: thumbnailURL:

description: source:

bibliographicCitation:

rightsHolder:

rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

#### measurementOrFact

relatedTo:

measurement Remarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Mide unos 20 cm. Las partes dorsales ligeramente teñidas de oliva; rufo en la cola (parece toda rufa desde abajo); las primarias tienen márgenes rufos; la garganta y el pecho de color gris oscuro y el abdomen amarillo intenso (Hilty & Brown 1986).

#### associatedParty

role:

personnelldentifier: personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone: postalCode: country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

# ancillaryData

dataObject:

additionalInformation: Registros biológicos: En el portal de Global Biodiversity Information Facility se reporta que para Colombia hay 133 registros. GBIF

Créditos específicos: Ficha elaborada con el apoyo de Carbones del Cerrejón Limited.

Metadatos: GBIF

Otros recursos en Internet: Arctos, Avibase, AVoCet, Colecciones en Línea, EoL, iSpecies, IUCN Red List, SI NMNH

xeno-canto

#### reference

geoPoint: location:

thumbnailURL:

# mediaURL

description:

# subject

source:

#### audience

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

# agent

mimeType: dataType: identifier:

\_id

\_bsontype: ObjectID id: VÕöø< G™\0-