Reporte de Fichas para el usuario: edith gómez
Ficha ID: 56e1979783c45700544e3d19 Fecha de creación: 2016-03-10T15:49:50.282Z
annualCycles
annualCycleUnstructured:
habitats
habitatUnstructured:
commonNamesAtomized
ancillaryData
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
source:
audience
bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
agent
mimeType: dataType: identifier: usedBy:
usedIn
distributionUnstructured: endDate: startDate:
ancillaryData
type: language: Español name: Avispa creation_date: 2016-03-10T15:49:50.282Z

uses Management And Conservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint: location:

thumbnailURL: description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: Peciolo del tergo abdominal I casi tres veces más largo que ancho; tergo abdominal II más largo que ancho y punteado; especie negra; clípeo con dos pequeños dientes; ángulos del propodeo con fuertes punteaduras y ventralmente con carenas transversas suaves. briefDescription:

references

keywords: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti,Brachygastra lecheguana,Brachygastra propodealis,Brachygastra scutellaris,Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides. Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea

editors

address: publisher: accessed: websites: chapter: series: pages: 585 issue: year: 1978

institution:

authors

authors: Richards, O. W.

title: The social wasps of the Americas excluding the Vespinae

source: type: book

tags: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata. Agelaia hamiltoni. Agelaia testacea, Angiopolybia obidensis. Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi,Parachartergus weyrauchi,Polybia bistriata,Polybia catillifex,Polybia eberhardae,Polybia emaciata,Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea abstract:

identifiers

last_modified: created: group_id: profile_id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
replacementNameFor
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
sanctioned
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
laterHomonymOf
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
conservedAgainst
basedOn
microReference: source: datatype: identifier: source: datatype: identifier: note:

microReference:

ruleConsidered:
basionym
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
spellingCorrentionOf
typeNameEntity
source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:
typificacion
typeOfType:
lectotypeMicroReferenceVoucher
lectotypePublicationVoucher
voucherReference
simple: microReference: year:
publishedIn
simple: source: datatype: identifier:
specialAuthorship
authors
year
simple:
canonicalAuthorship

authors

year

simple:

simple: Richards, 1978

canonicalName

infraspecificEpithet: specificEpithet: infragenericEpithet:

linkType: ref:

uninomial: simple: Clypearia weyrauchi

rank: SPECIES

simple: Clypearia weyrauchi Richards, 1978

attributes

nomenclaturalCode: isAnamorphic:

id:

synonymsAtomized

hierarchy

ancillaryData

dataObject: additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rigthsHolder:

rights: license: title:

modified: 2016-03-10T15:49:30.272Z created: 2016-03-10T15:49:30.272Z

agent

homepage: role: email: phone: address: position: organisation: lastName: firstName: mimeType: dataType: identifier: parentTaxon: Clypearia higherClassification: infraspecificEpithet: specificEpithet: taxonRank: SPECIES subGenus: genus: Clypearia family: Vespidae order: Hymenoptera classHierarchy: Insecta phylum: Arthropoda kingdom: Animalia recommended: classification:
invasiveness
invasivenessUnstructured:
reproduction
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured: 0
dataObject: additionalInformation:

reference geoPoint: location: thumbnailURL: mediaURL description: subject audience rightsHolder: rights: license: modified: created: agent mimeType: dataType: identifier: relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy: TraitBank measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: Sexual measurementType: Tipo de Reproducción measurementID: molecularData molecularDataUnstructured: migratory migratoryUnstructured: lifeForm lifeFormUnstructured: lifeCycle lifeCycleUnstructured: interactions interactionsUnstructured:

feeding

dataObject:

additionalInformation:

0

```
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured:
0
strategy: Depredadores
type: Heterótrofos - Omnívoros
environmentalEnvelope
environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured:
distance:
structureDispersed:
type:
purpose:
behavior
dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
```

behaviorUnstructured: Al igual que las demás especies del género Clypearia, son insectos depredadores con hábitos eusociales. ancillaryData dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier: measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

agent

mimeType: dataType: identifier:

distributionUnstructured: Esta especie se ha registrado en el norte de Sudamérica (Colombia, Ecuador, Perú y Bolivia). Para el país se han reportado poblaciones de esta especie en la región de la Amazonia.

distributionAtomized

stateProvince: Putumayo

ancillaryData

locality: municipality: country: Colombia temporalCoverage

endDate: startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured: areaOfOccupancy: extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint: location: thumbnailURL: description:

source:

bibliographicCitation:	
rightsHolder:	
rights:	
license:	
modified:	
created:	
mimeType:	
dataType:	
identifier:	

measurementOrFact

relatedTo:

measurementRemarks: measurementMethod:

measurementDeterminedDate:

measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID:

abstract: Margen frontal lamelado del pronoto delgado; segmento abdominal I en forma de copa; dorso torácico no aplanado; metanoto con el área basal estrecha y horizontal, su área posterior es vertical abruptamente.

associatedParty

role:

personnelldentifier: personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone: postalCode: country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation: Información de tipos: El holotipo está depositado en el Museum of Comparative Zoology, University of Harvard Cambridge, EEUU (MCZ).

Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

reference

geoPoint: location: thumbnailURL:

mediaURL description: subject source: audience bibliographicCitation: rightsHolder: rights: license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: agent mimeType: dataType: identifier: _id _bsontype: ObjectID id: Vá——fÄW DãÑ•