

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c7bb78f0106c67230e740d

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

isbn: 13: 978-0-7922-5314-3

keywords: Butorides virescens,Empidonax alnorum,Empidonax traillii,Geothlypis philadelphia,Leiothlypis peregrina,Leucophaeus atricilla,Myiarchus crinitus,Patagioenas fasciata,Petrochelidon pyrrhonota,Setophaga castanea,Setophaga striata,Spiza americana

institution:

address:

publisher: the National Geographic Society

accessed:

websites:

chapter:

series:

pages: 503

issue:

year: 2006

0: Dunn, J

1: Alderfer, J

title: Field guide to the birds of North America. Fifth Edition.

source:

type: book

tags: Butorides virescens,Empidonax alnorum,Empidonax traillii,Geothlypis philadelphia,Leiothlypis peregrina,Leucophaeus atricilla,Myiarchus crinitus,Patagioenas fasciata,Petrochelidon pyrrhonota,Setophaga castanea,Setophaga striata,Spiza americana

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

isbn: 0-618-57424-7

keywords: Anas clypeata,Anas platyrhynchos,Ardea herodias,Bartramia longicauda,Buteo swainsoni,Cairina moschata,Calidris melanotos,Columbina talpacoti,Contopus cooperi,Contopus virens,Crotophaga ani,Dendrocygna autumnalis,Dolichonyx oryzivorus,Egretta caerulea,Egretta thula,Empidonax alnorum,Empidonax traillii,Falco columbarius,Falco sparverius,Geothlypis philadelphia,Hirundo rustica,Leiothlypis peregrina,Myiarchus crinitus,Myiodynastes luteiventris,Nomonyx dominicus,Nycticorax nycticorax,Pandion haliaetus,Parkesia noveboracensis,Petrochelidon pyrrhonota,Podilymbus podiceps,Protonotaria citrea,Sayornis nigricans

institution:

edition: Hillstar E

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 391

issue:

year: 2005

0: Kaufman, K

title: Guía de campo a las aves de Norteamérica

source:

type: book

tags: *Anas clypeata*, *Anas platyrhynchos*, *Ardea herodias*, *Bartramia longicauda*, *Buteo swainsoni*, *Cairina moschata*, *Calidris melanotos*, *Columbina talpacoti*, *Contopus cooperi*, *Contopus virens*, *Crotophaga ani*, *Dendrocygna autumnalis*, *Dolichonyx oryzivorus*, *Egretta caerulea*, *Egretta thula*, *Empidonax alnorum*, *Empidonax traillii*, *Falco columbarius*, *Falco sparverius*, *Geothlypis philadelphia*, *Hirundo rustica*, *Leiothlypis peregrina*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Pandion haliaetus*, *Parkesia noveboracensis*, *Petrochelidon pyrrhonota*, *Podilymbus podiceps*, *Protonotaria citrea*, *Sayornis nigricans*

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

2

keywords: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dubusia taeniata*, *Egretta caerulea*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Glaucidium jardinii*, *Grallaria ruficapilla*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Leiothlypis peregrina*, *Lesbia nuna*, *Lesbia victoriae*, *Leucophaeus atricilla*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Metallura tyrianthina*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Myiarchus crinitus*

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:  
series:  
pages: 276  
issue:  
year: 2000  
0: Asociación Bogotana de Ornitología  
title: Aves de la Sabana de Bogotá: guía de campo  
source:  
type: book  
tags: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dubusia taeniata*, *Egretta caerulea*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Glaucidium jardinii*, *Grallaria ruficapilla*, *Hemispingus superciliaris*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Leiothlypis peregrina*, *Lesbia nuna*, *Lesbia victoriae*, *Leucophaeus atricilla*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Metallura tyrianthina*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Myiarchus crinitus*  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

3

link: [http://awsassets.panda.org/downloads/migratorias{\\\_}aves{\\\_}42{\\\_}final.pdf](http://awsassets.panda.org/downloads/migratorias{\_}aves{\_}42{\_}final.pdf)  
keywords: *Myiarchus crinitus*  
0: Naranjo, L G  
1: Amaya, J D  
2: Eusse-González, D  
3: Cifuentes-Sarmiento, Y  
institution:  
address:  
publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia  
accessed:  
websites:  
chapter:  
series:  
pages: 708  
issue:  
volume: 1  
year: 2012  
0: Tello, Sofía  
title: *Myiarchus crinitus*  
source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves  
type: book\_section  
tags: *Myiarchus crinitus*

abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:

habitatUnstructured: El Atrapamoscas Copetón en sus zonas de reproducción habita en bosques de denso follaje y caducifolios (Dunn & Aldefer, 2006; Kaufman, 2005); en las áreas donde pasa el invierno se encuentra principalmente en los bordes de los bosques, parques, vegetación secundaria, frecuentemente buscando árboles y arbustos con frutos (Asociación Bogotana de Ornitología, 2000). También es común encontrar a esta especie en el dosel de árboles o arbustos atrapando insectos en vuelo; sus hábitats son borde de selva húmeda, bosques secundarios o áreas despejadas y matorrales; plantaciones de árboles y bosques de galería (Tello, 2012).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:  
usedIn  
distributionUnstructured:  
endDate:  
startDate:

[ancillaryData](#)

type:  
language: Inglés  
name: Great Crested Flycatcher

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

link: [http://awsassets.panda.org/downloads/migratorias{\\\_}aves{\\\_}42{\\\_}final.pdf](http://awsassets.panda.org/downloads/migratorias{\_}aves{\_}42{\_}final.pdf)

keywords: Myiarchus crinitus

0: Naranjo, L G

1: Amaya, J D

2: Eusse-González, D

3: Cifuentes-Sarmiento, Y

institution:

address:

publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia

accessed:

websites:

chapter:

series:

pages: 708

issue:

volume: 1

year: 2012

0: Tello, Sofía

title: Myiarchus crinitus

source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves

type: book\_section

tags: Myiarchus crinitus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: El Atrapamoscas Copetón mide 20 cm, plumaje de coloración oliva en la cabeza, espalda café oliva, alas con bordes rufos y cola rufa más acentuada. Garganta gris hasta el pecho, barriga amarilla brillante. Levanta la cresta cuando está alterado. El canto whee eep es diagnóstico (Tello, 2012).

briefDescription:

## references

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon

assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dubusia taeniata,Egretta caerulea,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreiventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Gallinula melanops,Geothlypis philadelphia,Glaucidium jardinii,Grallaria ruficapilla,Hemispingus supercilialis,Hemispingus verticalis,Henicorhina leucophrys,Hirundo rustica,Icterus chrysater,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Leiothlypis peregrina,Lesbia nuna,Lesbia victoriae,Leucophaeus atricilla,Margarornis squamiger,Mecocerculus leucophrys,Megascops choliba,Melanerpes rubricapillus,Metallura tyrianthina,Mimus gilvus,Mniotilta varia,Molothrus bonariensis,Myiarchus crinitus

## editors

editors: Stiles, F G  
editors: Bohórquez, C I  
editors: Cadena, C D  
editors: De la Zerda, S  
editors: Hernández, M  
editors: Roselli, L  
editors: Kelsey, M  
editors: Valencia, I D  
editors: Knapp, D  
institution:  
edition: Primera Ed  
address:  
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca  
accessed:  
websites:  
chapter:  
series:  
pages: 276  
issue:  
year: 2000

## authors

authors: Asociación Bogotana de Ornitología  
title: Aves de la Sabana de Bogotá: guía de campo  
source:  
type: book  
tags: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dubusia taeniata,Egretta caerulea,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreiventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Gallinula melanops,Geothlypis philadelphia,Glaucidium jardinii,Grallaria ruficapilla,Hemispingus supercilialis,Hemispingus verticalis,Henicorhina

leucophrys,Hirundo rustica,Icterus chrysater,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Leiothlypis peregrina,Lesbia nuna,Lesbia victoriae,Leucophaeus atricilla,Margarornis squamiger,Mecocerculus leucophrys,Megascops choliba,Melanerpes rubricapillus,Metallura tyrianthina,Mimus gilvus,Mniotilta varia,Molothrus bonariensis,Myiarchus crinitus

abstract:

identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

basionym

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:

datatype:

identifier:

source:

datatype:

identifier:

source:

datatype:

identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)



lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:

publishedIn

simple:  
source:  
datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: (Linnaeus, 1758)

canonicalName

infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: Myiarchus crinitus  
rank: SPECIES  
simple: Myiarchus crinitus (Linnaeus, 1758)

attributes

nomenclaturalCode:  
isAnamorphic:  
id:

synonymsAtomized

ancillaryData

synonymStatus:  
synonymName  
anyTwo:

anyOne

providerLink:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:

ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typeVoucherEntity

typeOfType:  
simple:  
microReference:  
year:  
source:  
datatype:  
identifier:

combinationAuthorship

basionymAuthorship

simple:  
authorship

simple:  
simple: Bangs, 1898

epithet

infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:

genus

linkType:  
ref:  
uninomial:  
simple: Myiarchus crinitus boreus  
rank: SUBSPECIES  
simple: Myiarchus crinitus boreus Bangs, 1898  
nomenclaturalCode:  
isAnamorphic:  
id:

[hierarchy](#)

[ancillaryData](#)

dataObject:  
additionalInformation:

[reference](#)

geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:  
rightsHolder:  
rights:  
license:  
title:  
modified: 2016-02-22T14:05:27.799Z  
created: 2016-02-22T14:05:27.799Z

[agent](#)

homepage:  
role:  
email:  
phone:  
address:  
position:

organisation:  
lastName:  
firstName:  
mimeType:  
dataType:  
identifier:  
parentTaxon: Myiarchus  
higherClassification:  
infraspecificEpithet:  
specificEpithet:  
taxonRank: SPECIES  
subGenus:  
genus: Myiarchus  
family: Tyrannidae  
order: Passeriformes  
classHierarchy: Aves  
phylum: Chordata  
kingdom: Animalia  
recommended:  
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:  
additionalInformation:

0

isbn: 0-618-57424-7

keywords: Anas clypeata, Anas platyrhynchos, Ardea herodias, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Columbina talpacoti, Contopus cooperi, Contopus virens, Crotophaga ani, Dendrocygna autumnalis, Dolichonyx oryzivorus, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Falco sparverius, Geothlypis philadelphia, Hirundo rustica, Leiothlypis peregrina, Myiarchus crinitus, Myiodynastes luteiventris, Nomonyx dominicus, Nycticorax nycticorax, Pandion haliaetus, Parkesia noveboracensis, Petrochelidon pyrrhonota, Podilymbus podiceps, Protonotaria citrea, Sayornis nigricans

institution:

edition: Hillstar E

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 391

issue:

year: 2005

0: Kaufman, K

title: Guía de campo a las aves de Norteamérica

source:

type: book

tags: Anas clypeata, Anas platyrhynchos, Ardea herodias, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Columbina talpacoti, Contopus cooperi, Contopus virens, Crotophaga ani, Dendrocygna

autumnalis,Dolichonyx oryzivorus,Egretta caerulea,Egretta thula,Empidonax alnorum,Empidonax traillii,Falco columbarius,Falco sparverius,Geothlypis philadelphia,Hirundo rustica,Leiothlypis peregrina,Myiarchus crinitus,Myiodynastes luteiventris,Nomonyx dominicus,Nycticorax nycticorax,Pandion haliaetus,Parkesia noveboracensis,Petrochelidon pyrrhonota,Podilymbus podiceps,Protonotaria citrea,Sayornis nigricans

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

link: <http://eol.org/pages/916843/overview>

keywords: Myiarchus crinitus

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

0: EOL

title: Myiarchus crinitus

source: Encyclopedia of life

type: miscellany

tags: Myiarchus crinitus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

reproductionUnstructured: Es una especie monógama. Durante la época reproductiva los machos defienden fuertemente su territorio, que establecen a inicios de mayo. La hembra construye el nido con diferentes materiales como hojas, plumas, hierba, corteza, entre otros (EOL, n.d.), además tiene el extraño hábito de añadir en un pedazo muda de piel de serpiente (Kaufman, 2005). Las hembras ponen entre cuatro y ocho huevos con manchas marrón y púrpura; la incubación dura 15 días aproximadamente y entre los 13 y 15 días los juveniles abandonan el nido (EOL, n.d.).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

0

link: <http://eol.org/pages/916843/overview>

keywords: Myiarchus crinitus

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

0: EOL

title: Myiarchus crinitus

source: Encyclopedia of life

type: miscellany

tags: Myiarchus crinitus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

feedingUnstructured: El Atrapamoscas Copetón es una especie insectívora, que ocasionalmente consume frutos (EOL, n.d.).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Metallura tyrianthina, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Myiarchus crinitus

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca



accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dubusia taeniata*, *Egretta caerulea*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Glaucidium jadinii*, *Grallaria ruficapilla*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Leiothlypis peregrina*, *Lesbia nuna*, *Lesbia victoriae*, *Leucophaeus atricilla*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Metallura tyrianthina*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Myiarchus crinitus*

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Aglaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigristrois*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Cistothorus platensis*, *Coccyzus pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta caerulea*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jadinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura*

andicola, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiostyris alpestris, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla, Machetornis rixosa, Margarornis

squamiger, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxypogon guerinii*, *Oxyura jamaicensis*, *Pachyramphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systemellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

2

isbn: 0-618-57424-7

keywords: *Anas clypeata*, *Anas platyrhynchos*, *Ardea herodias*, *Bartramia longicauda*, *Buteo swainsoni*, *Cairina moschata*, *Calidris melanotos*, *Columbina talpacoti*, *Contopus cooperi*, *Contopus virens*, *Crotophaga ani*, *Dendrocygna autumnalis*, *Dolichonyx oryzivorus*, *Egretta caerulea*, *Egretta thula*, *Empidonax alnorum*, *Empidonax traillii*, *Falco columbarius*, *Falco sparverius*, *Geothlypis philadelphia*, *Hirundo rustica*, *Leiothlypis peregrina*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Pandion haliaetus*, *Parkesia noveboracensis*, *Petrochelidon pyrrhonota*, *Podilymbus podiceps*, *Protonotaria citrea*, *Sayornis nigricans*

institution:

edition: Hillstar E

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 391

issue:

year: 2005

0: Kaufman, K

title: Guía de campo a las aves de Norteamérica

source:

type: book

tags: *Anas clypeata*, *Anas platyrhynchos*, *Ardea herodias*, *Bartramia longicauda*, *Buteo swainsoni*, *Cairina moschata*, *Calidris melanotos*, *Columbina talpacoti*, *Contopus cooperi*, *Contopus virens*, *Crotophaga ani*, *Dendrocygna autumnalis*, *Dolichonyx oryzivorus*, *Egretta caerulea*, *Egretta thula*, *Empidonax alnorum*, *Empidonax traillii*, *Falco columbarius*, *Falco sparverius*, *Geothlypis philadelphia*, *Hirundo rustica*, *Leiothlypis peregrina*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Pandion haliaetus*, *Parkesia noveboracensis*, *Petrochelidon pyrrhonota*, *Podilymbus podiceps*, *Protonotaria citrea*, *Sayornis nigricans*

abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

3

isbn: 978-958-44-1101-3  
keywords: Buteo swainsoni, Coccycua pumila, Coccyzus melacoryphus, Contopus virens, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Dolichonyx oryzivorus, Egretta thula, Empidonax traillii, Falco sparverius, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Machetornis rixosa, Milvago chimachima, Mimus gilvus, Myiarchus crinitus, Nomonyx dominicus, Nycticorax nycticorax, Pardirallus maculatus, Phaetusa simplex, Phalacrocorax brasilianus, Phimosus infuscatus, Pitangus sulphuratus, Pygochelidon cyanoleuca, Sayornis nigricans, Setophaga striata  
institution:  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
pages: 301  
issue:  
year: 2007  
0: McNish, T  
title: Las aves de los Llanos de la Orinoquia

source:  
type: book  
tags: Buteo swainsoni, Coccycua pumila, Coccyzus melacoryphus, Contopus virens, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Dolichonyx oryzivorus, Egretta thula, Empidonax traillii, Falco sparverius, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Machetornis rixosa, Milvago chimachima, Mimus gilvus, Myiarchus crinitus, Nomonyx dominicus, Nycticorax nycticorax, Pardirallus maculatus, Phaetusa simplex, Phalacrocorax brasilianus, Phimosus infuscatus, Pitangus sulphuratus, Pygochelidon cyanoleuca, Sayornis nigricans, Setophaga striata  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

4

link: [http://awsassets.panda.org/downloads/migratorias{\\\_}aves{\\\_}42{\\\_}final.pdf](http://awsassets.panda.org/downloads/migratorias{\_}aves{\_}42{\_}final.pdf)  
keywords: Myiarchus crinitus  
0: Naranjo, L G  
1: Amaya, J D  
2: Eusse-González, D  
3: Cifuentes-Sarmiento, Y  
institution:  
address:  
publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia  
accessed:  
websites:  
chapter:

series:  
pages: 708  
issue:  
volume: 1  
year: 2012  
0: Tello, Sofía  
title: Myiarchus crinitus  
source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves  
type: book\_section  
tags: Myiarchus crinitus  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

5

isbn: 978-0-679-45122-8  
keywords: Anas clypeata,Ardea herodias,Buteo swainsoni,Cairina moschata,Crotophaga ani,Dendrocygna autumnalis,Dendrocygna viduata,Egretta caerulea,Egretta thula,Empidonax alnorum,Geothlypis philadelphia,Ixobrychus exilis,Leiothlypis peregrina,Myiarchus crinitus,Myiodynastes luteiventris,Nomonyx dominicus,Parkesia noveboracensis,Petrochelidon pyrrhonota,Pheucticus ludovicianus,Protonotaria citrea,Sayornis nigricans  
institution:  
edition: 1 st ed.  
address:  
publisher: National Audubon Society  
accessed:  
websites:  
chapter:  
series:  
pages: 545  
issue:  
year: 2000  
0: Sibley, D A  
title: The SIBLEY Guide to Birds  
source:  
type: book  
tags: Anas clypeata,Ardea herodias,Buteo swainsoni,Cairina moschata,Crotophaga ani,Dendrocygna autumnalis,Dendrocygna viduata,Egretta caerulea,Egretta thula,Empidonax alnorum,Geothlypis philadelphia,Ixobrychus exilis,Leiothlypis peregrina,Myiarchus crinitus,Myiodynastes luteiventris,Nomonyx dominicus,Parkesia noveboracensis,Petrochelidon pyrrhonota,Pheucticus ludovicianus,Protonotaria citrea,Sayornis nigricans  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Con frecuencia se le ve posado alto en el denso follaje (Kaufman, 2005); en árboles y arbustos en fruta (Hilty & Brown, 1986). Hace vuelos cortos para sacar insectos o frutos del follaje; no hace salidas largas al aire como lo hace un Tyrannus (Asociación Bogotana de Ornitología, 2000). Es poco común, es más escuchado que observado; desde dentro del dosel del bosque emite su canto característico; además del canto, su mayor tamaño y el rufo en las alas son características que facilitan su identificación (McNish, 2007). Los Atrapamoscas Copetones migran desde Canadá a finales de agosto – septiembre y a finales de septiembre, llegan a Costa Rica, Panamá y Colombia entre septiembre y octubre. De regreso salen de Colombia, Panamá y Costa Rica entre abril y mayo y los primeros sitios que llegan son Texas, Louisiana y Florida Central (Tello, 2012). Un característico reclamo wiya y brrrk brrrk brrrk! (Kaufman, 2005; Sibley, 2000). En las zonas de invernada emite un juiip con inflexión ascendente; también una serie de notas prrrriip guturales (más frecuente en las zonas templadas de Norteamérica) (Hilty & Brown, 1986).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:

additionalInformation:

## reference

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccyzua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiostyris alpestris, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus cinerascens, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

## editors

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

## authors

authors: Hilty, Steven

authors: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccyzua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliosus, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiostyris alpestris, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla, Mactornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus cinerascens, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne subis, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicrodon microrhynchus, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaidura macroura, Zonotrichia capensis

abstract:

## identifiers

last\_modified:

created:

group\_id:

profile\_id:



id:  
geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

[agent](#)

mimeType:  
dataType:  
identifier:

distributionUnstructured: El Atrapamoscas Copetón se distribuye desde el este de Alberta hasta el sur de Manitoba, siguiendo en el sur de Ontario y sur de Quebec además de Nueva Brunswick y Nueva Escocia. En Estados Unidos la distribución incluye la Florida y los cayos de la costa Atlántica y los estados de la región central. También puede reproducirse en el noreste de Coahuila, México. Inverna principalmente desde el sur de México (Veracruz, Oaxaca y península de Yucatán) a través de Centroamérica hasta el este de Panamá y el norte de Sudamérica en Colombia, oeste de Venezuela y algunos pocos registros en el Ecuador. En México es muy común en la vertiente Atlántica pero ausente a altas elevaciones desde Chiapas hasta el sur. En Costa Rica es común en la vertiente del Pacífico. En Colombia inverna en la cordillera Occidental, pero hay algunos registros para la cordillera Oriental. De los pocos registros para Venezuela se tienen localidades de Zulia, Mérida y el río Orinoco y en Ecuador muy pocos pero recientes registros en la provincia de Napo. Se ha reportado en muy pocas oportunidades en Cuba y Bermudas (Tello, 2012). En Colombia comprende la zona Andina con mayor concentración en el occidente (zona pacífica) y el norte del país. Posiblemente en el oriente. Específicamente se ha registrado en los departamentos de Antioquia, Boyacá, Cauca, Chocó, Caldas, Cundinamarca, Guajira, Quindío, Risaralda, San Andrés y Providencia y Valle del Cauca (Tello, 2012). Presente hasta los 1100 msnm (Hilty & Brown, 1986).

[distributionAtomized](#)

stateProvince: Córdoba

[ancillaryData](#)

locality:  
municipality:  
country: Colombia

temporalCoverage

endDate:  
startDate:

distributionScope

type:

## threatStatus

## ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

## measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

0

link: <http://maps.iucnredlist.org/map.html?id=22700433>

keywords: Myiarchus crinitus

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: BirdLife International and NatureServe

title: Bird Species Distribution Maps of the World. 2012. Myiarchus crinitus. The IUCN Red List of Threatened Species. Version 2014.3

source:

type: miscellany

tags: Myiarchus crinitus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

populationBiologyUnstructured: Tiene un alto rango de distribución mundial, la tendencia de la población es estable y su categoría de amenaza está en Preocupación Menor (BirdLife International and NatureServe, 2014).

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: El Atrapamoscas Copetón mide 20 cm; es poco común; es más escuchado que observado; desde dentro del dosel del bosque emite su canto característico (McNish, 2007). Se parece al Atrapamoscas Crestado (*Myiarchus tyrannulus*), pero es más oscuro; partes superiores ligeramente teñidas de oliva (coronilla no claramente café); más rufo en la cola (parece toda rufa desde abajo); primarias con márgenes rufos; garganta y pecho gris más oscuros y abdomen amarillo más intenso (no amarillo blanquecino pálido). La voz es simplemente diagnóstica (Hilty & Brown, 1986). Es muy común encontrar a esta especie en el dosel de árboles o arbustos atrapando insectos en vuelo; sus hábitats son borde de selva húmeda, bosques secundarios o áreas despejadas y matorrales; plantaciones de árboles y bosques de galería (Tello, 2012). Tiene un alto rango de distribución mundial; la tendencia de la población es estable y su categoría de amenaza está en Preocupación Menor (BirdLife International and NatureServe, 2014).

## associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

## ancillaryData

dataObject:

additionalInformation:

## reference

geoPoint:

location:

thumbnailURL:

mediaURL: <http://avibase.bsc-eoc.org/species.jsp?lang=ES&avibaseid=8D6D5F2BEA07A0CF>

description:

## subject

source: <http://avibase.bsc-eoc.org>

## audience

bibliographicCitation:

rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

## agent

contentType:  
dataType:  
identifier:

\_id

\_bsonType: ObjectId  
id: VÇ»xđ lg#çM