Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e7228383c45700544e3f05

Fecha de creación: 2016-03-14T20:43:55.340Z

creation_date: 2016-03-14T20:43:55.340Z usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject: additionalInformation:

additionaliniormation

0

keywords: Agamopus lampros, Ateuchus aeneomicans, Canthidium aurifex, Canthidium centrale, Canthidium elegantulum, Canthidium perceptibile, Canthon deyrollei, Canthon lituratus, Copris incertus, Dichotomius agenor, Onthophagus landolti landolti, Onthophagus transisthmius, Pseudocanthon perplexus, Uroxys bidentis, Uroxys depressifrons, Uroxys gatunensis, Uroxys gorgon, Uroxys micros

institution: address: publisher: accessed: websites: chapter: series:

pages: 1--204

issue: volume: 18 year: 1981 0: Howden, H. 1: Young, O.P.

title: Panamanian Scarabaeinae: taxonomy, distribution, and habits (Coleoptera, Scarabaeidae)[New taxa]

source: Contributions of the American Entomological Institute (USA)

type: article

tags: Agamopus lampros,Ateuchus aeneomicans,Canthidium aurifex,Canthidium centrale,Canthidium elegantulum,Canthidium perceptibile,Canthon deyrollei,Canthon lituratus,Copris incertus,Dichotomius agenor,Onthophagus landolti landolti,Onthophagus transisthmius,Pseudocanthon perplexus,Uroxys bidentis,Uroxys depressifrons,Uroxys gatunensis,Uroxys gorgon,Uroxys micros

abstract: last_modified: created: group_id: profile_id: id:

geoPoint: location:

thumbnailURL: description:

source:

bibliographicCitation:

rightsHolder:

rights: Claudia Alejandra Medina

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: Región fronto clipeal con tres tubérculos. Superficie no carinada entre ellos; longitud 4.5

mm. (Howden & Young, 1981).

briefDescription:

references

keywords: Canthidium angusticeps, Canthidium aurifex, Canthidium basipunctatum, Canthidium bicolor, Canthidium centrale, Canthidium convexifrons, Canthidium cupreum, Canthidium deplanatum, Canthidium elegantulum, Canthidium escalerai, Canthidium euchalceum, Canthidium funebre, Canthidium gerstaeckeri, Canthidium lebasi, Canthidium onitoides, Canthidium perceptibile, Canthidium rufinum, Canthidium splendidum, Canthidium steinheili

editors

institution: address: publisher: accessed: websites: chapter: series: pages: 19--40

issue: year: 1986

authors

authors: Mart\'inez, A. authors: Halffter, G.

title: Situación del género Canthidium Erichson (Coleoptera: Scarabaeidae: Scarabaeinae)

source: Acta Zoológica Mexicana

type: article

tags: Canthidium angusticeps,Canthidium aurifex,Canthidium basipunctatum,Canthidium bicolor,Canthidium centrale,Canthidium convexifrons,Canthidium cupreum,Canthidium deplanatum,Canthidium elegantulum,Canthidium escalerai,Canthidium euchalceum,Canthidium funebre,Canthidium gerstaeckeri,Canthidium lebasi,Canthidium onitoides,Canthidium perceptibile,Canthidium rufinum,Canthidium splendidum,Canthidium steinheili

abstract:

identifiers

last_modified: created: group_id: profile id:

id.

taxonRecordName

ancillaryData

providerSpecificData anyTwo: providerLink: publicationStatus microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: replacementNameFor microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: sanctioned microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: laterHomonymOf microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: conservedAgainst basedOn microReference:

source: datatype:

identifier: source: datatype: identifier: note: ruleConsidered:
basionym
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
spellingCorrentionOf
typeNameEntity
source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:
typificacion
typeOfType:
lectotypeMicroReferenceVoucher
lectotypePublicationVoucher
voucherReference
simple: microReference: year:
publishedIn
simple: source: datatype: identifier:
specialAuthorship
authors

year

simple:

canonicalAuthorship

authors

year

simple:

simple: Bates, 1887

canonicalName

infraspecificEpithet: specificEpithet: infragenericEpithet:

linkType: ref: uninomial:

simple: Canthidium aurifex

rank: SPECIES

simple: Canthidium aurifex Bates, 1887

attributes

nomenclaturalCode: isAnamorphic:

id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rigthsHolder:

rights: license: title:

modified: 2016-03-14T20:43:35.327Z created: 2016-03-14T20:43:35.327Z

agent

homepage:

role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:

parentTaxon: Canthidium higherClassification: infraspecificEpithet: specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Canthidium family: Scarabaeidae order: Coleoptera classHierarchy: Insecta phylum: Arthropoda kingdom: Animalia recommended: classification:

commonNameAtomized

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Claudia Alejandra Medina

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: agent mimeType: dataType: identifier: usedBy: usedIn distributionUnstructured: endDate: startDate: ancillaryData type: language: Español name: Escarabajo coprófago invasiveness invasivenessUnstructured: reproduction dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Claudia Alejandra Medina license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured: dataObject: additionalInformation: reference geoPoint: location: thumbnailURL: mediaURL

description:
subject
audience
rightsHolder: rights: license: modified: created:
agent
mimeType: dataType: identifier: relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy: TraitBank measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: Sexual measurementType: Tipo de Reproducción measurementID:
molecularData
molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions
interactionsUnstructured:
feeding
feedingUnstructured:
environmentalEnvelope
environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured: distance:

structureDispersed: type: purpose:
behavior
behaviorUnstructured:
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID:
annualCycle
annualCycleUnstructured:
habitat
habitatUnstructured:
endemicAtomized
distribution
ancillaryData
dataObject: additionalInformation:
reference

keywords: Canthidium angusticeps,Canthidium aurifex,Canthidium basipunctatum,Canthidium bicolor,Canthidium centrale,Canthidium convexifrons,Canthidium cupreum,Canthidium deplanatum,Canthidium elegantulum,Canthidium

escalerai, Canthidium euchalceum, Canthidium funebre, Canthidium gerstaeckeri, Canthidium lebasi, Canthidium onitoides, Canthidium perceptibile, Canthidium rufinum, Canthidium splendidum, Canthidium steinheili

editors

institution: address: publisher: accessed: websites: chapter: series:

pages: 19--40

issue: year: 1986

authors

authors: Mart\'inez, A. authors: Halffter, G.

title: Situación del género Canthidium Erichson (Coleoptera: Scarabaeidae: Scarabaeinae)

source: Acta Zoológica Mexicana

type: article

tags: Canthidium angusticeps, Canthidium aurifex, Canthidium basipunctatum, Canthidium bicolor, Canthidium centrale, Canthidium convexifrons, Canthidium cupreum, Canthidium deplanatum, Canthidium elegantulum, Canthidium escalerai, Canthidium euchalceum, Canthidium funebre, Canthidium gerstaeckeri, Canthidium lebasi, Canthidium onitoides, Canthidium perceptibile, Canthidium rufinum, Canthidium splendidum, Canthidium steinheili abstract:

identifiers

last_modified:

created:

group_id:

profile id:

id:

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Claudia Alejandra Medina

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

agent

mimeType:

dataType:

identifier:

distributionUnstructured: Esta especie ha sido registrada en Colombia, Costa Rica, Ecuador y Panamá (Martínez & Halffter, 1986; Medina et al., 2001). En el país se distribuye en los departamentos de Antioquia, Chocó y Vaupés (Medina et al., 2001).

distributionAtomized

stateProvince: Antioquia

ancillaryData

locality: municipality: country: Colombia

temporalCoverage

endDate: startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured: areaOfOccupancy: extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint: location: thumbnailURL: description:

source:

bibliographicCitation:

rightsHolder:

rights:
license:
modified:
created:
mimeType:

dataType:

identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID: abstract: Canthidium aurifex es una especie de la familia Scarabaeidae que se distribuye en Sudamérica y Centroamérica (Martínez & Halffter, 1986; Medina et al., 2001). Estos insectos alcanzan longitudes de 4.5 mm y presentan una región fronto clipeal con tres tubérculos (Howden & Young, 1981).
associatedParty
role: personnelldentifier: personnelDirectory: homepage: email: enataliagomezm@gmail.com phone: postalCode: country: Colombia state: Bogotá Distrito Capital city: Bogotá D.C. address: organisation: position: lastName: Gómez Melendro firstName: Edith Natalia
ancillaryData
dataObject: additionalInformation: Créditos específicos: La elaboración de estas fichas recibió el apoyo de la Organización de los Estados Américanos (OEA) y la Red Interamericana de Información sobre Biodiversidad (IABIN). Se agradece a la Colección Entomológica del Instituto Humboldt IAVH por facilitar información que contribuyó a la elaboración de las fichas y al Sistema de Información en Biodiversidad de Colombia (SIB) por el apoyo técnico. Imagen: Canthidium aurifex.jpg
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
source:
audience

bibliographicCitation:
rightsHolder:
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
agent
mimeType:
dataType:
identifier:
_id

_bsontype: ObjectID id: $V c f \ddot{f} W D \tilde{a} \tilde{o}$