

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d5d0443c16479905cba8fc

Fecha de creación: 2016-03-01T17:24:22.957Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Es común encontrar a esta especie en matorrales y árboles cercanos ríos, lagos, pantanos, lagunas y en manglares (Hilty & Brown 1986). En La Guajira habita en el bosque deciduo tropical, de galería y secundario (Cerrejón 2009-2010).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Español

name: Garrapatero mayor

creation_date: 2016-03-01T17:24:22.957Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: Mide unos 46 cm. Tiene ojos blancos prominentes, el pico negro comprimido con culmen arqueado en la porción basal. Su cuerpo es delgado de color negro azul lustroso, con lustre verdoso en las alas y púrpura en la cola que es larga y redondeada (Hilty & Brown 1986).

briefDescription:

[references](#)

taxonRecordName

[ancillaryData](#)

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

replacementNameFor

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:

identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: Gmelin, 1788

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:

ref:
uninomial:
simple: Crotophaga major
rank: SPECIES
simple: Crotophaga major Gmelin, 1788

attributes

nomenclaturalCode:

isAnamorphic:

id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-01T17:24:02.955Z

created: 2016-03-01T17:24:02.955Z

[agent](#)

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataType:

identifier:

parentTaxon: Crotophaga

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Crotophaga
family: Cuculidae
order: Cuculiformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

feedingUnstructured: Su dieta se compone de insectos y variedad de presas pequeñas (Hilty & Brown 1986). En La Guajira esta especie forrajea desde el suelo hasta el dosel y se alimenta de insectos (Cerrejón 2009-2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Gregario, se ha observado en grupos de 4 a 100 o más (Hilty & Brown 1986). Esta es un ave ruidosa que genera sonidos electrónicos extraños como un gorgoreo cro-coro que repite rápidamente, también vocaliza gruñidos, siseos y zumbidos (Hilty & Brown 1986).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:
distributionUnstructured: En bases de datos se encontró que:š
Esta especie se distribuye en: Argentina, Brasil, Colombia, Ecuador, Guyana, Panamá, Paraguay, Perú, Venezuela (Collections MNH SI 2012); Brasil, Colombia, Suriname (Arctos 2012). En bases de datos y monitoreos de fauna se encontró que:
Esta especie se distribuye en: La Guajira (Cerrejón 2009-2010); Amazonas, Atlántico, Bogotá, Bolívar, Caquetá, Casanare, Chocó, Cundinamarca, Huila, Magdalena, Santander, Tolima, Vaupés (ICN 2004); Bolívar, Caquetá, Cesar, Chocó, Magdalena, Sucre (Collections MNH SI 2012); Huila (Arctos 2012). En las regiones Amazonía, Andina, Caribe, Orinoquía y Pacífica del país.

distributionAtomized

stateProvince: Tolima

ancillaryData

locality:
municipality:
country: Colombia

temporalCoverage

endDate:
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:
municipality:
county:
stateProvince:
country: Global
0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

abstract: Mide unos 46 cm. Tiene ojos blancos prominentes, el pico negro comprimido con culmen arqueado en la porción basal. Su cuerpo es delgado de color negro azul lustroso, con lustre verdoso en las alas y púrpura en la cola que es larga y redondeada (Hilty & Brown 1986).

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Registros biológicos: Portal de datos GBIF
Metadatos: Portal de datos GBIF
Créditos específicos: "f-6† VÆ &÷& F 6öâ VÂ ÷—ò FR 6 &&öæW2 FVÂ 6W'&V~6â Æ—Ö—FVBâ
Otros recursos en Internet: Arctos, Avibase, AVoCet, Colecciones en Línea, EoL, iSpecies, IUCN Red List, SI NMNH
xeno-canto

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:
subject

source:
audience

bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id
_bsontype: ObjectID
id: VÖÐD< G™\0•Ä