

Reporte de Fichas para el usuario: Edgar Herrera

Ficha ID: 56e7985783c45700544e41e1

Fecha de creación: 2016-03-15T05:07:04.382Z

creation_date: 2016-03-15T05:07:04.382Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

keywords: Accipiter collaris, Accipiter cooperii, Accipiter poliogaster, Accipiter superciliosus, Amazona ochrocephala, Andigena hypoglaucha, Aramus guarauna, Aratinga pertinax, Ardea alba, Ardea cocoi, Ardea herodias, Arremonops conirostris, Arremonops tocuyensis, Atalotriccus pilaris, Attila spadiceus (Gmelin 1789), Basileuterus flaveolus (S. F. Baird 1865), Bubo virginianus (Gmelin 1788), Busarellus nigricollis, Buteogallus anthracinus, Buteogallus meridionalis, Buteogallus schistaceus, Buteogallus solitarius, Buteogallus urubitinga, Cacicus cela, Cairina moschata, Campylorhamphus trochilirostris (Lichtenstein 1820), Campylorhynchus griseus (Swainson 1838), Campylorhynchus nuchalis (Cabanis 1847), Caprimulgus cayennensis (Gmelin 1789), Cardinalis phoeniceus Bonaparte 1838, Carduelis psaltria (Say 1823), Cathartes aura, Cathartes burrovianus, Cathartes melambrotus, Certhiaxis cinnamomeus (Gmelin 1788), Ceryle torquata, Chlorostilbon gibsoni (Fraser 1840), Chondrohierax uncinatus, Chordeiles acutipennis (Hermann 1783), Chrysomus icterocephalus, Circus buffoni, Circus cinereus, Circus cyaneus, Coccyzus pumila, Columbina minuta (Linnaeus 1766), Conirostrum leucogenys (Lafresnaye 1852), Coragyps atratus, Coryphospingus pileatus (Wied-Neuwied 1821), Crax alector, Crotophaga major, Crotophaga sulcirostris Swainson 1827, Cryptoleucopteryx plumbea, Cyanocorax affinis (Pelzelin 1856), Cyclarhis gujanensis (J. F. Gmelin 1789), Dryocopus lineatus (Linnaeus 1766), Elaenia chiriquensis Lawrence 1865, Elanoides forficatus, Elanus leucurus, Falco deiroleucus, Fluvicola pica (Boddaert 1783), Forpus passerinus (Linnaeus 1758), Forpus xanthopterygius (Spix 1824), Furnarius leucopus Swainson 1838, Galbula ruficauda Cuvier 1816, Gallinula chloropus (Linnaeus 1758), Gampsonyx swainsonii, Geranospiza caerulescens, Glaucidium brasilianum (Gmelin 1788), Harpagus bidentatus, Harpia harpyja, Helicolestes hamatus, Hemitriccus margaritaceiventer (Orbigny & Lafresna, Himantopus mexicanus (Statius Muller 1776), Hylophilus flavipes Lafresnaye 1845, Hypnelus ruficollis (Wagler 1829), Icterus auricapillus Cassin 1848, Icterus galbula (Linnaeus 1758), Icterus nigrogularis (Hahn 1819), Ictinia mississippiensis, Ictinia plumbea, Inezia subflava (P. L. Sclater & Salvin 1873), Inezia tenuirostris (Cory 1913), Jabiru mycteria, Jacana jacana, Lepidopyga goudoti (Bourcier 1843), Leptotila verreauxi (Bonaparte 1855), Leucopternis melanops, Leucopternis semiplumbea, Machetornis rixosa, Megaceryle alcyon (Linnaeus 1758), Megaceryle torquata (Linnaeus 1766), Megarynchus pitangua (Linnaeus 1766), Megascops choliba, Melanerpes rubricapillus, Micrastur plumbeus, Mitu tomentosa, Molothrus bonariensis, Morphnarchus princeps, Morphnus guianensis, Myiarchus crinitus, Myiarchus venezuelensis Lawrence 1865, Myiopagis gaimardii (Orbigny 1840), Myiozetetes similis (Spix 1825), Myospiza humeralis (Bosc 1792), Nemosia pileata (Boddaert 1783), Neochen jubata, Opisthocomus hoazin, Oroaetus isidori, Ortalis ruficauda Jardine 1847, Pandion haliaetus, Parabuteo unicinctus, Patagioenas corensis (Jacquin 1784), Phaeomyias murina (Spix 1825), Pheucticus ludovicianus, Piculus chrysochloros (Vieillot 1818), Picumnus cinnamomeus Wagler 1829, Pilherodius pileatus (Boddaert 1783), Pionus sordidus (Linnaeus 1758), Pitangus lictor (Lichtenstein 1823), Pitangus sulphuratus, Polioptila plumbea (J. F. Gmelin 1788), Pseudastur albicollis, Pulsatrix perspicillata (Latham 1790), Pyrocephalus rubinus, Quiscalus lugubris, Quiscalus mexicanus (Gmelin 1788), Rostrhamus sociabilis, Sakesphorus canadensis (Linnaeus 1766), Saltator coerulescens Vieillot

1817, *Sarcoramphus papa*, *Sarkidiornis melanotos* (Pennant 1769), *Spiza americana*, *Spizaetus tyrannus*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx ruficollis* (Vieillot 1817), *Sterna supercilialis*, *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tachycineta bicolor* (Vieillot 1808), *Tapera naevia* (Linnaeus 1766), *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Zenaida auriculata*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

issue:

year: 1986

0: Hilty, S

1: Brown, W. L.

title: A Guide to the Birds of Colombia

source:

type: book

tags: *Accipiter collaris*, *Accipiter cooperii*, *Accipiter poliogaster*, *Accipiter superciliosus*, *Amazona ochrocephala*, *Andigena hypoglaucha*, *Aramus guarauna*, *Aratinga pertinax*, *Ardea alba*, *Ardea cocoi*, *Ardea herodias*, *Arremonops conirostris*, *Arremonops tocuyensis*, *Atalotriccus pilaris*, *Attila spadiceus* (Gmelin 1789), *Basileuterus flaveolus* (S. F. Baird 1865), *Bubo virginianus* (Gmelin 1788), *Busarellus nigricollis*, *Buteogallus anthracinus*, *Buteogallus meridionalis*, *Buteogallus schistaceus*, *Buteogallus solitarius*, *Buteogallus urubitinga*, *Cacicus cela*, *Cairina moschata*, *Campylorhamphus trochilirostris* (Lichtenstein 1820), *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Caprimulgus cayennensis* (Gmelin 1789), *Cardinalis phoeniceus* Bonaparte 1838, *Carduelis psaltria* (Say 1823), *Cathartes aura*, *Cathartes burrovianus*, *Cathartes melambrotus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Ceryle torquata*, *Chlorostilbon gibsoni* (Fraser 1840), *Chondrohierax uncinatus*, *Chordeiles acutipennis* (Hermann 1783), *Chrysomus icterocephalus*, *Circus buffoni*, *Circus cinereus*, *Circus cyaneus*, *Coccyzus pumila*, *Columbina minuta* (Linnaeus 1766), *Conirostrum leucogenys* (Lafresnaye 1852), *Coragyps atratus*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Crax alector*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cryptoleucopteryx plumbea*, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Dryocopus lineatus* (Linnaeus 1766), *Elaenia chiriquensis* Lawrence 1865, *Elanoides forficatus*, *Elanus leucurus*, *Falco deiroleucus*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Furnarius leucopus* Swainson 1838, *Galbula ruficauda* Cuvier 1816, *Gallinula chloropus* (Linnaeus 1758), *Gampsonyx swainsonii*, *Geranospiza caerulescens*, *Glaucidium brasilianum* (Gmelin 1788), *Harpagus bidentatus*, *Harpia harpyja*, *Helicolestes hamatus*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresna), *Himantopus mexicanus* (Statius Muller 1776), *Hylophilus flavipes* Lafresnaye 1845, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Ictinia mississippiensis*, *Ictinia plumbea*, *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Jabiru mycteria*, *Jacana jacana*, *Lepidopyga goudoti* (Bourcier 1843), *Leptotila verreauxi* (Bonaparte 1855), *Leucopternis melanops*, *Leucopternis semiplumbea*, *Machetornis rixosa*, *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Micrastur plumbeus*, *Mitu tomentosa*, *Molothrus bonariensis*, *Morphnarchus princeps*, *Morphnus guianensis*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neochen jubata*, *Opisthocomus hoazin*, *Oroaetus isidori*, *Ortalis ruficauda* Jardine 1847, *Pandion haliaetus*, *Parabuteo unicinctus*, *Patagioenas corensis* (Jacquin 1784), *Phaeomyias murina* (Spix 1825), *Pheucticus ludovicianus*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pionus sordidus* (Linnaeus 1758), *Pitangus lictor* (Lichtenstein 1823), *Pitangus sulphuratus*, *Polioptila plumbea* (J. F. Gmelin 1788), *Pseudastur albicollis*, *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Quiscalus mexicanus* (Gmelin 1788), *Rostrhamus sociabilis*, *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Sarcoramphus papa*, *Sarkidiornis melanotos* (Pennant 1769), *Spiza americana*, *Spizaetus tyrannus*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx*

ruficollis (Vieillot 1817), *Sterna superciliaris*, *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tachycineta bicolor* (Vieillot 1808), *Tapera naevia* (Linnaeus 1766), *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Zenaida auriculata*

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

fullDescriptionUnstructured: Mide unos 13 cm. Cabeza y nuca gris pálido con una lista central negra en la coronilla y una superciliar, que se prolonga hasta la nuca de color arcilla blanquecina. El resto de las partes dorsales color verde oliva; por debajo de color gris claro (Hilty & Brown 1986).

briefDescription:

references

link: <http://collections.mnh.si.edu/search/>

keywords: *Amazona ochrocephala*, *Ameiva festiva*, *Anas discors*, *Anhinga anhinga* (Linnaeus 1766), *Aramides cajanea*, *Aramus guarauna*, *Aratinga acuticaudata*, *Ardea herodias*, *Arremonops conirostris*, *Arremonops tocuyensis*, *Atalotriccus pilaris*, *Attila spadiceus* (Gmelin 1789), *Basileuterus flaveolus* (S. F. Baird 1865), *Bubo virginianus* (Gmelin 1788), *Burhinus bistriatus* (Wagler 1829), *Campylorhynchus trochilirostris* (Lichtenstein 1820), *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Caprimulgus cayennensis* (Gmelin 1789), *Cardinalis phoeniceus* Bonaparte 1838, *Carduelis psaltria* (Say 1823), *Carollia brevicauda* (Schinz 1821), *Centurio senex* Gray 1842, *Certhiaxis cinnamomeus* (Gmelin 1788), *Chlorostilbon gibsoni* (Fraser 1840), *Chordeiles acutipennis* (Hermann 1783), *Chrysolampis mosquitos* (Linnaeus 1758), *Chrysomus icterocephalus*, *Coccyzus pumila*, *Colibri coruscans*, *Columbina minuta* (Linnaeus 1766), *Columbina passerina* (Linnaeus 1758), *Conirostrum leucogenys* (Lafresnaye 1852), *Coryphospingus pileatus* (Wied-Neuwied 1821), *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Dendrocygna autumnalis*, *Dryocopus lineatus* (Linnaeus 1766), *Egretta thula*, *Egretta tricolor* (Statius Muller 1776), *Elaenia chiriquensis* Lawrence 1865, *Engystomops pustulosus*, *Eumops glaucinus* (Wagner 1843), *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Furnarius leucopus* Swainson 1838, *Galbula ruficauda* Cuvier 1816, *Gallinula chloropus* (Linnaeus 1758), *Glaucidium brasilianum* (Gmelin 1788), *Glossophaga longirostris* Miller 1898, *Gonatodes vittatus*, *Heliodoxa leadbeateri* (Bourcier 1843), *Hemitriccus margaritaceiventer* (Orbigny & Lafresnaye), *Himantopus mexicanus* (Statius Muller 1776), *Hylophilus flavipes* Lafresnaye 1845, *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Lasiurus ega* (Gervais 1856), *Lepidoblepharis sanctaemartae* Ruthven 1916, *Lepidopygia goudoti* (Bourcier 1843), *Leptonycteris curasoae* Miller 1900, *Leptotila verreauxi* (Bonaparte 1855), *Leucippus fallax* (Bourcier 1843), *Lophostoma silvicolium* (d'Orbigny 1836), *Machetornis rixosa*, *Macrophyllum macrophyllum* (Schinz 1821), *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Micronycteris hirsuta* (Peters 1869), *Molothrus bonariensis*, *Mormoops megalophylla* (Peters

1864), *Mycteria americana* Linnaeus 1758, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Noctilio albiventris* (Desmarest 1818), *Nycticorax nycticorax*, *Ortalis ruficauda* Jardine 1847, *Pachyrhamphus rufus* (Boddaert 1783), *Patagioenas corensis* (Jacquin 1784), *Phaeomyias murina* (Spix 1825), *Pheucticus ludovicianus*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pionus sordidus* (Linnaeus 1758), *Piranga flava* (Vieillot 1822), *Platalea ajaja* Linnaeus 1758, *Pleurodema brachyops*, *Polioptila plumbea* (J. F. Gmelin 1788), *Pseudopaludicola pusilla*, *Pteronotus davyi* (Gray 1838), *Pteronotus gymnonotus* (Natterer 1843), *Pteronotus parnellii* (Gray 1843), *Pteronotus personatus* (Wagner 1843), *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Quiscalus mexicanus* (Gmelin 1788), *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Spiza americana*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx ruficollis* (Vieillot 1817), *Streptoprocne zonaris* (Shaw 1796), *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Tachycineta albiventer* (Boddaert 1783), *Tachycineta bicolor* (Vieillot 1808), *Tapera naevia* (Linnaeus 1766), *Thamnophilus doliaatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tolmomyias sulphurescens* (Spix 1825), *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Zenaida auriculata*

editors

institution:
address:
publisher:
accessed:
websites:
chapter:
series:
issue:
year: 2012

authors

authors: Smithsonian National Museum of Natural History

title: Search Museum Collections Records

source: SMITHSONIAN NATIONAL MUSEUM OF NATURAL HISTORY

type: miscellany

tags: *Amazona ochrocephala*, *Ameiva festiva*, *Anas discors*, *Anhinga anhinga* (Linnaeus 1766), *Aramides cajanea*, *Aramus guarauna*, *Aratinga acuticaudata*, *Ardea herodias*, *Arremonops conirostris*, *Arremonops tocuyensis*, *Atalotriccus pilaris*, *Attila spadiceus* (Gmelin 1789), *Basileuterus flaveolus* (S. F. Baird 1865), *Bubo virginianus* (Gmelin 1788), *Burhinus bistriatus* (Wagler 1829), *Campylorhamphus trochilirostris* (Lichtenstein 1820), *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Caprimulgus cayennensis* (Gmelin 1789), *Cardinalis phoeniceus* Bonaparte 1838, *Carduelis psaltria* (Say 1823), *Carollia brevicauda* (Schinz 1821), *Centurio senex* Gray 1842, *Certhiaxis cinnamomeus* (Gmelin 1788), *Chlorostilbon gibsoni* (Fraser 1840), *Chordeiles acutipennis* (Hermann 1783), *Chrysolampis mosquitos* (Linnaeus 1758), *Chrysomus icterocephalus*, *Coccyzus pumila*, *Colibri coruscans*, *Columbina minuta* (Linnaeus 1766), *Columbina passerina* (Linnaeus 1758), *Conirostrum leucogenys* (Lafresnaye 1852), *Coryphospingus pileatus* (Wied-Neuwied 1821), *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cyanocorax affinis* (Pelzeln 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Dendrocygna autumnalis*, *Dryocopus lineatus* (Linnaeus 1766), *Egretta thula*, *Egretta tricolor* (Statius Muller 1776), *Elaenia chiriquensis* Lawrence 1865, *Engystomops pustulosus*, *Eumops glaucinus* (Wagner 1843), *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Furnarius leucopus* Swainson 1838, *Galbula ruficauda* Cuvier 1816, *Gallinula chloropus* (Linnaeus 1758), *Glaucidium brasilianum* (Gmelin 1788), *Glossophaga longirostris* Miller 1898, *Gonatodes vittatus*, *Heliodoxa leadbeateri* (Bourcier 1843), *Hemitriccus margaritaceiventer* (Orbigny & Lafresna), *Himantopus mexicanus* (Statius Muller 1776), *Hylophilus flavipes* Lafresnaye 1845, *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Lasiurus ega* (Gervais 1856), *Lepidoblepharis sanctaemartae* Ruthven 1916, *Lepidopyga goudoti* (Bourcier 1843), *Leptonycteris curasoae* Miller 1900, *Leptotila verreauxi* (Bonaparte 1855), *Leucippus fallax* (Bourcier 1843), *Lophostoma silvicolium* (d'Orbigny 1836), *Machetornis rixosa*, *Macrophyllum macrophyllum* (Schinz 1821), *Megaceryle alcyon* (Linnaeus

1758),Megaceryle torquata (Linnaeus 1766),Megarynchus pitangua (Linnaeus 1766),Megascops choliba,Melanerpes rubricapillus,Micronycteris hirsuta (Peters 1869),Molothrus bonariensis,Mormoops megalophylla (Peters 1864),Mycteria americana Linnaeus 1758,Myiarchus crinitus,Myiarchus venezuelensis Lawrence 1865,Myiopagis gaimardii (Orbigny 1840),Myiozetetes similis (Spix 1825),Myospiza humeralis (Bosc 1792),Nemosia pileata (Boddaert 1783),Noctilio albiventris (Desmarest 1818),Nycticorax nycticorax,Ortalis ruficauda Jardine 1847,Pachyramphus rufus (Boddaert 1783),Patagioenas corensis (Jacquin 1784),Phaeomyias murina (Spix 1825),Pheucticus ludovicianus,Piculus chrysochloros (Vieillot 1818),Picumnus cinnamomeus Wagler 1829,Pilherodius pileatus (Boddaert 1783),Pionus sordidus (Linnaeus 1758),Piranga flava (Vieillot 1822),Platalea ajaja Linnaeus 1758,Pleurodema brachyops,Polioptila plumbea (J. F. Gmelin 1788),Pseudopaludicola pusilla,Pteronotus davyi (Gray 1838),Pteronotus gymnonotus (Natterer 1843),Pteronotus parnellii (Gray 1843),Pteronotus personatus (Wagner 1843),Pulsatrix perspicillata (Latham 1790),Pyrocephalus rubinus,Quiscalus lugubris,Quiscalus mexicanus (Gmelin 1788),Sakesphorus canadensis (Linnaeus 1766),Saltator coerulescens Vieillot 1817,Spiza americana,Sporophila bouvronides (Lesson 1831),Sporophila intermedia Cabanis 1851,Sporophila nigricollis (Vieillot 1823),Stelgidopteryx ruficollis (Vieillot 1817),Streptoprocne zonaris (Shaw 1796),Sturnella magna,Sublegatus arenarum (Salvin 1863),Tachycineta albiventer (Boddaert 1783),Tachycineta bicolor (Vieillot 1808),Tapera naevia (Linnaeus 1766),Thamnophilus doliatus (Linnaeus 1764),Thraupis glaucocolpa Cabanis 1850,Thraupis palmarum,Thryothorus leucotis (Lafresnaye 1845),Tolmomyias sulphurescens (Spix 1825),Tyrannus dominicensis (Gmelin 1788),Tyto alba,Zenaida auriculata

abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:

note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:

identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: Todd, 1912

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Arremonops tocuyensis
rank: SPECIES
simple: Arremonops tocuyensis Todd, 1912

attributes

nomenclaturalCode:

isAnamorphic:

id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-15T05:06:44.381Z

created: 2016-03-15T05:06:44.381Z

[agent](#)

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataType:

identifier:

parentTaxon: Arremonops

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Arremonops
family: Emberizidae
order: Passeriformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

commonNameAtomized

ancillaryData

usedBy:
usedIn
distributionUnstructured:
endDate:
startDate:

ancillaryData

type:
language: Español
name: Pinzón guajiro

invasiveness
invasivenessUnstructured:
reproduction
reproductionUnstructured:
molecularData
molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions
interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:

0

keywords: *Acacia glomerosa*, *Acacia tamarindifolia*, *Acacia tortuosa*, *Alternanthera halimifolia*, *Amazona ochrocephala*, *Ameiva festiva*, *Anas discors*, *Anemopaegma orbiculatum*, *Anhinga anhinga* (Linnaeus 1766), *Anoda crenatiflora*, *Aphelandra pulcherrima*, *Aramides cajanea*, *Aramus guarauna*, *Aratinga acuticaudata*, *Ardea herodias*, *Aristolochia anguicida*, *Arremonops conirostris*, *Arremonops tocuyensis*, *Artemisia vulgaris*, *Aspidosperma cuspa*, *Atalotriccus pilaris*, *Ayenia magna*, *Basileuterus flaveolus* (S. F. Baird 1865), *Bastardia viscosa*, *Batis maritima*, *Bauhinia pauletia*, *Boerhavia scandens*, *Bonellia frutescens*, *Bourreria cumanensis*, *Bouteloua aristidoides*, *Bromelia chrysantha*, *Bubo virginianus* (Gmelin 1788), *Burhinus bistriatus* (Wagler 1829), *Butorides virescens*, *Caesalpinia bonduc*, *Caesalpinia mollis*, *Calotropis gigantea*, *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Canavalia brasiliensis*, *Canavalia rosea*, *Capparis frondosa*, *Capparis linearis*, *Capparis pulcherrima*, *Caprimulgus cayennensis* (Gmelin 1789), *Capsicum baccatum*, *Cardinalis phoeniceus* Bonaparte 1838, *Cardiospermum corindum*, *Carduelis psaltria* (Say 1823), *Castela erecta*, *Celtis iguanaea*, *Cenchrus brownii*, *Cenchrus ciliaris*, *Cenchrus echinatus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Chlorostilbon gibsoni* (Fraser 1840), *Chordeiles acutipennis* (Hermann 1783), *Chrysolampis mosquitus* (Linnaeus 1758), *Chrysomus icterocephalus*, *Cissus trifoliata*, *Citrullus lanatus*, *Cleome spinosa*, *Cnidoscopus urens*, *Coccoloba microstachya*, *Columbina minuta* (Linnaeus 1766), *Columbina passerina* (Linnaeus 1758), *Combretum fruticosum*, *Conirostrum leucogenys* (Lafresnaye 1852), *Conyza canadensis*, *Conyza perijaensis*, *Cordia macuirensis*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Craniolaria annua*, *Crotalaria purdiana*, *Crotalaria retusa*, *Croton ferrugineus*, *Croton malambo*, *Croton niveus*, *Croton ovalifolius*, *Croton punctatus*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Cyperus articulatus*, *Datura innoxia*, *Dendrocygna autumnalis*, *Desmodium incanum*, *Diphysa carthagenensis*, *Ditaxis argothamnoides*, *Dolichandra unguis-cati*, *Dryocopus lineatus* (Linnaeus 1766), *Dysphania ambrosioides*, *Egletes prostrata*, *Egretta thula*, *Egretta tricolor* (Statius Muller 1776), *Elaenia chiriquensis* Lawrence 1865, *Eleocharis elegans*, *Eleocharis interstincta*, *Engystomops pustulosus*, *Eragrostis ciliaris*, *Erythrina berterioana*, *Euphorbia hirta*, *Evolvulus convolvuloides*, *Evolvulus sericeus*, *Exostema caribaeum*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Fridericia chica*, *Fridericia dichotoma*, *Fridericia mollissima*, *Funastrum clausum*, *Furnarius leucopus* Swainson 1838, *Galactia striata*, *Galbula ruficauda* Cuvier 1816, *Glaucidium brasilianum* (Gmelin 1788), *Glossophaga longirostris* Miller 1898, *Gonatodes vittatus*, *Handroanthus serratifolius*, *Heliotropium angiospermum*, *Heliotropium indicum*, *Heliotropium procumbens*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresna), *Hibiscus phoeniceus*, *Hillia costanensis*, *Himantopus mexicanus* (Statius Muller 1776), *Hippomane mancinella*, *Hylophilus flavipes* Lafresnaye 1845, *Hymenocallis caribaea*, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Ipomoea incarnata*, *Jacquinia armillaris*, *Laguncularia racemosa*, *Lantana fucata*, *Lepidoblepharis sanctaemartae* Ruthven 1916, *Leptodactylus poecilochilus*, *Leptonycteris curasoae* Miller 1900, *Leptotila verreauxi* (Bonaparte 1855), *Licania cuspidata*, *Lonchocarpus atropurpureus*, *Lonchocarpus sanctae-marthae*, *Ludwigia octovalvis*, *Luffa operculata*, *Machetornis rixosa*, *Mallotus rhamnifolius*, *Manihot carthaginensis*, *Matelea maritima*, *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Melocactus intortus*, *Melochia parvifolia*, *Melochia tomentosa*, *Mentzelia aspera*, *Merremia aegyptia*, *Merremia umbellata*, *Micronycteris hirsuta* (Peters 1869), *Mimosa arenosa*, *Mimosa leiocarpa*, *Mimosa pellita*, *Mollugo verticillata*, *Molothrus bonariensis*, *Morisonia americana*, *Mormoops megalophylla* (Peters 1864), *Musa paradisiaca*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neurolaena lobata*, *Noctilio leporinus* (Linnaeus 1758), *Nycticorax nycticorax*, *Pachyramphus rufus* (Boddaert 1783), *Parkia velutina*, *Parkinsonia praecox*, *Passiflora foetida*, *Patagioenas corensis* (Jacquin 1784), *Paullinia cururu*, *Phaeomyias murina* (Spix 1825), *Phaseolus lunatus*, *Pheucticus ludovicianus*, *Phoradendron mucronatum*, *Phthirusa stelis*, *Phyla nodiflora*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pilosocereus lanuginosus*, *Pionus sordidus* (Linnaeus 1758), *Piptadenia viridiflora*, *Piranga flava* (Vieillot 1822), *Pitangus lictor* (Lichtenstein 1823), *Platalea ajaja* Linnaeus 1758, *Pleurodema brachyops*, *Polioptila plumbea* (J. F. Gmelin 1788), *Porophyllum ruderae*, *Pseudopaludicola pusilla*, *Pteronotus gymnonotus* (Natterer 1843), *Pteronotus parnellii* (Gray 1843), *Pteronotus personatus* (Wagner 1843), *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Rhizophora mangle*, *Rivina humilis*, *Rosenbergiodendron formosum*, *Ruellia paniculata*, *Ruprechtia ramiflora*, *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Sarcostemma glaucum*, *Scoparia dulcis*, *Senna atomaria*, *Senna bicapsularis*, *Senna candolleana*, *Senna italica*, *Senna occidentalis*, *Senna tora*, *Serjania mexicana*, *Sesuvium portulacastrum*, *Sida ciliaris*, *Solandra grandiflora*, *Solanum bicolor*, *Spiza americana*, *Sporobolus coromandelianus*, *Sporobolus pyramidalis*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis

1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx ruficollis* (Vieillot 1817), *Stemodia maritima*, *Streptoprocne zonaris* (Shaw 1796), *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tabernaemontana amygdalifolia*, *Tachycineta albiventer* (Boddaert 1783), *Tachycineta bicolor* (Vieillot 1808), *Talinum paniculatum*, *Tapera naevia* (Linnaeus 1766), *Tephrosia cinerea*, *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tiaris bicolor* (Linnaeus 1766), *Tiaris obscurus* (Orbigny & Lafresnaye 1837), *Tillandsia flexuosa*, *Tityra inquisitor* (Lichtenstein 1823), *Tolmomyias sulphurescens* (Spix 1825), *Tournefortia hirsutissima*, *Trianthema portulacastrum*, *Tribulus cistoides*, *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Vitex compressa*, *Wissadula periplocifolia*, *Zenaida auriculata*

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 417

issue:

year: 2010

0: Cerrejón

title: Monitoreo de Fauna en el Valle del Cerrejón

source:

type: report

tags: *Acacia glomerosa*, *Acacia tamarindifolia*, *Acacia tortuosa*, *Alternanthera halimifolia*, *Amazona ochrocephala*, *Ameiva festiva*, *Anas discors*, *Anemopaegma orbiculatum*, *Anhinga anhinga* (Linnaeus 1766), *Anoda crenatiflora*, *Aphelandra pulcherrima*, *Aramides cajanea*, *Aramus guarauna*, *Aratinga acuticaudata*, *Ardea herodias*, *Aristolochia anguicida*, *Arremonops conirostris*, *Arremonops tocuyensis*, *Artemisia vulgaris*, *Aspidosperma cuspa*, *Atalotriccus pilaris*, *Ayenia magna*, *Basileuterus flaveolus* (S. F. Baird 1865), *Bastardia viscosa*, *Batis maritima*, *Bauhinia pauletia*, *Boerhavia scandens*, *Bonellia frutescens*, *Bourreria cumanaensis*, *Bouteloua aristidoides*, *Bromelia chrysantha*, *Bubo virginianus* (Gmelin 1788), *Burhinus bistriatus* (Wagler 1829), *Butorides virescens*, *Caesalpinia bonduc*, *Caesalpinia mollis*, *Calotropis gigantea*, *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Canavalia brasiliensis*, *Canavalia rosea*, *Capparis frondosa*, *Capparis linearis*, *Capparis pulcherrima*, *Caprimulgus cayennensis* (Gmelin 1789), *Capsicum baccatum*, *Cardinalis phoeniceus* Bonaparte 1838, *Cardiospermum corindum*, *Carduelis psaltria* (Say 1823), *Castela erecta*, *Celtis iguanaea*, *Cenchrus brownii*, *Cenchrus ciliaris*, *Cenchrus echinatus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Chlorostilbon gibsoni* (Fraser 1840), *Chordeiles acutipennis* (Hermann 1783), *Chrysolampis mosquitos* (Linnaeus 1758), *Chrysomus icterocephalus*, *Cissus trifoliata*, *Citrullus lanatus*, *Cleome spinosa*, *Cnidoscopus urens*, *Coccoloba microstachya*, *Columbina minuta* (Linnaeus 1766), *Columbina passerina* (Linnaeus 1758), *Combretum fruticosum*, *Conirostrum leucogenys* (Lafresnaye 1852), *Conyza canadensis*, *Conyza perijaensis*, *Cordia macuirensis*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Craniolaria annua*, *Crotalaria purdiana*, *Crotalaria retusa*, *Croton ferrugineus*, *Croton malambo*, *Croton niveus*, *Croton ovalifolius*, *Croton punctatus*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Cyperus articulatus*, *Datura innoxia*, *Dendrocycna autumnalis*, *Desmodium incanum*, *Diphysa carthagenensis*, *Ditaxis argothamnoides*, *Dolichandra unguis-cati*, *Dryocopus lineatus* (Linnaeus 1766), *Dysphania ambrosioides*, *Egletes prostrata*, *Egretta thula*, *Egretta tricolor* (Statius Muller 1776), *Elaenia chiriquensis* Lawrence 1865, *Eleocharis elegans*, *Eleocharis interstincta*, *Engystomops pustulosus*, *Eragrostis ciliaris*, *Erythrina berteriana*, *Euphorbia hirta*, *Evolvulus convolvuloides*, *Evolvulus sericeus*, *Exostema caribaeum*, *Fluicicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Fridericia chica*, *Fridericia dichotoma*, *Fridericia mollissima*, *Funastrum clausum*, *Furnarius leucopus* Swainson 1838, *Galactia striata*, *Galbula ruficauda* Cuvier 1816, *Glaucidium brasilianum* (Gmelin 1788), *Glossophaga longirostris* Miller 1898, *Gonatodes vittatus*, *Handroanthus serratifolius*, *Heliotropium angiospermum*, *Heliotropium indicum*, *Heliotropium procumbens*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresna), *Hibiscus phoeniceus*, *Hillia costanensis*, *Himantopus mexicanus* (Statius Muller 1776), *Hippomane mancinella*, *Hylophilus flavipes* Lafresnaye 1845, *Hymenocallis caribaea*, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Ipomoea incarnata*, *Jacquinia armillaris*, *Laguncularia racemosa*, *Lantana fucata*, *Lepidoblepharis*

sanctaemartae Ruthven 1916,Leptodactylus poecilochilus,Leptonycteris curasoae Miller 1900,Leptotila verreauxi (Bonaparte 1855),Licania cuspidata,Lonchocarpus atropurpureus,Lonchocarpus sanctae-marthae,Ludwigia octovalvis,Luffa operculata,Machetornis rixosa,Mallotus rhamnifolius,Manihot carthaginensis,Matelea maritima,Megaceryle alcyon (Linnaeus 1758),Megaceryle torquata (Linnaeus 1766),Megarynchus pitangua (Linnaeus 1766),Megascops choliba,Melanerpes rubricapillus,Melocactus intortus,Melochia parvifolia,Melochia tomentosa,Mentzelia aspera,Merremia aegyptia,Merremia umbellata,Micronycteris hirsuta (Peters 1869),Mimosa arenosa,Mimosa leiocarpa,Mimosa pellita,Mollugo verticillata,Molothrus bonariensis,Morisonia americana,Mormoops megalophylla (Peters 1864),Musa paradisiaca,Myiarchus crinitus,Myiarchus venezuelensis Lawrence 1865,Myiopagis gaimardii (Orbigny 1840),Myiozetetes similis (Spix 1825),Myospiza humeralis (Bosc 1792),Nemosia pileata (Boddaert 1783),Neurolaena lobata,Noctilio leporinus (Linnaeus 1758),Nycticorax nycticorax,Pachyrhamphus rufus (Boddaert 1783),Parkia velutina,Parkinsonia praecox,Passiflora foetida,Patagioenas corensis (Jacquin 1784),Paullinia cururu,Phaeomyias murina (Spix 1825),Phaseolus lunatus,Pheucticus ludovicianus,Phoradendron mucronatum,Phthirusa stelis,Phyla nodiflora,Piculus chrysochloros (Vieillot 1818),Picumnus cinnamomeus Wagler 1829,Pilherodius pileatus (Boddaert 1783),Pilosocereus lanuginosus,Pionus sordidus (Linnaeus 1758),Piptadenia viridiflora,Piranga flava (Vieillot 1822),Pitangus lictor (Lichtenstein 1823),Platalea ajaja Linnaeus 1758,Pleurodema brachyops,Polioptila plumbea (J. F. Gmelin 1788),Porophyllum ruderae,Pseudopaludicola pusilla,Pteronotus gymnotus (Natterer 1843),Pteronotus parnellii (Gray 1843),Pteronotus personatus (Wagner 1843),Pulsatrix perspicillata (Latham 1790),Pyrocephalus rubinus,Quiscalus lugubris,Rhizophora mangle,Rivina humilis,Rosenbergiodendron formosum,Ruellia paniculata,Ruprechtia ramiflora,Sakesphorus canadensis (Linnaeus 1766),Saltator coerulescens Vieillot 1817,Sarcostemma glaucum,Scoparia dulcis,Senna atomaria,Senna bicapsularis,Senna candolleana,Senna italica,Senna occidentalis,Senna tora,Serjania mexicana,Sesuvium portulacastrum,Sida ciliaris,Solandra grandiflora,Solanum bicolor,Spiza americana,Sporobolus coromandelianus,Sporobolus pyramidalis,Sporophila bouvronides (Lesson 1831),Sporophila intermedia Cabanis 1851,Sporophila nigricollis (Vieillot 1823),Stelgidopteryx ruficollis (Vieillot 1817),Stemodia maritima,Streptoprocne zonaris (Shaw 1796),Sturnella magna,Sublegatus arenarum (Salvin 1863),Synallaxis candei Orbigny & Lafresnaye 1838,Tabernaemontana amygdalifolia,Tachycineta albiventer (Boddaert 1783),Tachycineta bicolor (Vieillot 1808),Talinum paniculatum,Tapera naevia (Linnaeus 1766),Tephrosia cinerea,Thamnophilus doliatus (Linnaeus 1764),Thraupis glaucocolpa Cabanis 1850,Thraupis palmarum,Thryothorus leucotis (Lafresnaye 1845),Tiaris bicolor (Linnaeus 1766),Tiaris obscurus (Orbigny & Lafresnaye 1837),Tillandsia flexuosa,Tityra inquisitor (Lichtenstein 1823),Tolmomyias sulphurescens (Spix 1825),Tournefortia hirsutissima,Trianthema portulacastrum,Tribulus cistoides,Tyrannus dominicensis (Gmelin 1788),Tyto alba,Vireo flavoviridis (Cassin 1851),Vitex compressa,Wissadula periplocifolia,Zenaida auriculata

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

data Type:

identifier:

feedingUnstructured: Su dieta en La Guajira se compone principalmente de granos (Cerrejón 2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: *Accipiter collaris*, *Accipiter cooperii*, *Accipiter poliogaster*, *Accipiter superciliosus*, *Amazona ochrocephala*, *Andigena hypoglaucha*, *Aramus guarauna*, *Aratinga pertinax*, *Ardea alba*, *Ardea cocoi*, *Ardea herodias*, *Arremonops conirostris*, *Arremonops tocuyensis*, *Atalotriccus pilaris*, *Attila spadiceus* (Gmelin 1789), *Basileuterus flaveolus* (S. F. Baird 1865), *Bubo virginianus* (Gmelin 1788), *Busarellus nigricollis*, *Buteogallus anthracinus*, *Buteogallus meridionalis*, *Buteogallus schistaceus*, *Buteogallus solitarius*, *Buteogallus urubitinga*, *Cacicus cela*, *Cairina moschata*, *Campylorhamphus trochilirostris* (Lichtenstein 1820), *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Caprimulgus cayennensis* (Gmelin 1789), *Cardinalis phoeniceus* Bonaparte 1838, *Carduelis psaltria* (Say 1823), *Cathartes aura*, *Cathartes burrovianus*, *Cathartes melambrotus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Ceryle torquata*, *Chlorostilbon gibsoni* (Fraser 1840), *Chondrohierax uncinatus*, *Chordeiles acutipennis* (Hermann 1783), *Chrysomus icterocephalus*, *Circus buffoni*, *Circus cinereus*, *Circus cyaneus*, *Coccyzus pumila*, *Columbina minuta* (Linnaeus 1766), *Conirostrum leucogenys* (Lafresnaye 1852), *Coragyps atratus*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Crax alector*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cryptoleucopteryx plumbea*, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Dryocopus lineatus* (Linnaeus 1766), *Elaenia chiriquensis* Lawrence 1865, *Elanoides forficatus*, *Elanus leucurus*, *Falco deiroleucus*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Furnarius leucopus* Swainson 1838, *Galbula ruficauda* Cuvier 1816, *Gallinula chloropus* (Linnaeus 1758), *Gampsonyx swainsonii*, *Geranospiza caerulescens*, *Glaucidium brasilianum* (Gmelin 1788), *Harpagus bidentatus*, *Harpia harpyja*, *Helicolestes hamatus*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresna), *Himantopus mexicanus* (Statius Muller 1776), *Hylophilus flavipes* Lafresnaye 1845, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Ictinia mississippiensis*, *Ictinia plumbea*, *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Jabiru mycteria*, *Jacana jacana*, *Lepidopyga goudoti* (Bourcier 1843), *Leptotila verreauxi* (Bonaparte 1855), *Leucopternis melanops*, *Leucopternis semiplumbea*, *Machetornis rixosa*, *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Micrastur plumbeus*, *Mitu tomentosa*, *Molothrus bonariensis*, *Morphnarchus princeps*, *Morphnus guianensis*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neochen jubata*, *Opisthocomus hoazin*, *Oroaetus isidori*, *Ortalis ruficauda* Jardine 1847, *Pandion haliaetus*, *Parabuteo unicinctus*, *Patagioenas corensis* (Jacquin 1784), *Phaeomyias murina* (Spix 1825), *Pheucticus ludovicianus*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pionus sordidus* (Linnaeus 1758), *Pitangus lictor* (Lichtenstein 1823), *Pitangus sulphuratus*, *Polioptila plumbea* (J. F. Gmelin 1788), *Pseudastur albicollis*, *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Quiscalus mexicanus* (Gmelin 1788), *Rostrhamus sociabilis*, *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Sarcoramphus papa*, *Sarkidiornis melanotos* (Pennant 1769), *Spiza americana*, *Spizaetus tyrannus*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx*

ruficollis (Vieillot 1817), *Sterna supercilialis*, *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tachycineta bicolor* (Vieillot 1808), *Tapera naevia* (Linnaeus 1766), *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Zenaida auriculata*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

issue:

year: 1986

0: Hilty, S

1: Brown, W. L.

title: A Guide to the Birds of Colombia

source:

type: book

tags: *Accipiter collaris*, *Accipiter cooperii*, *Accipiter poliogaster*, *Accipiter superciliosus*, *Amazona ochrocephala*, *Andigena hypoglaucha*, *Aramus guarauna*, *Aratinga pertinax*, *Ardea alba*, *Ardea cocoi*, *Ardea herodias*, *Arremonops conirostris*, *Arremonops tocuyensis*, *Atalotriccus pilaris*, *Attila spadiceus* (Gmelin 1789), *Basileuterus flaveolus* (S. F. Baird 1865), *Bubo virginianus* (Gmelin 1788), *Busarellus nigricollis*, *Buteogallus anthracinus*, *Buteogallus meridionalis*, *Buteogallus schistaceus*, *Buteogallus solitarius*, *Buteogallus urubitinga*, *Cacicus cela*, *Cairina moschata*, *Campylorhamphus trochilirostris* (Lichtenstein 1820), *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Caprimulgus cayennensis* (Gmelin 1789), *Cardinalis phoeniceus* Bonaparte 1838, *Carduelis psaltria* (Say 1823), *Cathartes aura*, *Cathartes burrovianus*, *Cathartes melambrotus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Ceryle torquata*, *Chlorostilbon gibsoni* (Fraser 1840), *Chondrohierax uncinatus*, *Chordeiles acutipennis* (Hermann 1783), *Chrysomus icterocephalus*, *Circus buffoni*, *Circus cinereus*, *Circus cyaneus*, *Coccyzus pumila*, *Columbina minuta* (Linnaeus 1766), *Conirostrum leucogenys* (Lafresnaye 1852), *Coragyps atratus*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Crax alector*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cryptoleucopteryx plumbea*, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Dryocopus lineatus* (Linnaeus 1766), *Elaenia chiriquensis* Lawrence 1865, *Elanoides forficatus*, *Elanus leucurus*, *Falco deiroleucus*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Furnarius leucopus* Swainson 1838, *Galbula ruficauda* Cuvier 1816, *Gallinula chloropus* (Linnaeus 1758), *Gampsonyx swainsonii*, *Geranospiza caerulescens*, *Glaucidium brasilianum* (Gmelin 1788), *Harpagus bidentatus*, *Harpia harpyja*, *Helicolestes hamatus*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresnaye), *Himantopus mexicanus* (Statius Muller 1776), *Hylophilus flavipes* Lafresnaye 1845, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Ictinia mississippiensis*, *Ictinia plumbea*, *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Jabiru mycteria*, *Jacana jacana*, *Lepidopyga goudoti* (Bourcier 1843), *Leptotila verreauxi* (Bonaparte 1855), *Leucopternis melanops*, *Leucopternis semiplumbea*, *Machetornis rixosa*, *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Micrastur plumbeus*, *Mitu tomentosa*, *Molothrus bonariensis*, *Morphnarchus princeps*, *Morphnus guianensis*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neochen jubata*, *Opisthocomus hoazin*, *Oroaetus isidori*, *Ortalis ruficauda* Jardine 1847, *Pandion haliaetus*, *Parabuteo unicinctus*, *Patagioenas corensis* (Jacquin 1784), *Phaeomyias murina* (Spix 1825), *Pheucticus ludovicianus*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pionus sordidus* (Linnaeus 1758), *Pitangus lictor* (Lichtenstein 1823), *Pitangus sulphuratus*, *Polioptila plumbea* (J. F. Gmelin 1788), *Pseudastur albicollis*, *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Quiscalus mexicanus* (Gmelin 1788), *Rostrhamus sociabilis*, *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Sarcoramphus papa*, *Sarkidiornis melanotos* (Pennant 1769), *Spiza americana*, *Spizaetus tyrannus*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx ruficollis* (Vieillot 1817), *Sterna supercilialis*, *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tachycineta bicolor* (Vieillot 1808), *Tapera naevia* (Linnaeus 1766), *Thamnophilus*

doliatus (Linnaeus 1764),Thraupis glaucocolpa Cabanis 1850,Thraupis palmarum,Thryothorus leucotis (Lafresnaye 1845),Tyrannus dominicensis (Gmelin 1788),Tyto alba,Vireo flavoviridis (Cassin 1851),Zenaida auriculata

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Prefiere regiones áridas. Semiterrestre y algo arisco, usualmente solitario o en parejas separadas. Salta por el suelo o en matorrales o sube unos pocos metros en arbustos especialmente para cantar. Canto distintivo se oye con más frecuencia temprano en la mañana (Hilty & Brown 1986).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

annualCycle

annualCycleUnstructured:

habitat

0

dataObject:

additionalInformation:

0

keywords: *Accipiter collaris*, *Accipiter cooperii*, *Accipiter poliogaster*, *Accipiter superciliosus*, *Amazona ochrocephala*, *Andigena hypoglaucha*, *Aramus guarauna*, *Aratinga pertinax*, *Ardea alba*, *Ardea cocoi*, *Ardea herodias*, *Arremonops conirostris*, *Arremonops tocuyensis*, *Atalotriccus pilaris*, *Attila spadiceus* (Gmelin 1789), *Basileuterus flaveolus* (S. F. Baird 1865), *Bubo virginianus* (Gmelin 1788), *Busarellus nigricollis*, *Buteogallus anthracinus*, *Buteogallus meridionalis*, *Buteogallus schistaceus*, *Buteogallus solitarius*, *Buteogallus urubitinga*, *Cacicus cela*, *Cairina moschata*, *Campylorhamphus trochilirostris* (Lichtenstein 1820), *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Caprimulgus cayennensis* (Gmelin 1789), *Cardinalis phoeniceus* Bonaparte 1838, *Carduelis psaltria* (Say 1823), *Cathartes aura*, *Cathartes burrovianus*, *Cathartes melambrotus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Ceryle torquata*, *Chlorostilbon gibsoni* (Fraser 1840), *Chondrohierax uncinatus*, *Chordeiles acutipennis* (Hermann 1783), *Chrysomus icterocephalus*, *Circus buffoni*, *Circus cinereus*, *Circus cyaneus*, *Coccyzus pumila*, *Columbina minuta* (Linnaeus 1766), *Conirostrum leucogenys* (Lafresnaye 1852), *Coragyps atratus*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Crax alector*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cryptoleucopteryx plumbea*, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Dryocopus lineatus* (Linnaeus 1766), *Elaenia chiriquensis* Lawrence 1865, *Elanoides forficatus*, *Elanus leucurus*, *Falco deiroleucus*, *Fluvicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Furnarius leucopus* Swainson 1838, *Galbula ruficauda* Cuvier 1816, *Gallinula chloropus* (Linnaeus 1758), *Gampsonyx swainsonii*, *Geranospiza caerulescens*, *Glaucidium brasilianum* (Gmelin 1788), *Harpagus bidentatus*, *Harpia harpyja*, *Helicolestes hamatus*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresna, *Himantopus mexicanus* (Statius Muller 1776), *Hylophilus flavipes* Lafresnaye 1845, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Ictinia mississippiensis*, *Ictinia plumbea*, *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Jabiru mycteria*, *Jacana jacana*, *Lepidopyga goudoti* (Bourcier 1843), *Leptotila verreauxi* (Bonaparte 1855), *Leucopternis melanops*, *Leucopternis semiplumbea*, *Machetornis rixosa*, *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Micrastur plumbeus*, *Mitu tomentosa*, *Molothrus bonariensis*, *Morphnarchus princeps*, *Morphnus guianensis*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neochen jubata*, *Opisthocomus hoazin*, *Oroaetus isidori*, *Ortalis ruficauda* Jardine 1847, *Pandion haliaetus*, *Parabuteo unicinctus*, *Patagioenas corensis* (Jacquin 1784), *Phaeomyias murina* (Spix 1825), *Pheucticus ludovicianus*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pionus sordidus* (Linnaeus 1758), *Pitangus lictor* (Lichtenstein 1823), *Pitangus sulphuratus*, *Polioptila plumbea* (J. F. Gmelin 1788), *Pseudastur albicollis*, *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Quiscalus mexicanus* (Gmelin 1788), *Rostrhamus sociabilis*, *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Sarcoramphus papa*, *Sarkidiornis melanotos* (Pennant 1769), *Spiza americana*, *Spizaetus tyrannus*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis 1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx ruficollis* (Vieillot 1817), *Sterna superciliaris*, *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tachycineta bicolor* (Vieillot 1808), *Tapera naevia* (Linnaeus 1766), *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Zenaida auriculata*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:
issue:
year: 1986
0: Hilty, S
1: Brown, W. L.
title: A Guide to the Birds of Colombia
source:
type: book
tags: Accipiter collaris, Accipiter cooperii, Accipiter poliogaster, Accipiter superciliosus, Amazona ochrocephala, Andigena hypoglaucha, Aramus guarauna, Aratinga pertinax, Ardea alba, Ardea cocoi, Ardea herodias, Arremonops conirostris, Arremonops tocuyensis, Atalotriccus pilaris, Attila spadiceus (Gmelin 1789), Basileuterus flaveolus (S. F. Baird 1865), Bubo virginianus (Gmelin 1788), Busarellus nigricollis, Buteogallus anthracinus, Buteogallus meridionalis, Buteogallus schistaceus, Buteogallus solitarius, Buteogallus urubitinga, Cacicus cela, Cairina moschata, Campylorhamphus trochilirostris (Lichtenstein 1820, Campylorhynchus griseus (Swainson 1838), Campylorhynchus nuchalis (Cabanis 1847), Caprimulgus cayennensis (Gmelin 1789), Cardinalis phoeniceus Bonaparte 1838, Carduelis psaltria (Say 1823), Cathartes aura, Cathartes burrovianus, Cathartes melambrotus, Certhiaxis cinnamomeus (Gmelin 1788), Ceryle torquata, Chlorostilbon gibsoni (Fraser 1840), Chondrohierax uncinatus, Chordeiles acutipennis (Hermann 1783), Chrysomus icterocephalus, Circus buffoni, Circus cinereus, Circus cyaneus, Coccyzus pumila, Columbina minuta (Linnaeus 1766), Conirostrum leucogenys (Lafresnaye 1852), Coragyps atratus, Coryphospingus pileatus (Wied-Neuwied 1821), Crax alector, Crotophaga major, Crotophaga sulcirostris Swainson 1827, Cryptoleucopteryx plumbea, Cyanocorax affinis (Pelzelin 1856), Cyclarhis gujanensis (J. F. Gmelin 1789), Dryocopus lineatus (Linnaeus 1766), Elaenia chiriquensis Lawrence 1865, Elanoides forficatus, Elanus leucurus, Falco deiroleucus, Fluvicola pica (Boddaert 1783), Forpus passerinus (Linnaeus 1758), Forpus xanthopterygius (Spix 1824), Furnarius leucopus Swainson 1838, Galbula ruficauda Cuvier 1816, Gallinula chloropus (Linnaeus 1758), Gampsonyx swainsonii, Geranospiza caerulescens, Glaucidium brasilianum (Gmelin 1788), Harpagus bidentatus, Harpia harpyja, Helicolestes hamatus, Hemitriccus margaritaceiventer (Orbigny \& Lafresna, Himantopus mexicanus (Statius Muller 1776), Hylophilus flavipes Lafresnaye 1845, Hypnelus ruficollis (Wagler 1829), Icterus auricapillus Cassin 1848, Icterus galbula (Linnaeus 1758), Icterus nigrogularis (Hahn 1819), Ictinia mississippiensis, Ictinia plumbea, Inezia subflava (P. L. Sclater \& Salvin 1873), Inezia tenuirostris (Cory 1913), Jabiru mycteria, Jacana jacana, Lepidopyga goudoti (Bourcier 1843), Leptotila verreauxi (Bonaparte 1855), Leucopternis melanops, Leucopternis semiplumbea, Machetornis rixosa, Megaceryle alcyon (Linnaeus 1758), Megaceryle torquata (Linnaeus 1766), Megarynchus pitangua (Linnaeus 1766), Megascops choliba, Melanerpes rubricapillus, Micrastur plumbeus, Mitu tomentosa, Molothrus bonariensis, Morphnarchus princeps, Morphnus guianensis, Myiarchus crinitus, Myiarchus venezuelensis Lawrence 1865, Myiopagis gaimardii (Orbigny 1840), Myiozetetes similis (Spix 1825), Myospiza humeralis (Bosc 1792), Nemosia pileata (Boddaert 1783), Neechen jubata, Opisthocomus hoazin, Oroaetus isidori, Ortalis ruficauda Jardine 1847, Pandion haliaetus, Parabuteo unicinctus, Patagioenas corensis (Jacquin 1784), Phaeomyias murina (Spix 1825), Pheucticus ludovicianus, Piculus chrysochloros (Vieillot 1818), Picumnus cinnamomeus Wagler 1829, Pilherodius pileatus (Boddaert 1783), Pionus sordidus (Linnaeus 1758), Pitangus lictor (Lichtenstein 1823), Pitangus sulphuratus, Polioptila plumbea (J. F. Gmelin 1788), Pseudastur albicollis, Pulsatrix perspicillata (Latham 1790), Pyrocephalus rubinus, Quiscalus lugubris, Quiscalus mexicanus (Gmelin 1788), Rostrhamus sociabilis, Sakesphorus canadensis (Linnaeus 1766), Saltator coerulescens Vieillot 1817, Sarcoramphus papa, Sarkidiornis melanotos (Pennant 1769), Spiza americana, Spizaetus tyrannus, Sporophila bouvronides (Lesson 1831), Sporophila intermedia Cabanis 1851, Sporophila nigricollis (Vieillot 1823), Stelgidopteryx ruficollis (Vieillot 1817), Sterna supercilialis, Sturnella magna, Sublegatus arenarum (Salvin 1863), Synallaxis candei Orbigny \& Lafresnaye 1838, Tachycineta bicolor (Vieillot 1808), Tapera naevia (Linnaeus 1766), Thamnophilus doliatus (Linnaeus 1764), Thraupis glaucocolpa Cabanis 1850, Thraupis palmarum, Thryothorus leucotis (Lafresnaye 1845), Tyrannus dominicensis (Gmelin 1788), Tyto alba, Vireo flavoviridis (Cassin 1851), Zenaida auriculata
abstract:
last_modified:
created:
group_id:
profile_id:
id:

keywords: *Acacia glomerosa*, *Acacia tamarindifolia*, *Acacia tortuosa*, *Alternanthera halimifolia*, *Amazona ochrocephala*, *Ameiva festiva*, *Anas discors*, *Anemopaegma orbiculatum*, *Anhinga anhinga* (Linnaeus 1766), *Anoda crenatiflora*, *Aphelandra pulcherrima*, *Aramides cajanea*, *Aramus guarauna*, *Aratinga acuticaudata*, *Ardea herodias*, *Aristolochia anguicida*, *Arremonops conirostris*, *Arremonops tocuyensis*, *Artemisia vulgaris*, *Aspidosperma cuspa*, *Atalotriccus pilaris*, *Ayenia magna*, *Basileuterus flaveolus* (S. F. Baird 1865), *Bastardia viscosa*, *Batis maritima*, *Bauhinia pauletia*, *Boerhavia scandens*, *Bonellia frutescens*, *Bourreria cumanensis*, *Bouteloua aristidoides*, *Bromelia chrysantha*, *Bubo virginianus* (Gmelin 1788), *Burhinus bistriatus* (Wagler 1829), *Butorides virescens*, *Caesalpinia bonduc*, *Caesalpinia mollis*, *Calotropis gigantea*, *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Canavalia brasiliensis*, *Canavalia rosea*, *Capparis frondosa*, *Capparis linearis*, *Capparis pulcherrima*, *Caprimulgus cayennensis* (Gmelin 1789), *Capsicum baccatum*, *Cardinalis phoeniceus* Bonaparte 1838, *Cardiospermum corindum*, *Carduelis psaltria* (Say 1823), *Castela erecta*, *Celtis iguanaea*, *Cenchrus brownii*, *Cenchrus ciliaris*, *Cenchrus echinatus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Chlorostilbon gibsoni* (Fraser 1840), *Chordeiles acutipennis* (Hermann 1783), *Chrysolampis mosquitus* (Linnaeus 1758), *Chrysomus icterocephalus*, *Cissus trifoliata*, *Citrullus lanatus*, *Cleome spinosa*, *Cnidoscopus urens*, *Coccoloba microstachya*, *Columbina minuta* (Linnaeus 1766), *Columbina passerina* (Linnaeus 1758), *Combretum fruticosum*, *Conirostrum leucogenys* (Lafresnaye 1852), *Conyza canadensis*, *Conyza perijaensis*, *Cordia macuirensis*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Craniolaria annua*, *Crotalaria purdiana*, *Crotalaria retusa*, *Croton ferrugineus*, *Croton malambo*, *Croton niveus*, *Croton ovalifolius*, *Croton punctatus*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Cyperus articulatus*, *Datura innoxia*, *Dendrocygna autumnalis*, *Desmodium incanum*, *Diphysa carthagenensis*, *Ditaxis argothamnoides*, *Dolichandra unguis-cati*, *Dryocopus lineatus* (Linnaeus 1766), *Dysphania ambrosioides*, *Egletes prostrata*, *Egretta thula*, *Egretta tricolor* (Statius Muller 1776), *Elaenia chiriquensis* Lawrence 1865, *Eleocharis elegans*, *Eleocharis interstincta*, *Engystomops pustulosus*, *Eragrostis ciliaris*, *Erythrina berterioana*, *Euphorbia hirta*, *Evolvulus convolvuloides*, *Evolvulus sericeus*, *Exostema caribaeum*, *Fluicicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Fridericia chica*, *Fridericia dichotoma*, *Fridericia mollissima*, *Funastrum clausum*, *Furnarius leucopus* Swainson 1838, *Galactia striata*, *Galbula ruficauda* Cuvier 1816, *Glaucidium brasilianum* (Gmelin 1788), *Glossophaga longirostris* Miller 1898, *Gonatodes vittatus*, *Handroanthus serratifolius*, *Heliotropium angiospermum*, *Heliotropium indicum*, *Heliotropium procumbens*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresna), *Hibiscus phoeniceus*, *Hillia costanensis*, *Himantopus mexicanus* (Statius Muller 1776), *Hippomane mancinella*, *Hylophilus flavipes* Lafresnaye 1845, *Hymenocallis caribaea*, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Ipomoea incarnata*, *Jacquinia armillaris*, *Laguncularia racemosa*, *Lantana fucata*, *Lepidoblepharis sanctaemartae* Ruthven 1916, *Leptodactylus poecilochilus*, *Leptonycteris curasoae* Miller 1900, *Leptotila verreauxi* (Bonaparte 1855), *Licania cuspidata*, *Lonchocarpus atropurpureus*, *Lonchocarpus sanctae-marthae*, *Ludwigia octovalvis*, *Luffa operculata*, *Machetornis rixosa*, *Mallotus rhamnifolius*, *Manihot carthaginensis*, *Matelea maritima*, *Megaceryle alcyon* (Linnaeus 1758), *Megaceryle torquata* (Linnaeus 1766), *Megarynchus pitangua* (Linnaeus 1766), *Megascops choliba*, *Melanerpes rubricapillus*, *Melocactus intortus*, *Melochia parvifolia*, *Melochia tomentosa*, *Mentzelia aspera*, *Merremia aegyptia*, *Merremia umbellata*, *Micronycteris hirsuta* (Peters 1869), *Mimosa arenosa*, *Mimosa leiocarpa*, *Mimosa pellita*, *Mollugo verticillata*, *Molothrus bonariensis*, *Morisonia americana*, *Mormoops megalophylla* (Peters 1864), *Musa paradisiaca*, *Myiarchus crinitus*, *Myiarchus venezuelensis* Lawrence 1865, *Myiopagis gaimardii* (Orbigny 1840), *Myiozetetes similis* (Spix 1825), *Myospiza humeralis* (Bosc 1792), *Nemosia pileata* (Boddaert 1783), *Neurolaena lobata*, *Noctilio leporinus* (Linnaeus 1758), *Nycticorax nycticorax*, *Pachyramphus rufus* (Boddaert 1783), *Parkia velutina*, *Parkinsonia praecox*, *Passiflora foetida*, *Patagioenas corensis* (Jacquin 1784), *Paullinia cururu*, *Phaeomyias murina* (Spix 1825), *Phaseolus lunatus*, *Pheucticus ludovicianus*, *Phoradendron mucronatum*, *Phthirusa stelis*, *Phyla nodiflora*, *Piculus chrysochloros* (Vieillot 1818), *Picumnus cinnamomeus* Wagler 1829, *Pilherodius pileatus* (Boddaert 1783), *Pilosocereus lanuginosus*, *Pionus sordidus* (Linnaeus 1758), *Piptadenia viridiflora*, *Piranga flava* (Vieillot 1822), *Pitangus lictor* (Lichtenstein 1823), *Platalea ajaja* Linnaeus 1758, *Pleurodema brachyops*, *Polioptila plumbea* (J. F. Gmelin 1788), *Porophyllum ruderales*, *Pseudopaludicola pusilla*, *Pteronotus gymnonotus* (Natterer 1843), *Pteronotus parnellii* (Gray 1843), *Pteronotus personatus* (Wagner 1843), *Pulsatrix perspicillata* (Latham 1790), *Pyrocephalus rubinus*, *Quiscalus lugubris*, *Rhizophora mangle*, *Rivina humilis*, *Rosenbergiodendron formosum*, *Ruellia paniculata*, *Ruprechtia ramiflora*, *Sakesphorus canadensis* (Linnaeus 1766), *Saltator coerulescens* Vieillot 1817, *Sarcostemma glaucum*, *Scoparia dulcis*, *Senna atomaria*, *Senna bicapsularis*, *Senna candolleana*, *Senna italica*, *Senna occidentalis*, *Senna tora*, *Serjania mexicana*, *Sesuvium portulacastrum*, *Sida ciliaris*, *Solandra grandiflora*, *Solanum bicolor*, *Spiza americana*, *Sporobolus coromandelianus*, *Sporobolus pyramidalis*, *Sporophila bouvronides* (Lesson 1831), *Sporophila intermedia* Cabanis

1851, *Sporophila nigricollis* (Vieillot 1823), *Stelgidopteryx ruficollis* (Vieillot 1817), *Stemodia maritima*, *Streptoprocne zonaris* (Shaw 1796), *Sturnella magna*, *Sublegatus arenarum* (Salvin 1863), *Synallaxis candei* Orbigny & Lafresnaye 1838, *Tabernaemontana amygdalifolia*, *Tachycineta albiventer* (Boddaert 1783), *Tachycineta bicolor* (Vieillot 1808), *Talinum paniculatum*, *Tapera naevia* (Linnaeus 1766), *Tephrosia cinerea*, *Thamnophilus doliatus* (Linnaeus 1764), *Thraupis glaucocolpa* Cabanis 1850, *Thraupis palmarum*, *Thryothorus leucotis* (Lafresnaye 1845), *Tiaris bicolor* (Linnaeus 1766), *Tiaris obscurus* (Orbigny & Lafresnaye 1837), *Tillandsia flexuosa*, *Tityra inquisitor* (Lichtenstein 1823), *Tolmomyias sulphurescens* (Spix 1825), *Tournefortia hirsutissima*, *Trianthema portulacastrum*, *Tribulus cistoides*, *Tyrannus dominicensis* (Gmelin 1788), *Tyto alba*, *Vireo flavoviridis* (Cassin 1851), *Vitex compressa*, *Wissadula periplocifolia*, *Zenaida auriculata*

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 417

issue:

year: 2010

0: Cerrejón

title: Monitoreo de Fauna en el Valle del Cerrejón

source:

type: report

tags: *Acacia glomerosa*, *Acacia tamarindifolia*, *Acacia tortuosa*, *Alternanthera halimifolia*, *Amazona ochrocephala*, *Ameiva festiva*, *Anas discors*, *Anemopaegma orbiculatum*, *Anhinga anhinga* (Linnaeus 1766), *Anoda crenatiflora*, *Aphelandra pulcherrima*, *Aramides cajanea*, *Aramus guarauna*, *Aratinga acuticaudata*, *Ardea herodias*, *Aristolochia anguicida*, *Arremonops conirostris*, *Arremonops tocuyensis*, *Artemisia vulgaris*, *Aspidosperma cuspa*, *Atalotriccus pilaris*, *Ayenia magna*, *Basileuterus flaveolus* (S. F. Baird 1865), *Bastardia viscosa*, *Batis maritima*, *Bauhinia pauletia*, *Boerhavia scandens*, *Bonellia frutescens*, *Bourreria cumanaensis*, *Bouteloua aristidoides*, *Bromelia chrysantha*, *Bubo virginianus* (Gmelin 1788), *Burhinus bistriatus* (Wagler 1829), *Butorides virescens*, *Caesalpinia bonduc*, *Caesalpinia mollis*, *Calotropis gigantea*, *Campylorhynchus griseus* (Swainson 1838), *Campylorhynchus nuchalis* (Cabanis 1847), *Canavalia brasiliensis*, *Canavalia rosea*, *Capparis frondosa*, *Capparis linearis*, *Capparis pulcherrima*, *Caprimulgus cayennensis* (Gmelin 1789), *Capsicum baccatum*, *Cardinalis phoeniceus* Bonaparte 1838, *Cardiospermum corindum*, *Carduelis psaltria* (Say 1823), *Castela erecta*, *Celtis iguanaea*, *Cenchrus brownii*, *Cenchrus ciliaris*, *Cenchrus echinatus*, *Certhiaxis cinnamomeus* (Gmelin 1788), *Chlorostilbon gibsoni* (Fraser 1840), *Chordeiles acutipennis* (Hermann 1783), *Chrysolampis mosquitos* (Linnaeus 1758), *Chrysomus icterocephalus*, *Cissus trifoliata*, *Citrullus lanatus*, *Cleome spinosa*, *Cnidoscopus urens*, *Coccoloba microstachya*, *Columbina minuta* (Linnaeus 1766), *Columbina passerina* (Linnaeus 1758), *Combretum fruticosum*, *Conirostrum leucogenys* (Lafresnaye 1852), *Conyza canadensis*, *Conyza perijaensis*, *Cordia macuirensis*, *Coryphospingus pileatus* (Wied-Neuwied 1821), *Craniolaria annua*, *Crotalaria purdiana*, *Crotalaria retusa*, *Croton ferrugineus*, *Croton malambo*, *Croton niveus*, *Croton ovalifolius*, *Croton punctatus*, *Crotophaga major*, *Crotophaga sulcirostris* Swainson 1827, *Cyanocorax affinis* (Pelzelin 1856), *Cyclarhis gujanensis* (J. F. Gmelin 1789), *Cyperus articulatus*, *Datura innoxia*, *Dendrocycna autumnalis*, *Desmodium incanum*, *Diphysa carthagenensis*, *Ditaxis argothamnoides*, *Dolichandra unguis-cati*, *Dryocopus lineatus* (Linnaeus 1766), *Dysphania ambrosioides*, *Egletes prostrata*, *Egretta thula*, *Egretta tricolor* (Statius Muller 1776), *Elaenia chiriquensis* Lawrence 1865, *Eleocharis elegans*, *Eleocharis interstincta*, *Engystomops pustulosus*, *Eragrostis ciliaris*, *Erythrina berteriana*, *Euphorbia hirta*, *Evolvulus convolvuloides*, *Evolvulus sericeus*, *Exostema caribaeum*, *Fluicicola pica* (Boddaert 1783), *Forpus passerinus* (Linnaeus 1758), *Forpus xanthopterygius* (Spix 1824), *Fridericia chica*, *Fridericia dichotoma*, *Fridericia mollissima*, *Funastrum clausum*, *Furnarius leucopus* Swainson 1838, *Galactia striata*, *Galbula ruficauda* Cuvier 1816, *Glaucidium brasilianum* (Gmelin 1788), *Glossophaga longirostris* Miller 1898, *Gonatodes vittatus*, *Handroanthus serratifolius*, *Heliotropium angiospermum*, *Heliotropium indicum*, *Heliotropium procumbens*, *Hemitriccus margaritaceiventer* (Orbigny & Lafresna), *Hibiscus phoeniceus*, *Hillia costanensis*, *Himantopus mexicanus* (Statius Muller 1776), *Hippomane mancinella*, *Hylophilus flavipes* Lafresnaye 1845, *Hymenocallis caribaea*, *Hypnelus ruficollis* (Wagler 1829), *Icterus auricapillus* Cassin 1848, *Icterus galbula* (Linnaeus 1758), *Icterus nigrogularis* (Hahn 1819), *Inezia subflava* (P. L. Sclater & Salvin 1873), *Inezia tenuirostris* (Cory 1913), *Ipomoea incarnata*, *Jacquinia armillaris*, *Laguncularia racemosa*, *Lantana fucata*, *Lepidoblepharis*

sanctaemartae Ruthven 1916, Leptodactylus poecilochilus, Leptonycteris curasoae Miller 1900, Leptotila verreauxi (Bonaparte 1855), Licania cuspidata, Lonchocarpus atropurpureus, Lonchocarpus sanctae-marthae, Ludwigia octovalvis, Luffa operculata, Machetornis rixosa, Mallotus rhamnifolius, Manihot carthaginensis, Matelea maritima, Megaceryle alcyon (Linnaeus 1758), Megaceryle torquata (Linnaeus 1766), Megarynchus pitangua (Linnaeus 1766), Megascops choliba, Melanerpes rubricapillus, Melocactus intortus, Melochia parvifolia, Melochia tomentosa, Mentzelia aspera, Merremia aegyptia, Merremia umbellata, Micronycteris hirsuta (Peters 1869), Mimosa arenosa, Mimosa leiocarpa, Mimosa pellita, Mollugo verticillata, Molothrus bonariensis, Morisonia americana, Mormoops megalophylla (Peters 1864), Musa paradisiaca, Myiarchus crinitus, Myiarchus venezuelensis Lawrence 1865, Myiopagis gaimardii (Orbigny 1840), Myiozetetes similis (Spix 1825), Myospiza humeralis (Bosc 1792), Nemosia pileata (Boddaert 1783), Neurolaena lobata, Noctilio leporinus (Linnaeus 1758), Nycticorax nycticorax, Pachyramphus rufus (Boddaert 1783), Parkia velutina, Parkinsonia praecox, Passiflora foetida, Patagioenas corensis (Jacquin 1784), Paullinia cururu, Phaeomyias murina (Spix 1825), Phaseolus lunatus, Pheucticus ludovicianus, Phoradendron mucronatum, Phthirusa stelis, Phyla nodiflora, Piculus chrysochloros (Vieillot 1818), Picumnus cinnamomeus Wagler 1829, Pilherodius pileatus (Boddaert 1783), Pilosocereus lanuginosus, Pionus sordidus (Linnaeus 1758), Piptadenia viridiflora, Piranga flava (Vieillot 1822), Pitangus lictor (Lichtenstein 1823), Platalea ajaja Linnaeus 1758, Pleurodema brachyops, Polioptila plumbea (J. F. Gmelin 1788), Porophyllum ruderae, Pseudopaludicola pusilla, Pteronotus gymnotus (Natterer 1843), Pteronotus parnellii (Gray 1843), Pteronotus personatus (Wagner 1843), Pulsatrix perspicillata (Latham 1790), Pyrocephalus rubinus, Quiscalus lugubris, Rhizophora mangle, Rivina humilis, Rosenbergiodendron formosum, Ruellia paniculata, Ruprechtia ramiflora, Sakesphorus canadensis (Linnaeus 1766), Saltator coerulescens Vieillot 1817, Sarcostemma glaucum, Scoparia dulcis, Senna atomaria, Senna bicapsularis, Senna candolleana, Senna italica, Senna occidentalis, Senna tora, Serjania mexicana, Sesuvium portulacastrum, Sida ciliaris, Solandra grandiflora, Solanum bicolor, Spiza americana, Sporobolus coromandelianus, Sporobolus pyramidalis, Sporophila bouvronides (Lesson 1831), Sporophila intermedia Cabanis 1851, Sporophila nigricollis (Vieillot 1823), Stelgidopteryx ruficollis (Vieillot 1817), Stemodia maritima, Streptoprocne zonaris (Shaw 1796), Sturnella magna, Sublegatus arenarum (Salvin 1863), Synallaxis candei Orbigny & Lafresnaye 1838, Tabernaemontana amygdalifolia, Tachycineta albiventer (Boddaert 1783), Tachycineta bicolor (Vieillot 1808), Talinum paniculatum, Tapera naevia (Linnaeus 1766), Tephrosia cinerea, Thamnophilus doliatus (Linnaeus 1764), Thraupis glaucocolpa Cabanis 1850, Thraupis palmarum, Thryothorus leucotis (Lafresnaye 1845), Tiaris bicolor (Linnaeus 1766), Tiaris obscurus (Orbigny & Lafresnaye 1837), Tillandsia flexuosa, Tityra inquisitor (Lichtenstein 1823), Tolmomyias sulphurescens (Spix 1825), Tournefortia hirsutissima, Trianthema portulacastrum, Tribulus cistoides, Tyrannus dominicensis (Gmelin 1788), Tyto alba, Vireo flavoviridis (Cassin 1851), Vitex compressa, Wissadula periplocifolia, Zenaida auriculata

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

habitatUnstructured: Poco común a medianamente común en suelo o en marañas en montes bajos, áridos y deciduos y áreas de matorral. Arisco e inconspicuo (Hilty & Brown 1986). Esta especie prefiere los hábitats xerofíticos y subxerofíticos de La Guajira (Cerrejón 2010).

endemicAtomized

distribution

ancillaryData

dataObject:

additionalInformation:

reference

link: <http://collections.mnh.si.edu/search/>

keywords: Amazona ochrocephala, Ameiva festiva, Anas discors, Anhinga anhinga (Linnaeus 1766), Aramides cajanea, Aramus guarauna, Aratinga acuticaudata, Ardea herodias, Arremonops conirostris, Arremonops tocuyensis, Atalotriccus pilaris, Attila spadiceus (Gmelin 1789), Basileuterus flaveolus (S. F. Baird 1865), Bubo virginianus (Gmelin 1788), Burhinus bistriatus (Wagler 1829), Campylorhynchus trochilirostris (Lichtenstein 1820), Campylorhynchus griseus (Swainson 1838), Campylorhynchus nuchalis (Cabanis 1847), Caprimulgus cayennensis (Gmelin 1789), Cardinalis phoeniceus Bonaparte 1838, Carduelis psaltria (Say 1823), Carolia brevicauda (Schinz 1821), Centurio senex Gray 1842, Certhia cinnamomeus (Gmelin 1788), Chlorostilbon gibsoni (Fraser 1840), Chordeiles acutipennis (Hermann 1783), Chrysolampis mosquitos (Linnaeus 1758), Chrysomus icterocephalus, Coccyzus pumila, Colibri coruscans, Columbina minuta (Linnaeus 1766), Columbina passerina (Linnaeus 1758), Conirostrum leucogenys (Lafresnaye 1852), Coryphospingus pileatus (Wied-Neuwied 1821), Crotophaga major, Crotophaga sulcirostris Swainson 1827, Cyanocorax affinis (Pelzeln 1856), Cyclarhis gujanensis (J. F. Gmelin 1789), Dendrocygna autumnalis, Dryocopus lineatus (Linnaeus 1766), Egretta thula, Egretta tricolor (Statius Muller 1776), Elaenia chiriquensis Lawrence 1865, Engystomops pustulosus, Eumops glaucinus (Wagner 1843), Fluvicola pica (Boddaert 1783), Forpus passerinus (Linnaeus 1758), Furnarius leucopus Swainson 1838, Galbula ruficauda Cuvier 1816, Gallinula chloropus (Linnaeus 1758), Glaucidium brasilianum (Gmelin 1788), Glossophaga longirostris Miller 1898, Gonatodes vittatus, Heliodoxa leadbeateri (Bourcier 1843), Hemitriccus margaritaceiventer (Orbigny & Lafresna, Himantopus mexicanus (Statius Muller 1776), Hylophilus flavipes Lafresnaye 1845, Icterus auricapillus Cassin 1848, Icterus galbula (Linnaeus 1758), Icterus nigrogularis (Hahn 1819), Inezia subflava (P. L. Sclater & Salvin 1873), Inezia tenuirostris (Cory 1913), Lasiurus ega (Gervais 1856), Lepidoblepharis sanctaemartae Ruthven 1916, Lepidopyga goudoti (Bourcier 1843), Leptonycteris curasoae Miller 1900, Leptotila verreauxi (Bonaparte 1855), Leucippus fallax (Bourcier 1843), Lophostoma silviculum (d'Orbigny 1836), Machetornis rixosa, Macrophyllum macrophyllum (Schinz 1821), Megaceryle alcyon (Linnaeus 1758), Megaceryle torquata (Linnaeus 1766), Megarynchus pitangua (Linnaeus 1766), Megascops choliba, Melanerpes rubricapillus, Micronycteris hirsuta (Peters 1869), Molothrus bonariensis, Mormoops megalophylla (Peters 1864), Mycteria americana Linnaeus 1758, Myiarchus crinitus, Myiarchus venezuelensis Lawrence 1865, Myiopagis gaimardii (Orbigny 1840), Myiozetetes similis (Spix 1825), Myospiza humeralis (Bosc 1792), Nemosia pileata (Boddaert 1783), Noctilio albiventris (Desmarest 1818), Nycticorax nycticorax, Ortalis ruficauda Jardine 1847, Pachyramphus rufus (Boddaert 1783), Patagioenas corensis (Jacquin 1784), Phaeomyias murina (Spix 1825), Pheucticus ludovicianus, Piculus chrysochloros (Vieillot 1818), Picumnus cinnamomeus Wagler 1829, Pilherodius pileatus (Boddaert 1783), Pionus sordidus (Linnaeus 1758), Piranga flava (Vieillot 1822), Platalea ajaja Linnaeus 1758, Pleurodema brachyops, Polioptila plumbea (J. F. Gmelin 1788), Pseudopaludicola pusilla, Pteronotus davyi (Gray 1838), Pteronotus gymnonotus (Natterer 1843), Pteronotus parnellii (Gray 1843), Pteronotus personatus (Wagner 1843), Pulsatrix perspicillata (Latham 1790), Pyrocephalus rubinus, Quiscalus lugubris, Quiscalus mexicanus (Gmelin 1788), Sakesphorus canadensis (Linnaeus 1766), Saltator coerulescens Vieillot 1817, Spiza americana, Sporophila bouvronides (Lesson 1831), Sporophila intermedia Cabanis 1851, Sporophila nigricollis (Vieillot 1823), Stelgidopteryx ruficollis (Vieillot 1817), Streptoprocne zonaris (Shaw 1796), Sturnella magna, Sublegatus arenarum (Salvin 1863), Tachycineta albiventer (Boddaert 1783), Tachycineta bicolor (Vieillot 1808), Tapera naevia (Linnaeus 1766), Thamnophilus doliatus (Linnaeus 1764), Thraupis glaucocolpa Cabanis 1850, Thraupis palmarum, Thryothorus leucotis (Lafresnaye 1845), Tolmomyias sulphurescens (Spix 1825), Tyrannus dominicensis (Gmelin 1788), Tyto alba, Zenaida auriculata

editors

institution:

address:
publisher:
accessed:
websites:
chapter:
series:
issue:
year: 2012

authors

authors: Smithsonian National Museum of Natural History

title: Search Museum Collections Records

source: SMITHSONIAN NATIONAL MUSEUM OF NATURAL HISTORY

type: miscellany

tags: Amazona ochrocephala, Ameiva festiva, Anas discors, Anhinga anhinga (Linnaeus 1766), Aramides cajanea, Aramus guarauna, Aratinga acuticaudata, Ardea herodias, Arremonops conirostris, Arremonops tocuyensis, Atalotriccus pilaris, Attila spadiceus (Gmelin 1789), Basileuterus flaveolus (S. F. Baird 1865), Bubo virginianus (Gmelin 1788), Burhinus bistriatus (Wagler 1829), Campylorhynchus trochilirostris (Lichtenstein 1820), Campylorhynchus griseus (Swainson 1838), Campylorhynchus nuchalis (Cabanis 1847), Caprimulgus cayennensis (Gmelin 1789), Cardinalis phoeniceus Bonaparte 1838, Carduelis psaltria (Say 1823), Carollia brevicauda (Schinz 1821), Centurio senex Gray 1842, Certhiaxis cinnamomeus (Gmelin 1788), Chlorostilbon gibsoni (Fraser 1840), Chordeiles acutipennis (Hermann 1783), Chrysolampis mosquitus (Linnaeus 1758), Chrysomus icterocephalus, Coccyzus pumila, Colibri coruscans, Columbina minuta (Linnaeus 1766), Columbina passerina (Linnaeus 1758), Conirostrum leucogenys (Lafresnaye 1852), Coryphospingus pileatus (Wied-Neuwied 1821), Crotophaga major, Crotophaga sulcirostris Swainson 1827, Cyanocorax affinis (Pelzelin 1856), Cyclarhis gujanensis (J. F. Gmelin 1789), Dendrocoryna autumnalis, Dryocopus lineatus (Linnaeus 1766), Egretta thula, Egretta tricolor (Statius Muller 1776), Elaenia chiriquensis Lawrence 1865, Engystomops pustulosus, Eumops glaucinus (Wagner 1843), Fluvicola pica (Boddaert 1783), Forpus passerinus (Linnaeus 1758), Furnarius leucopus Swainson 1838, Galbula ruficauda Cuvier 1816, Gallinula chloropus (Linnaeus 1758), Glaucidium brasilianum (Gmelin 1788), Glossophaga longirostris Miller 1898, Gonatodes vittatus, Heliodoxa leadbeateri (Bourcier 1843), Hemitriccus margaritaceiventer (Orbigny & Lafresna, Himantopus mexicanus (Statius Muller 1776), Hylophilus flavipes Lafresnaye 1845, Icterus auricapillus Cassin 1848, Icterus galbula (Linnaeus 1758), Icterus nigrogularis (Hahn 1819), Inezia subflava (P. L. Sclater & Salvin 1873), Inezia tenuirostris (Cory 1913), Lasiurus ega (Gervais 1856), Lepidoblepharis sanctaemartae Ruthven 1916, Lepidopyga goudoti (Bourcier 1843), Leptonycteris curasoae Miller 1900, Leptotila verreauxi (Bonaparte 1855), Leucippus fallax (Bourcier 1843), Lophostoma silviculum (d'Orbigny 1836), Mactornis rixosa, Macrophyllum macrophyllum (Schinz 1821), Megaceryle alcyon (Linnaeus 1758), Megaceryle torquata (Linnaeus 1766), Megarynchus pitangua (Linnaeus 1766), Megascops choliba, Melanerpes rubricapillus, Micronycteris hirsuta (Peters 1869), Molothrus bonariensis, Mormoops megalophylla (Peters 1864), Mycteria americana Linnaeus 1758, Myiarchus crinitus, Myiarchus venezuelensis Lawrence 1865, Myiopagis gaimardii (Orbigny 1840), Myiozetetes similis (Spix 1825), Myospiza humeralis (Bosc 1792), Nemosia pileata (Boddaert 1783), Noctilio albiventris (Desmarest 1818), Nycticorax nycticorax, Ortalis ruficauda Jardine 1847, Pachyrhamphus rufus (Boddaert 1783), Patagioenas corensis (Jacquin 1784), Phaeomyias murina (Spix 1825), Pheucticus ludovicianus, Piculus chrysochloros (Vieillot 1818), Picumnus cinnamomeus Wagler 1829, Pilherodius pileatus (Boddaert 1783), Pionus sordidus (Linnaeus 1758), Piranga flava (Vieillot 1822), Platalea ajaja Linnaeus 1758, Pleurodema brachyops, Polioptila plumbea (J. F. Gmelin 1788), Pseudopaludicola pusilla, Pteronotus davyi (Gray 1838), Pteronotus gymnotus (Natterer 1843), Pteronotus parnellii (Gray 1843), Pteronotus personatus (Wagner 1843), Pulsatrix perspicillata (Latham 1790), Pyrocephalus rubinus, Quiscalus lugubris, Quiscalus mexicanus (Gmelin 1788), Sakesphorus canadensis (Linnaeus 1766), Saltator coerulescens Vieillot 1817, Spiza americana, Sporophila bouvronides (Lesson 1831), Sporophila intermedia Cabanis 1851, Sporophila nigricollis (Vieillot 1823), Stelgidopteryx ruficollis (Vieillot 1817), Streptoprocne zonaris (Shaw 1796), Sturnella magna, Sublegatus arenarum (Salvin 1863), Tachycineta albiventer (Boddaert 1783), Tachycineta bicolor (Vieillot 1808), Tapera naevia (Linnaeus 1766), Thamnophilus doliatus (Linnaeus 1764), Thraupis glaucocolpa Cabanis 1850, Thraupis palmarum, Thryothorus leucotis (Lafresnaye 1845), Tolmomyias sulphurescens (Spix 1825), Tyrannus dominicensis (Gmelin 1788), Tyto alba, Zenaida auriculata

abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:
distributionUnstructured: Se encuentra en alturas hasta 200 m (Hilty & Brown 1986).En bases de datos y monitoreos de fauna se encontró que:
Esta especie se distribuye en: La Guajira (Cerrejón 2010, Collections MNH 2012, ICN 2004), Magdalena (Collections MNH 2012). Su región natural es la región Caribe de Colombia. A nivel mundial en bases de datos se encontró que:¿
Esta especie se distribuye en: Colombia (Collections MNH 2012).

distributionAtomized

stateProvince: La Guajira

ancillaryData

locality:
municipality:
country: Colombia
temporalCoverage
endDate:
startDate:
distributionScope
type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

0

link: <http://www.iucnredlist.org>

keywords: Arremonops tocuyensis

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2012

0: BirdLife International

title: Arremonops tocuyensis. En: IUCN 2012. IUCN Red List of Threatened Species. Versión 2012.2

source:

type: miscellany

tags: Arremonops tocuyensis

abstract:

last_modified:

created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
populationBiologyUnstructured: El tamaño de la población mundial no ha sido cuantificado, pero esta especie se describe como "bastante común" (Birdlife International 2012).

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

abstract: Mide unos 13 cm. Cabeza y nuca gris pálido con una lista central negra en la coronilla y una superciliar, que se prolonga hasta la nuca de color arcilla blanquecina. El resto de las partes dorsales color verde oliva; por debajo de color gris claro (Hilty & Brown 1986).

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: edgarestebanhc@gmail.com
phone:
postalCode:
country:
state:
city:
address:
organisation:
position:
lastName: Herrera Collazos
firstName: Edgar Esteban

ancillaryData

dataObject:
additionalInformation: Registros biológicos:
Portal de datos GBIF

reference

geoPoint:
location:
thumbnailURL:
mediaURL: <http://data.gbif.org/species/2491041/>
description:

subject

source: <http://data.gbif.org/>

audience

bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id

_bsontype: ObjectID
id: Vç~WfÄW Dä