

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d4da303c16479905cba8d3

Fecha de creación: 2016-02-29T23:54:26.397Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

habitatUnstructured: Terrestre, bosques, montes secundarios y áreas intervenidas muy abiertas. Una de las aves más comunes y conspicuas en Colombia en terreno abierto o semiabierto con árboles, en áreas residenciales y en claros y orillas de ríos en zonas selváticas.

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Español

name: Tirano tropical

creation\_date: 2016-02-29T23:54:26.397Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataFormat:

identifier:

fullDescriptionUnstructured: Largo total promedio 22 cm. Cabeza gris con mascara negruzca; parche naranja oculto en la coronilla; espalda oliva grisáceo; alas y cola ligeramente ahorquillada café negruzco; garganta gris pálido; bajas partes inferiores amarillas con fuerte lavado oliva en el pecho.

briefDescription:

[references](#)

taxonRecordName

[ancillaryData](#)

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

replacementNameFor

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

conservedAgainst

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:  
datatype:

identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion  
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:  
  
publishedIn  
  
simple:  
source:  
datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: Vieillot, 1819  
  
canonicalName  
  
infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: Tyrannus melancholicus  
rank: SPECIES  
simple: Tyrannus melancholicus Vieillot, 1819

attributes

nomenclaturalCode:

isAnamorphic:

id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-02-29T23:54:06.394Z

created: 2016-02-29T23:54:06.394Z

[agent](#)

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataType:

identifier:

parentTaxon: Tyrannus

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Tyrannus  
family: Tyrannidae  
order: Passeriformes  
classHierarchy: Aves  
phylum: Chordata  
kingdom: Animalia  
recommended:  
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Clara Maria Pérez Gallego  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
reproductionUnstructured: El nido es una taza frágil usualmente apoyada en horqueta a 5 - 15 m; 2 - 3 huevos blancos crema manchados y rayados café rojizo.

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:  
additionalInformation:

geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Clara Maria Pérez Gallego  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
feedingUnstructured: Se alimenta principalmente de insectos.

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

behaviorUnstructured: Diurno y hacia los estratos medios y bajos de bosques, rastrojos y áreas abiertas. Se posa conspicuo en copa de árboles, alambres y otras perchas expuestas, usualmente a media altura. Vuela en pos de insectos a casi cualquier altura, ocasionalmente al suelo o al agua; a veces persigue insectos con ágiles maniobras. Notablemente pugnaz. Voz con una llamada usual con un trino agudo trí"í"í"í"ip . Canto al amanecer, más elaborado, es una serie de pip seguida por un trino ascendente. Más bien vocal y uno de los primeros cantos en la mañana.

ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:  
additionalInformation:

[reference](#)

geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:  
rightsHolder:  
rights: Clara Maria Pérez Gallego  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:



created:

[agent](#)

mimeType:

dataType:

identifier:

distributionUnstructured: S México hasta C Argentina y Trinidad. Ampla distribución en todo el país y sus regiones naturales hasta los 2700 msnm.

[distributionAtomized](#)

temporalCoverage

endDate:

startDate:

distributionScope

type:

[threatStatus](#)

[ancillaryData](#)

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

[measurementDeterminedBy](#)

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract:

[associatedParty](#)

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

[ancillaryData](#)

dataObject:

additionalInformation: Registros biológicos: En la Central Hidroeléctrica Jaguas se registró en los bosques de La Arenosa, Santa Ana, Almenara, El Edén, en los rastrojos Papayos, Puente Cable, Vivero, Frente Papayos, San Lorenzo, Helipuerto-Fuga y en las áreas intervenidas de Embalse, Vertedero, Compuertas y Zona Baja. En la Central Hidroeléctrica San Carlos se registró en los bosques Quebrada la Llore, Cerro de las Torres y Almenara, en los rastrojos Portería Presa y Central San Carlos y en las áreas intervenidas de Vertedero, Casino Viejo, Vía Domo y Embalse.

Imagen: Tyrannus\_melancholicus.jpg

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:  
rightsHolder: Clara Maria Pérez Gallego  
rights:  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

agent

mimeType:  
dataType:  
identifier:

\_id  
\_bsontype: ObjectID  
id: VÔÛ0< G™\º•0