Reporte de Fichas para el usuario: edith gómez
Ficha ID: 56e1948b83c45700544e3d17 Fecha de creación: 2016-03-10T15:36:50.366Z
annualCycles
annualCycleUnstructured:
habitats
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: habitatUnstructured: La especie habita bosques húmedos de Sudamérica.
commonNamesAtomized
ancillaryData
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
source:
audience
bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
agent

mime i ype:
dataType:
identifier:
usedBy:
usedIn
distribution Instructured

distributionUnstructured:

endDate: startDate:

### ancillaryData

type:

language: Español name: Avispa

creation\_date: 2016-03-10T15:36:50.366Z usesManagementAndConservation

# ancillaryData

managementAndConservationUnstructured:

# managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint: location: thumbnailURL: description: source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: Costado anterodorsal de la propleura doblado hacia afuera en toda su longitud.

briefDescription:

### references

keywords: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus

metanotalis,Clypearia sulcata,Clypearia weyrauchi,Epipona tatua,Leipomeles dorsata,Metapolybia acincta,Metapolybia docilis,Metapolybia nigra,Metapolybia suffusa,Mischocyttarus adolphi,Mischocyttarus alfkenii excrucians,Mischocyttarus alfkenii trinitatis,Mischocyttarus carbonarius carbonarius,Mischocyttarus flavicans,Mischocyttarus foveatus,Mischocyttarus injucundus tingomariae,Mischocyttarus insolitus,Mischocyttarus labiatus,Mischocyttarus latissimus,Mischocyttarus metoecus,Mischocyttarus petiolatus,Mischocyttarus pseudomimeticus,Mischocyttarus punctatus,Mischocyttarus tomentosus,Mischocyttarus vaqueroi,Parachartergus amazonensis,Parachartergus colobopterus,Parachartergus fasciipennis,Parachartergus fraternus,Parachartergus fulgidipennis,Parachartergus griseus,Parachartergus richardsi,Parachartergus weyrauchi,Polybia bistriata,Polybia catillifex,Polybia eberhardae,Polybia emaciata,Polybia fastidiosuscula,Polybia jurinei,Polybia liliacea,Polybia micans,Polybia occidentalis,Polybia platycephala sylvestris,Polybia quadricincta,Polybia raui raui,Polybia rejecta,Polybia scrobalis pronotalis,Polybia scrobalis scrobalis,Polybia sericea,Polybia striata,Polybia velutina,Protopolybia acutiscutis,Protopolybia amarella,Protopolybia chartergoides chartergoides,Protopolybia exigua binominata,Protopolybia exigua exigua,Protopolybia fuscatus,Protopolybia rugulosa,Protopolybia sedula,Pseudopolybia compressa,Pseudopolybia vespiceps,Synoeca septentrionalis,Synoeca surinama,Synoeca virginea

### editors

institution: address: publisher: accessed: websites: chapter: series: pages: 585 issue: year: 1978

### authors

authors: Richards, O. W.

title: The social wasps of the Americas excluding the Vespinae

source: type: book

tags: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta. Metapolybia docilis. Metapolybia nigra. Metapolybia suffusa. Mischocyttarus adolphi. Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea abstract:

### identifiers

last\_modified: created: group\_id: profile\_id:

id.

taxonRecordName

# ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

# replacementNameFor

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

### sanctioned

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

# laterHomonymOf

microReference:

source: datatype: identifier:

datatype: identifier: note: ruleConsidered:
conservedAgainst
basedOn
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
basionym
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: spellingCorrentionOf
typeNameEntity
source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:
typificacion
typeOfType:
lectotypeMicroReferenceVoucher
lectotypePublicationVoucher

voucherReference

simple: microReference:

source:

year:
publishedIn
simple: source: datatype: identifier:
specialAuthorship
authors
year
simple:
canonicalAuthorship
authors
year
simple: simple: (de Saussure, 1853)
canonicalName
infraspecificEpithet: specificEpithet: infragenericEpithet: linkType: ref: uninomial: simple: Clypearia sulcata rank: SPECIES simple: Clypearia sulcata (de Saussure, 1853)
attributes
nomenclaturalCode: isAnamorphic: id:
synonymsAtomized
hierarchy
ancillaryData
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:

### mediaURL

description:

# subject

source: The Catalogue of Life, 3rd January 2011

### audience

bibliographicCitation:

rigthsHolder:

rights: license: title:

modified: 2016-03-10T15:36:30.364Z created: 2016-03-10T15:36:30.364Z

# agent

homepage:

role: email: phone: address: position: organisation: lastName: firstName: mimeType: dataType: identifier:

parentTaxon: Clypearia higherClassification: infraspecificEpithet: specificEpithet: taxonRank: SPECIES

subGenus: genus: Clypearia family: Vespidae order: Hymenoptera classHierarchy: Insecta phylum: Arthropoda kingdom: Animalia recommended: classification:

invasiveness

invasivenessUnstructured:

reproduction

dataObject:

additionalInformation:

geoPoint: location:

thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured: 0
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
audience
rightsHolder: rights: license: modified: created:
agent
mimeType: dataType: identifier: relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy: TraitBank measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: Sexual measurementType: Tipo de Reproducción measurementID:
molecularData
molecularDataUnstructured:
migratory

migratoryUnstructured: lifeForm lifeFormUnstructured: lifeCycle lifeCycleUnstructured: interactions interactionsUnstructured: feeding dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: feedingUnstructured: 0 strategy: Depredadores type: Heterótrofos - Omnívoros environmentalEnvelope environmentalEnvelopeUnstructured: ecologicalSignificance ecologicalSignificanceUnstructured: dispersal dispersalUnstructured: distance: structureDispersed: type: purpose: behavior

dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: behaviorUnstructured: Son insectos depredadores con habitos sociales. Y se les encuentra anidando en árboles habitados por hormigas del género Azteca. Cuando los adultos emergen de la celda ésta es limpiada y reutilizada. Dentro de un mismo nido puede haber variación en la coloración de los individuos.
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID:

# distribution

endemicAtomized

# ancillaryData

dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
source:

### audience

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

### agent

mimeType: dataType: identifier:

distributionUnstructured: Esta especie se ha registrado en Colombia, Ecuador, Venezuela, Guyana, Guyana Francesa y la región Norte de Brasil. En Colombia se distribuye en la región de la Amazonia.

### distributionAtomized

stateProvince: Putumayo

# ancillaryData

locality: municipality: country: Colombia

temporalCoverage

endDate: startDate:

distributionScope

type:

# threatStatus

### territory

territoryUnstructured: areaOfOccupancy: extentOfOccurrence:

populationBiology

populationBiologyUnstructured:
legislation
legislationUnstructured:
directThreats
directThreatsUnstructured:
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID: abstract: Margen frontal lamelado del pronoto delgado; segmento abdominal I en forma de copa; dorso torácico no aplanado; metanoto con el área basal estrecha y horizontal, su área posterior es vertical abruptamente.
associatedParty
role: personnelIdentifier: personnelDirectory: homepage: email: enataliagomezm@gmail.com phone: postalCode: country: Colombia state: Bogotá Distrito Capital city: Bogotá D.C. address: organisation: position: lastName: Gómez Melendro

firstName: Edith Natalia

### ancillaryData

dataObject:

additionalInformation: Información de tipos: Holotipo hembra coleccionado en Brasil y depositado en el British Museum of Natural History (BMNH), Londres, Reino Unido.

Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

### reference

geoPoint:
location:
thumbnailURL:

### mediaURL

description:

### subject

source:

### audience

bibliographicCitation:

rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

### agent

mimeType: dataType: identifier:

\_id

\_bsontype: ObjectID id: Vá" f ÄW DãÑp