

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e4a54b83c45700544e3e44

Fecha de creación: 2016-03-12T23:25:07.371Z

creation_date: 2016-03-12T23:25:07.371Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: En estas avispa el escutelo se ve, de perfil, curvado regularmente formando con el metanoto una única inclinación o convexidad y nunca se encuentra elevado y proyectado sobre el metanoto, el margen posterior del metanoto se prolonga hacia el área medio-basal del propodeo como un lóbulo o ángulo agudo; el segmento abdominal I es a menudo peciolado.

briefDescription:

[references](#)

keywords: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca trilli, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala

sylvestris, Polybia quadricincta, Polybia raii, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea

editors

institution:
address:
publisher:
accessed:
websites:
chapter:
series:
pages: 585
issue:
year: 1978

authors

authors: Richards, O. W.
title: The social wasps of the Americas excluding the Vespinae
source:
type: book

tags: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca trilli, Brachygastera albula, Brachygastera augusti, Brachygastera lecheguana, Brachygastera propodealis, Brachygastera scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raii, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: Duce, 1907
canonicalName
infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Protopolybia rugulosa
rank: SPECIES
simple: Protopolybia rugulosa Duce, 1907

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rightsHolder:
rights:
license:
title:
modified: 2016-03-12T23:24:47.370Z
created: 2016-03-12T23:24:47.370Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Protopolybia
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Protopolybia
family: Vespidae
order: Hymenoptera
classHierarchy: Insecta
phylum: Arthropoda
kingdom: Animalia
recommended:
classification:

commonNameAtomized

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:
created:
mimeType:
dataType:
identifier:
reproductionUnstructured:

0

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder:
rights:
license:
modified:
created:

agent

mimeType:
dataType:
identifier:
relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedBy: TraitBank
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue: Sexual
measurementType: Tipo de Reproducción
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

data Type:

identifier:

feedingUnstructured:

0

0

strategy: Depredadores

type:

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
behaviorUnstructured: Protopolybia rugulosa al igual que las demás especies del género Protopolybia exhibe hábitos predadores y eusociales.

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

annualCycle

annualCycleUnstructured:

habitat

habitatUnstructured:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:

additionalInformation:

reference

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

mimeType:

dataType:

identifier:

distributionUnstructured: Esta especie se distribuye en Colombia, Surinám, Guyana Francesa y Brasil. En el país se ha reportado en la región Amazónica.

distributionAtomized

stateProvince: Amazonas

ancillaryData

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Esta especie depredadora de la familia Vespidae se distribuye en Sudamérica. Se caracteriza por tener el clípeo 1,5 veces más largo que ancho; especie negra con algunas manchas amarillas en la mitad del escutelo y en los segmentos abdominales I-II.

[associatedParty](#)

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomez@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Información de tipos: Lectotipo: hembra proveniente de Brasil, y depositado en el British Museum of Natural History (BMNH), Londres, Reino Unido.
Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaría General de la CAN y el Instituto Sinchi para el proyecto “Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC”.

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

mimeType:
dataType:
identifier:

_id
_bsontype: ObjectID
id: Vã¥KfÄW Dää@