

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e33eda83c45700544e3dbe

Fecha de creación: 2016-03-11T21:55:44.827Z

creation_date: 2016-03-11T21:55:44.827Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: Cabeza: labio amarillo blanquecino; labro y frente anteriormente amarillo claro; anteclypeo y postclypeo amarillo claro con manchas verdosas; frente lateralmente y por encima gris verdoso. Tórax: lateralmente con bandas marrón oscuro, y verdosas; dorsalmente marrón oscuro. Patas ocre y espinas negras. Abdomen café claro con manchas verdosas; apéndices ocre mate, apéndices superiores más cortos y más curvos que en otras especies. Alas del macho con una banda marrón proximal al triángulo, venación rojiza; Todos los triángulos y subtriángulos libres, es decir sin venas transversas. Hembra con patrón de coloración similar al macho, pero con las bandas del tórax blanco azulosas en vez de verdosas.

briefDescription:

[references](#)

keywords: Argia cupraurea, Argia pulla Hagen in, Cora chiribiquete, Cora inca, Erythemis vesiculosa, Erythrodiplax kimminsi, Erythrodiplax umbrata, Erythrodiplax unimaculata, Mecistogaster linearis, Orthemis discolor, Perissolestes magdalenae, Perithemis lais, Perithemis mooma, Polythore beata, Polythore concinna, Polythore derivata, Polythore mutata, Telebasis corallina, Uracis fastigiata, Uracis imbuta

[editors](#)

institution:

address:

publisher: Instituto Nacional de Biodiversidad, Ministerio del Ambiente y Energía, Universidad Nacional de Costa Rica, Banco Mundial, GEF

accessed:

websites:

chapter:
series:
issue:
year: 2006

authors

authors: Esquivel, C
title: Libélulas de Mesoamérica y el Caribe. Dragonflies and damselflies of Middle America and the Caribbean
source:
type: book
tags: Argia cupraurea,Argia pulla Hagen in,Cora chiribiquete,Cora inca,Erythemis vesiculosa,Erythrodiplax kimminsi,Erythrodiplax umbrata,Erythrodiplax unimaculata,Mecistogaster linearis,Orthemis discolor,Perissolestes magdalenae,Perithemis lais,Perithemis mooma,Polythore beata,Polythore concinna,Polythore derivata,Polythore mutata,Telebasis corallina,Uracis fastigiata,Uracis imbuta
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

laterHomonymOf

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

basionym

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:

datatype:

identifier:

source:

datatype:

identifier:

source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:

canonicalAuthorship

authors

year

simple:
simple: Perty, 1834

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Perithemis lais
rank: SPECIES
simple: Perithemis lais Perty, 1834

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

ancillaryData

synonymStatus:

synonymName

anyTwo:

anyOne

providerLink:

microReference:

publishedIn

source:

datatype:

identifier:

reletedName

source:

datatype:

identifier:

note:

ruleConsidered:

microReference:

publishedIn

source:

datatype:

identifier:

reletedName

source:

datatype:

identifier:

note:

ruleConsidered:

microReference:

publishedIn

source:

datatype:

identifier:

reletedName

source:

datatype:

identifier:

note:

ruleConsidered:

microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:

datatype:
identifier:

combinationAuthorship

basionymAuthorship

simple:

authorship

simple:

simple: Ris, 1910

epithet

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

genus

linkType:

ref:

uninomial:

simple: Perithemis naias

rank: SPECIES

simple: Perithemis naias Ris, 1910

nomenclaturalCode:

isAnamorphic:

id:

hierarchy

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-11T21:55:24.826Z

created: 2016-03-11T21:55:24.826Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Perithemis
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Perithemis
family: Libellulidae
order: Odonata
classHierarchy: Insecta
phylum: Arthropoda
kingdom: Animalia
recommended:
classification:

commonNameAtomized

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:

created:

agent

contentType:

dataType:

identifier:

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

ancillaryData

type:

language: Español

name: Libélula

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

reproductionUnstructured: Cuando una hembra entra en el territorio del macho, éste la intercepta vibrando sus alas más rápido y levantando el abdomen, luego vuela hacia un sitio adecuado para la oviposición, si la hembra accede se posa y allí el macho la atrapa y comienza la cópula. La cópula es muy corta, mientras la hembra pone los huevos, el macho vuela muy cerca de ella. Para poner los huevos la hembra vuela sobre pequeños troncos que sobresalen del agua o sobre la vegetación flotante, y toca el sitio con la punta del abdomen repetidamente para adherir los huevos al sustrato.

0

dataObject:

additionalInformation:

reference

geoPoint:

location:
thumbnailURL:

mediaURL

description:
subject

audience

rightsHolder:
rights:
license:
modified:
created:

agent

contentType:
dataType:
identifier:
relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedBy: TraitBank
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue: Sexual
measurementType: Tipo de Reproducción
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:

description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured:

0

0

strategy: Depredadores
type:

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:
structureDispersed:
type:
purpose:

behavior

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
behaviorUnstructured: Los machos son territoriales y se posan muy cerca del agua, en la punta de ramitas de la vegetación emergente, y con el cuerpo y las alas completamente horizontales. Con frecuencia los machos se buscan para pelear o perseguirse, formando cadenas de persecución compuestas por machos de territorios

adyacentes. Son unos de los pocos anisópteros en los que el macho corteja a la hembra antes de la cópula.

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

annualCycle

annualCycleUnstructured:

habitat

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
habitatUnstructured: Esta especie habita lagunas o charcas en áreas abiertas

endemicAtomized

distribution

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:
distributionUnstructured: En el país se ha registrado en las regiones de la Amazonia, Caribe, Orinoquía y Pacífica.

distributionAtomized

ancillaryData

locality:
municipality:
country: México

temporalCoverage

endDate:
startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured:
areaOfOccupancy:
extentOfOccurrence:

populationBiology
populationBiologyUnstructured:

legislation
legislationUnstructured:

directThreats
directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

abstract: Perithemis lais es una especie de libélula depredadora que se distribuye en Centroamérica y Sudamérica. Se caracterizan por ser unos de los pocos anisópteros en los que el macho corteja a la hembra antes de la cópula.

[associatedParty](#)

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital

city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".
Registros biológicos: Colección Entomológica, Instituto de Ciencias Naturales, Universidad Nacional de Colombia. Bogotá, Colombia <http://www.biovirtual.unal.edu.co/colecciones/datos/animalia/consultasMammalia.jsp?clase=4>

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id

_bsontype: ObjectID
id: Vã>UfÄW DãÛà