Reporte de Fichas para el usuario: edith gómez Ficha ID: 56e0dc9483c45700544e3cf2 Fecha de creación: 2016-03-10T02:31:54.513Z annualCycles annualCycleUnstructured: habitats 0 dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: habitatUnstructured: Esta especie habita en bosques secos tropicales de Centroamérica y Sudamérica. commonNamesAtomized creation_date: 2016-03-10T02:31:54.513Z usesManagementAndConservation ancillaryData managementAndConservationUnstructured: managementAndConservationAtomized 0 dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType:

dataType: identifier: useNotes: En algunas regiones de México y sur de Sudamérica esta especie se usa como fuente de miel. potential: properties: ratingPopularity: economics: useTypeAtomized: conservationExplotationData: seasonOfAvailabilityUse: meansOfApplicationAdministration: productionDetails: vernacularNameUseAnnotations: organisms: users: partUsed: useValue: sourceOfInformationText:
references
identificationKeys
0
dataObject: additionalInformation:
0
keywords: Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Brachygastra albula,Brachygastra augusti,Brachygastra lecheguana institution: Universidad de Costa Rica address: publisher: accessed: websites: chapter: series: issue: year: 2010
0: Valverde, J.P. title: Clave taxonómica ilustrada para los géneros y las especies de avispas eusociales de Costa Rica (Hymenoptera: Vespidae, Polistinae)
source: type: thesis tags: Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Brachygastra albula,Brachygastra augusti,Brachygastra lecheguana abstract:
last_modified:
created: group_id:
profile_id:
id:
geoPoint:

location:
thumbnailURL:
description:
SOURCE.

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

keys: Valverde, J. P. (2010). Clave taxonómica ilustrada para los géneros y las especies de avispas eusociales de Costa Rica (Hymenoptera: Vespidae, Polistinae). Universidad de Costa Rica.

fullDescription

0

dataObject:

additionalInformation:

geoPoint: location: thumbnailURL: description: source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: Visto de perfil el escutelo es prominente con dos superficies y proyectándose sobre el metanoto y el segmento abdominal I es corto y estrecho pero no en forma de copa.

Escutelo con un ángulo marcado entre las superficies anterior y posterior; metanoto liso; si hay punteaduras, están en el margen dorsal, manchas amarillas en los tergos abdominales II-V; en vista dorsal escutelo emarginado posteriormente.

briefDescription:

references

keywords: Agelaia angulata angulata,Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Asteloeca traili,Brachygastra albula,Brachygastra augusti,Brachygastra lecheguana,Brachygastra propodealis,Brachygastra scutellaris,Chartergellus amazonicus,Chartergellus punctatior,Chartergus chartarius,Chartergus globiventris,Chartergus metanotalis,Clypearia sulcata,Clypearia weyrauchi,Epipona tatua,Leipomeles dorsata,Metapolybia acincta,Metapolybia docilis,Metapolybia nigra,Metapolybia suffusa,Mischocyttarus adolphi,Mischocyttarus alfkeni,Mischocyttarus alfkenii excrucians,Mischocyttarus alfkenii trinitatis,Mischocyttarus carbonarius carbonarius,Mischocyttarus flavicans,Mischocyttarus foveatus,Mischocyttarus injucundus tingomariae,Mischocyttarus insolitus,Mischocyttarus labiatus,Mischocyttarus latissimus,Mischocyttarus metoecus,Mischocyttarus petiolatus,Mischocyttarus pseudomimeticus,Mischocyttarus punctatus,Mischocyttarus tomentosus,Mischocyttarus vaqueroi,Parachartergus amazonensis,Parachartergus colobopterus,Parachartergus

fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea

editors

institution: address: publisher: accessed: websites: chapter: series: pages: 585 issue: year: 1978

authors

authors: Richards, O. W.

title: The social wasps of the Americas excluding the Vespinae

source: type: book

tags: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus wevrauchi, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea abstract:

identifiers

last_modified: created: group_id: profile_id: id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

replacementNameFor

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

sanctioned

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

laterHomonymOf

microReference:

source:
datatype:
identifier:
source:
datatype:
identifier:
note:

ruleConsidered:

conservedAgainst

basedOn microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: basionym microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: spellingCorrentionOf typeNameEntity source: datatype: identifier: source: datatype: identifier: source: datatype: identifier: typificacion typeOfType: lectotypeMicroReferenceVoucher **lectotypePublicationVoucher** voucherReference simple: microReference: year: publishedIn simple: source: datatype: identifier:

specialAuthorship

authors

year

simple:

canonicalAuthorship

authors

year

simple:

simple: (Latreille, 1824)

canonicalName

infraspecificEpithet: specificEpithet: infragenericEpithet:

linkType: ref: uninomial:

simple: Brachygastra lecheguana

rank: SPECIES

simple: Brachygastra lecheguana (Latreille, 1824)

attributes

nomenclaturalCode: isAnamorphic:

id:

synonymsAtomized

ancillaryData

synonymStatus:

synonymName

anyTwo:

anyOne

providerLink: microReference:

publishedIn

source: datatype: identifier:

reletedName

source: datatype:

identifier: note: ruleConsidered: microReference:
publishedIn
source: datatype: identifier:
reletedName
source: datatype: identifier: note:

ruleConsidered: microReference:

publishedIn

source: datatype: identifier:

reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

publishedIn

source: datatype: identifier:

reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

publishedIn

source: datatype: identifier:

reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:
publishedIn
source: datatype: identifier:
reletedName
source: datatype: identifier: note: ruleConsidered:
typeNameEntity
source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:
typeVoucherEntity
typeOfType: simple: microReference: year: source: datatype: identifier:
$combination \\ Authorship$
basionymAuthorship
simple:
authorship
simple: simple: Perty, 1833
epithet
infraspecificEpithet: specificEpithet: infragenericEpithet:
genus
linkType: ref:
uninomial: simple: Brachygastra analis

rank: SPECIES

simple: Brachygastra analis Perty, 1833

nomenclaturalCode: isAnamorphic:

id:

hierarchy

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rigthsHolder: rights: license: title:

modified: 2016-03-10T02:31:34.511Z created: 2016-03-10T02:31:34.511Z

agent

homepage:

role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:

parentTaxon: Brachygastra

higherClassification: infraspecificEpithet: specificEpithet: taxonRank: SPECIES

subGenus:

genus: Brachygastra family: Vespidae

order: Hymenoptera classHierarchy: Insecta phylum: Arthropoda kingdom: Animalia recommended: classification: invasiveness invasivenessUnstructured: reproduction dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured: dataObject: additionalInformation: reference geoPoint: location: thumbnailURL: mediaURL description: subject audience rightsHolder: rights: license: modified: created:

agent

mimeType: dataType: identifier: relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy: TraitBank measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: Sexual measurementType: Tipo de Reproducción measurementID: molecularData molecularDataUnstructured: migratory migratoryUnstructured: lifeForm lifeFormUnstructured: lifeCycle lifeCycleUnstructured: interactions interactionsUnstructured: feeding dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: feedingUnstructured: Estas avispas almacenan grandes cantidades de néctar en sus nidos; visita algunas plantas alucinógenas y algodonales en busca de néctar y presas. 0

n

strategy: Depredadores type: Heterótrofos - Omnívoros
environmentalEnvelope
environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured: distance: structureDispersed: type: purpose:
behavior
0
dataObject: additionalInformation:
0
keywords: Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Brachygastra albula,Brachygastra augusti,Brachygastra lecheguana institution: Universidad de Costa Rica address: publisher: accessed: websites: chapter: series: issue: year: 2010 0: Valverde, J.P.
title: Clave taxonómica ilustrada para los géneros y las especies de avispas eusociales de Costa Rica (Hymenoptera: Vespidae, Polistinae) source:
type: thesis tags: Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Brachygastra albula,Brachygastra augusti,Brachygastra lecheguana abstract:
last_modified: created: group_id: profile_id: id: geoPoint: location: thumbnailURL:
description: source:

bibliographicCitation: rightsHolder: rights: Equipo Coordinador del SiB Colombia license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: behaviorUnstructured: Construyen sus nidos con cada panal cubierto por una envoltura y fusionado a la envoltura del panal precedente (Valverde, 2010), adicionalmente, este presenta cualidades tóxicas en algunas épocas del año.
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID:
endemicAtomized
distribution
ancillaryData
dataObject: additionalInformation:
reference
geoPoint:

location: thumbnailURL:

mediaURL description: subject source: audience bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: agent mimeType: dataType: identifier: distributionUnstructured: Se distribuye en Centroamérica (Costa Rica y Panamá) y en Sudamérica (Colombia, Venezuela, Guyana, Guyana Francesa, Surinam, Ecuador, Perú, Paraguay, Uruguay, Brasil, Bolivia y el norte de Argentina) en áreas de hasta 1500 msnm de altura. En Colombia se ha registrado a lo largo del país desde La Guajira hasta el Amazonas. distributionAtomized stateProvince: Córdoba ancillaryData locality: municipality: country: Colombia temporalCoverage endDate: startDate: distributionScope type: threatStatus territory territoryUnstructured: areaOfOccupancy: extentOfOccurrence: populationBiology populationBiologyUnstructured: legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType:
identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID: abstract: Esta especie se distribuye en bosques secos tropicales de Centro y Sudamérica. Se caracteriza por presentar un escutelo en vista dorsal usualmente emarginado posteriormente y el digitus del macho obtuso distalmente en vista lateral (Valverde, 2010).
associatedParty
role: personnelIdentifier: personnelDirectory: homepage: email: enataliagomezm@gmail.com phone: postalCode: country: Colombia state: Bogotá Distrito Capital city: Bogotá D.C. address:

ancillaryData

lastName: Gómez Melendro firstName: Edith Natalia

organisation: position:

dataObject: additionalInformation: Información de tipos: El holotipo es una hembra coleccionada en Brasil que fue depositada en

el Musée National d'histoire Naturelle, Paris, Francia (MNHN) pero aparentemente hoy esta extraviada. Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation: rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

agent

mimeType: dataType: identifier:

id

_bsontype: ObjectID id: VàÜ"fÄW DãÏ