

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c7483ef0106c67230e73e7

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dubusia taeniata,Egretta caerulea,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreiventris

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dubusia taeniata,Egretta

caerulea,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreiventris

abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:

rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:
created:
mimeType:
dataType:
identifier:

habitatUnstructured: En áreas rurales la especie prefiere elevaciones mayores y con más vegetación natural que Eriocnemis vestita. Se desplaza en el nivel medio y bajo de los matorrales nativos y subpáramo donde visita las flores del uvo de monte (ericaeas), agarrándose de estas para libar. Prefiere sitios abiertos pero también se encuentra en el dosel del bosque (Asociación Bogotana de Ornitología, 2000).

commonNamesAtomized

ancillaryData

usedBy:
usedIn
distributionUnstructured:
endDate:
startDate:

ancillaryData

type:
language: Inglés
name: Coppery-bellied Puffleg
usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

fullDescriptionUnstructured: Mide 10 cm. Es similar al macho de *Eriocnemis vestita* , pero un poco más grande y con el pico relativamente más corto, cola más larga y los pompones aún más grandes y llamativos (Asociación Bogotana de Ornitología, 2000). Los adultos presentan una coloración general verde brillante oscuro; un parche cobrizo en el centro del abdomen (visible con buena luz), más grande y brillante en el macho. Las coberteras infracaudales azul- violeta brillante, cola negro azulado y ahorquillada. Picos y patas negras (Asociación Bogotana de Ornitología, 2000).

briefDescription:

references

link: Neotropical Birds Online http://neotropical.birds.cornell.edu/portal/species/overview?p{_}p{_}spp=268536

keywords: *Eriocnemis cupreovertris*

editors

editors: Schulenberg, T S

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2011

authors

authors: Sedgwick, C W

title: Coppery-bellied Puffleg (*Eriocnemis cupreovertris*)

source:

type: miscellany

tags: *Eriocnemis cupreovertris*

abstract:

identifiers

last_modified:

created:

group_id:

profile_id:

id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Fraser, 1840)
canonicalName
infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Eriocnemis cupreovertris
rank: SPECIES
simple: Eriocnemis cupreovertris (Fraser, 1840)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rightsHolder:
rights:
license:
title:
modified: 2016-02-19T16:45:14.344Z
created: 2016-02-19T16:45:14.344Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Eriocnemis
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Eriocnemis
family: Trochilidae
order: Apodiformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:

0

link: <http://www.hbw.com/node/55589>
keywords: Eriocnemis cupreiventris
0: del Hoyo, J
1: Elliott, A
2: Sargatal, J
3: Christie, D A
4: de Juana, E
institution:
address:

publisher: Lynx Edicions
accessed:
websites:
chapter:
series:
issue:
year: 1999
0: Heynen, I
1: Boesman, P
title: Coppery-bellied Puffleg (*Eriocnemis cupreovertris*)
source: Handbook of the Birds of the World Alive
type: book_section
tags: *Eriocnemis cupreovertris*
abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
contentType:
dataType:
identifier:
reproductionUnstructured: Se reproduce entre septiembre y enero. El nido es a menudo situado en la vegetación densa. Pone dos huevos que son incubados por la hembra (Heynen & Boesman, 1999).
molecularData
molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions
interactionsUnstructured:
feeding

dataObject:
additionalInformation:

0

link: <http://www.hbw.com/node/55589>

keywords: *Eriocnemis cupreovertris*

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 1999

0: Heynen, I

1: Boesman, P

title: Coppery-bellied Puffleg (*Eriocnemis cupreovertris*)

source: Handbook of the Birds of the World Alive

type: book_section

tags: *Eriocnemis cupreovertris*

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

feedingUnstructured: Consume principalmente néctar que extrae de las plantas con flores de corolas cortas como *Palicourea*, *Cavendishia* y *Pernettya*. También caza insectos desde sus sitios de percha (Heynen & Boesman, 1999).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus

fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dubusia taeniata,Egretta caerulea,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera

ensifera,Eriocnemis cupreiventris

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataFormat:

identifier:

behaviorUnstructured: Defiende territorios en parches ricos en flores de ericáceas en matorrales y subpáramo (Asociación Bogotana de Ornitología, 2000).

La especie tiene un reclamo fuerte y agudo pit o tsip emitido una o varias veces. No canta. (Asociación Bogotana de Ornitología, 2000).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataFormat:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:
measurementID:

endemicAtomized

distribution

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES: Apéndice II

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Casi amenazado (NT)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

0

link: <http://www.birdlife.org/datazone/species/factsheet/22687854>

keywords: Eriocnemis cupreovertris

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:
year: 2014
0: BirdLife International
title: Species factsheet: Eriocnemis cupreovertris
source:
type: miscellany
tags: Eriocnemis cupreovertris
abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
populationBiologyUnstructured: El tamaño de la población mundial no se ha cuantificado, pero esta especie es descrita como poco común a rara (BirdLife International, 2014).

legislation

legislationUnstructured:

directThreats

0

dataObject:
additionalInformation:

0

link: <http://www.birdlife.org/datazone/species/factsheet/22687854>
keywords: Eriocnemis cupreovertris
institution:
address:
publisher:
accessed:
websites:
chapter:
series:
issue:
year: 2014
0: BirdLife International
title: Species factsheet: Eriocnemis cupreovertris
source:
type: miscellany

tags: Eriocnemis cupreovertris

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

directThreatsUnstructured: Su hábitat en los Andes al este de Colombia ha sido ampliamente degradado en gran parte para establecer cultivos y pastos. En Mérida y Táchira (Venezuela), la deforestación a nivel local es severa debido principalmente a la colonización agrícola, con muchas áreas de páramo y bosque ya taladas para el cultivo comercial y de subsistencia (BirdLife International, 2014).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: El Pomponero Vientricobrizo es muy activo y agresivo. Son muy territoriales. Por lo general son solitarios y

se alimentan de las flores tubulares en arbustos bajos o pequeños árboles. Habita los bordes de los bosques montanos, laderas arbustivas, páramo abierto, matorrales y bosques achaparrados (Sedgwick, 2011).

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:
mediaURL: <http://www.xeno-canto.org/56044>
description:

subject

source: <http://www.xeno-canto.org>

audience

bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:

agent

contentType:
dataType:
identifier:

_id

_bsontype: ObjectId
id: VÇH>đ lg#ç>p