

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e0639e83c45700544e3cbf
Fecha de creación: 2016-03-09T17:55:48.105Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
habitatUnstructured: Habita en bosques húmedos tropicales de Centroamérica y Sudamérica.

[commonNamesAtomized](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

[agent](#)

contentType:
dataFormat:
identifier:
usedBy:

usedIn

distributionUnstructured:
endDate:
startDate:

ancillaryData

type:
language: Español
name: Avispa
creation_date: 2016-03-09T17:55:48.105Z

usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:
managementAndConservationAtomized

identificationKeys

0

dataObject:
additionalInformation:

0

keywords: Agelaia angulicollis,Agelaia cajennensis
institution: Universidad de Costa Rica
address:
publisher:
accessed:
websites:
chapter:
series:
issue:
year: 2010
0: Valverde, J.P.
title: Clave taxonómica ilustrada para los géneros y las especies de avispas eusociales de Costa Rica (Hymenoptera: Vespidae, Polistinae)
source:
type: thesis
tags: Agelaia angulicollis,Agelaia cajennensis
abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:

location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

keys: Valverde, J. P. (2010). Clave taxonómica ilustrada para los géneros y las especies de avispas eusociales de Costa Rica (Hymenoptera: Vespidae, Polistinae). Universidad de Costa Rica.

fullDescription

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

fullDescriptionUnstructured: Tórax y lóbulos laterales del clípeo, pálidos; clípeo en contacto con los ojos; espacio malar con un área brillante indistinta; parastigma el doble de largo que de ancho. Placa dorsal del mesepisterno mucho más estrecha y la sutura forma un ángulo con la parte anterior por debajo de los 60°. Escutelo con una línea central muy notoria, algunas veces superficial; mesoescuto con un margen continuo desde atrás hasta la parte anterior de la tégula.

briefDescription:

references

keywords: *Agelaia angulata angulata*, *Agelaia angulicollis*, *Agelaia brevistigma*, *Agelaia cajennensis*, *Agelaia fulvofasciata*, *Agelaia hamiltoni*, *Agelaia testacea*, *Angiopolybia obidensis*, *Angiopolybia pallens*, *Angiopolybia paraensis*, *Angiopolybia zischkai*, *Apoica thoracica*, *Asteloeca trilli*, *Brachygastra albula*, *Brachygastra augusti*, *Brachygastra lecheguana*, *Brachygastra propodealis*, *Brachygastra scutellaris*, *Chartergellus amazonicus*, *Chartergellus punctatior*, *Chartergus chartarius*, *Chartergus globiventris*, *Chartergus metanotalis*, *Clypearia sulcata*, *Clypearia weyrauchi*, *Epipona tatua*, *Leipomeles dorsata*, *Metapolybia acincta*, *Metapolybia docilis*, *Metapolybia nigra*, *Metapolybia suffusa*, *Mischocyttarus adolphi*, *Mischocyttarus alfkeni*, *Mischocyttarus alfkenii excrucians*, *Mischocyttarus alfkenii trinitatis*, *Mischocyttarus carbonarius carbonarius*, *Mischocyttarus flavicans*, *Mischocyttarus foveatus*, *Mischocyttarus injucundus tingomariae*, *Mischocyttarus insolitus*, *Mischocyttarus labiatus*, *Mischocyttarus latissimus*, *Mischocyttarus metoecus*, *Mischocyttarus petiolatus*, *Mischocyttarus pseudomimeticus*, *Mischocyttarus punctatus*, *Mischocyttarus tomentosus*, *Mischocyttarus vaqueroi*, *Parachartergus amazonensis*, *Parachartergus colobopterus*, *Parachartergus*

fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raii, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea

editors

institution:
address:
publisher:
accessed:
websites:
chapter:
series:
pages: 585
issue:
year: 1978

authors

authors: Richards, O. W.
title: The social wasps of the Americas excluding the Vespinae
source:
type: book

tags: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca trilli, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raii, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

[ancillaryData](#)

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

replacementNameFor

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

sanctioned

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

laterHomonymOf

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

basionym

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:

datatype:

identifier:

source:

datatype:

identifier:

source:

datatype:

identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:

microReference:

year:

publishedIn

simple:

source:

datatype:

identifier:

specialAuthorship

authors

year

simple:

canonicalAuthorship

authors

year

simple:

simple: (Fabricius, 1798)

canonicalName

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

linkType:

ref:

uninomial:

simple: Agelaia cajennensis

rank: SPECIES

simple: Agelaia cajennensis (Fabricius, 1798)

attributes

nomenclaturalCode:

isAnamorphic:

id:

synonymsAtomized

ancillaryData

synonymStatus:

synonymName

anyTwo:

anyOne

providerLink:

microReference:

publishedIn

source:

datatype:

identifier:

reletedName

source:

datatype:

identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:

ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:
datatype:
identifier:

combinationAuthorship

basionymAuthorship

simple:

authorship

simple:
simple: Ducke, 1904

epithet

infraspecificEpithet:
specificEpithet:
infragenericEpithet:

genus

linkType:
ref:
uninomial:
simple: Polybia lignicola

rank: SPECIES
simple: Polybia lignicola Ducke, 1904
nomenclaturalCode:
isAnamorphic:
id:

[hierarchy](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-03-09T17:55:28.105Z
created: 2016-03-09T17:55:28.105Z

[agent](#)

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Agelaia
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Agelaia
family: Vespidae

order: Hymenoptera
classHierarchy: Insecta
phylum: Arthropoda
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:

0

keywords: Agelaia angulicollis,Agelaia cajennensis
institution: Universidad de Costa Rica
address:
publisher:
accessed:
websites:
chapter:
series:
issue:

year: 2010

0: Valverde, J.P.

title: Clave taxonómica ilustrada para los géneros y las especies de avispas eusociales de Costa Rica
(Hymenoptera: Vespidae, Polistinae)

source:

type: thesis

tags: Agelaia angulicollis,Agelaia cajennensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

reproductionUnstructured: Construyen sus nidos en cavidades, que consiste de muchos panales apilados unidos por pedicelos, o un panal con envoltura, en la rama de un árbol (Valverde, 2010).

0

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder:
rights:
license:
modified:
created:

agent

contentType:
dataType:
identifier:
relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedBy: TraitBank
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue: Sexual
measurementType: Tipo de Reproducción
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:

0

keywords: Agelaia angulicollis,Agelaia cajennensis

institution: Universidad de Costa Rica

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2010

0: Valverde, J.P.

title: Clave taxonómica ilustrada para los géneros y las especies de avispa eusociales de Costa Rica
(Hymenoptera: Vespidae, Polistinae)

source:

type: thesis

tags: Agelaia angulicollis,Agelaia cajennensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

feedingUnstructured: La dieta en Polistinae se compone de dos elementos, néctar y presas (Valverde, 2010).

0

0

strategy: Depredadores

type: Heterótrofos - Omnívoros

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

behaviorUnstructured: Son insectos eusociales y de hábitos depredadores. Sus nidos se encuentran entre hojas de bromelias epífitas, en oquedades de los árboles o de construcciones.

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataObject:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:
measurementType:
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

[agent](#)

contentType:
dataType:
identifier:
distributionUnstructured: Se ha registrado en México, Honduras, Belice, Costa Rica, Panamá, Colombia, Ecuador, Perú, Guyana, Guyana Francesa, Trinidad y Tobago, Surinam, Brasil y Bolivia; en rangos altitudinales de hasta 500 msnm. En Colombia se distribuye en las regiones de la Orinoquia, Amazonia y Pacífica.

[distributionAtomized](#)

stateProvince: Amazonas

[ancillaryData](#)

locality:
municipality:
country: Colombia

temporalCoverage

endDate:
startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

mimeType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Se distribuye desde Centroamérica hasta Sudamérica en áreas con altitudes de hasta 500 msnm y hábitats de bosque húmedo tropical. Esta especie de avista presenta un margen lateral lamelado del pronoto muy estrecho y el ápice de la pterostigma redondeado, sin una emarginación (Valverde, 2010).

associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:
email: enataliagomez@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Información de tipos: El lectotipo está depositado en el Musée National d'histoire Naturelle, Paris, Francia (MNHN).
Créditos específicos: " Â -ç7F—GWFò Ö `6æ—6ò FR —çfW7F—v 6—öæW2 6—VçNÖf—6 2 6—æ6†'â Â 6—7FVÖ FR —æf÷:&Ö 6Ÿ6â Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id

_bsontype: ObjectId
id: VàcžfÄW DãËö