

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d4b4873c16479905cba8b3

Fecha de creación: 2016-02-29T21:13:45.428Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Esta especie se encuentra en áreas intervenidas (Hilty & Brown 2001). Es común en terrenos abiertos y secos, granjas, prados, áreas pobladas (Hilty & Brown 2001).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Alemán

name: Rosttäubchen

creation_date: 2016-02-29T21:13:45.428Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: C. talpacoti tiene un largo total promedio de 16.5cm. El macho es principalmente Rufo canela, con las partes inferiores más pálidas y la coronilla contrastante gris clara, la hembra es mucho más opaca especialmente por debajo y con la espalda rojiza (Hilty & Brown 2001).

briefDescription:

[references](#)

taxonRecordName

[ancillaryData](#)

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

replacementNameFor

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:

identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Temminck, 1810)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Columbina talpacoti
rank: SPECIES
simple: Columbina talpacoti (Temminck, 1810)

attributes

nomenclaturalCode:

isAnamorphic:

id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-02-29T21:13:25.427Z

created: 2016-02-29T21:13:25.427Z

[agent](#)

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataType:

identifier:

parentTaxon: Columbina

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Columbina
family: Columbidae
order: Columbiformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
reproductionUnstructured: Está especie hace un nido frágil y somero a 1m o más, con dos huevos (Hilty & Brown 2001).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:

geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured: La Tortolita rojiza se alimenta principalmente de semillas (Hilty & Brown 2001).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

behaviorUnstructured: Es una especie diurna (Hilty & Brown 2001). "W2 Vâ fR æ÷F &ÆVÖVçFR 6ö6– &ÆR ' 6öæf– &ÆRâ
Permanece en pareja o grupos de diferentes tamaños. Camina mecánicamente con pasos cortos y rápidos. En
querellas a menudo amenaza levantando un ala (Hilty & Brown 2001).

La Tortolita rojiza emite un sonido de ca juu ca-juu hueco y monótono, generalmente canta desde percha elevada
(Hilty & Brown 2001).

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:

created:

agent

contentType:

dataType:

identifier:

distributionUnstructured: Esta especie se encuentra desde el Sur México (ocasionalmente Sur de Texas) hasta el Norte de Argentina, Paraguay y Uruguay; accidental en Chile; noroccidente de Perú. Trinidad (Hilty & Brown 2001). En el país se distribuye en todo el centro del país y zonas secas (Hilty & Brown 2001), principalmente en la región del caribe y amazónica, encontrándose también en la región andina (Hilty & Brown 2001). Está especie en Colombia se encuentra hasta 2600m.s.n.m. (Hilty & Brown 2001).

distributionAtomized

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract:

[associatedParty](#)

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

[ancillaryData](#)

dataObject:
additionalInformation: Imagen: Columbina_talpacoti.jpg
Registros biológicos: En la Central Hidroeléctrica la Miel se registro esta especie en el bosque el Tigre, la Mula y Campiña, en los rastrojos de Reten Norcasia y el Encanto y en las áreas intervenidas de Cantera, Terraplen.

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder: Clara Maria Pérez Gallego
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id
_bsonType: ObjectID
id: VÔ´†< G™\0<0