

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d5f14a3c16479905cba904

Fecha de creación: 2016-03-01T19:45:16.277Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Es común en matorrales áridos y semiáridos, monte seco, áreas cultivadas y manglares (Hilty & Brown 1986). En La Guajira habita en los bosques deciduos tropical, de galería y secundario y en matorrales o rastrojos (Cerrejón 2009-2010).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Inglés

name: Red-crowned Woodpecker

creation\_date: 2016-03-01T19:45:16.277Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
fullDescriptionUnstructured: Se reconocen cuatro subespecies:  
Melanerpes rubricapillus paraguanae (Gilliard, 1940)  
Melanerpes rubricapillus rubricapillus (Cabanis, 1862)  
Melanerpes rubricapillus seductus Bangs, 1901  
Melanerpes rubricapillus subfuscus (Wetmore, 1957)  
briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:

datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

conservedAgainst

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:

publishedIn

simple:  
source:  
datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: (Cabanis, 1862)

canonicalName

infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:

ref:  
uninomial:  
simple: Melanerpes rubricapillus  
rank: SPECIES

simple: *Melanerpes rubricapillus* (Cabanis, 1862)

## attributes

nomenclaturalCode:

isAnamorphic:

id:

## synonymsAtomized

## hierarchy

## ancillaryData

dataObject:

additionalInformation:

## reference

geoPoint:

location:

thumbnailURL:

## mediaURL

description:

## subject

source: The Catalogue of Life, 3rd January 2011

## audience

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-01T19:44:56.274Z

created: 2016-03-01T19:44:56.274Z

## agent

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

mimeType:

dataType:

identifier:

parentTaxon: *Melanerpes*

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Melanerpes

family: Picidae

order: Piciformes

classHierarchy: Aves

phylum: Chordata

kingdom: Animalia

recommended:

classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

feedingUnstructured: A menudo come fruta (Hilty & Brown 1986). En La Guajira se observó que su dieta se compone principalmente de invertebrados y que forrajea en el dosel (Cerrejón 2009-2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

behaviorUnstructured: Es conspicuo en áreas abiertas, se le observa solitario o en parejas, a altura media o inferior. Cincela la superficie de la corteza, inspecciona extremos de ramas rotas (Hilty & Brown 1986). Es un ave ruidosa; krrr-r-r-r fuerte y arrastrado y uica fuerte si está excitado. Tamborilea principalmente al inicio de la temporada de reproducción (Skutch 1969 en Hilty & Brown 1986).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

mimeType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:

dataType:

identifier:

distributionUnstructured: En bases de datos se encontró que:

Esta especie se distribuye en: Costa Rica, México, Venezuela (Arctos 2012)

Las siguientes subespecies se distribuyen en:

Melanerpes rubricapillus costaricensis Costa Rica (Arctos 2012)

Melanerpes rubricapillus paraguanae Colombia (Collections MNH SI 2012)

Melanerpes rubricapillus rubricapillus Colombia (Arctos 2012; Collections MNH SI 2012), Panamá (Arctos 2012).

En bases de datos y monitoreos de fauna se encontró que:

Las siguientes subespecies se distribuyen en Colombia:

Melanerpes rubricapillus paraguanae Antioquia, La Guajira (ICN 2004); Bolívar, Cesar, La Guajira, Magdalena (Collections MNH SI 2012)



Melanerpes rubricapillus rubricapillus Atlántico, Bolívar, Caldas, Cesar, Córdoba, Cundinamarca, Huila, Magdalena, Norte de Santander, Santander, Sucre, Tolima (ICN 2004); Antioquia, Atlántico, Bolívar, Magdalena, Norte de Santander, Santander (Collections MNH SI 2012); Huila, Magdalena, Santander (Arctos 2012). En las regiones Andina y Caribe, hasta los 1700 m (Hilty & Brown 1986).

distributionAtomized

stateProvince: Córdoba

ancillaryData

locality:  
municipality:  
country: Colombia

temporalCoverage

endDate:  
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:  
locality:  
municipality:  
county:  
stateProvince:  
country: Global  
0: IUCN  
measurementValue: Preocupación menor (LC)  
relatedTo:  
measurementRemarks:  
measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementType: IUCN  
measurementID:

territory

territoryUnstructured:  
areaOfOccupancy:  
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Mide unos 18 cm. Los machos tienen la frente blanco amarillenta; coronilla y occipucio rojos; el resto de las partes dorsales barradas negro y blanco, rabadilla blanca; los lados de la cabeza hasta arriba de los ojos y las partes ventrales de color gris anteadado, el centro del vientre rojo. La hembra tiene la coronilla blanco anteadado, sólo el occipucio rojo (Hilty & Brown 1986).

[associatedParty](#)

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

ancillaryData

dataObject:  
additionalInformation: Registros biológicos: En el portal de Global Biodiversity Information Facility se reporta que para Colombia hay 752 registros. GBIF  
Créditos específicos: Ficha elaborada con el apoyo de Carbones del Cerrejón Limited.  
Metadatos: GBIF š  
Imagen: Melanerpes\_rubricapillus\_foto.jpg  
Otros recursos en Internet: Arctos, Avibase, Colecciones en Línea, EoL, iSpecies, IUCN Red List, SI NMNH, xenocanto

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

agent

mimeType:  
dataType:  
identifier:

\_id  
\_bsontype: ObjectID  
id: VÖñJ< G™\0”