

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e1befc83c45700544e3d36

Fecha de creación: 2016-03-10T18:37:55.125Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

habitatUnstructured: Habitan en bosques de galería de Sudamérica.

[commonNamesAtomized](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:
dataFormat:
identifier:
usedBy:

usedIn

distributionUnstructured:
endDate:
startDate:

ancillaryData

type:
language: Español
name: Abejita estacional amazónica
creation_date: 2016-03-10T18:37:55.125Z

usesManagementAndConservation

ancillaryData

dataObject:
additionalInformation:

reference

keywords: Eufriesea auripes,Eufriesea dressleri,Exaerete frontalis

editors

editors: Amat-G, G.
editors: Andrade-C, M. G.
editors: Amat G, E.
institution:
address:
publisher: Instituto de Ciencias Naturales-Universidad Nacional de Colombia, Conservación Internacional Colombia, Instituto Alexander von Humboldt, Ministerio de Ambiente, Vivienda y Desarrollo Territorial
accessed:
websites:
chapter:
series:
pages: 160--163
issue:
year: 2007

authors

authors: Bonilla, M A
title: Abejita estacional amazónica (Eufriesea auripes)
source: Libro Rojo de los Invertebrados Terrestres de Colombia
type: book_section
tags: Eufriesea auripes,Eufriesea dressleri,Exaerete frontalis
abstract:

identifiers

last_modified:
created:
group_id:

profile_id:
id:
geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

[agent](#)

contentType:
dataType:
identifier:
managementAndConservationUnstructured: Para Bonilla (2007) es fundamental hacer un muestreo intensivo de las comunidades de abejas euglosinas en diferentes áreas del país con el ánimo de establecer los tamaños actuales de las poblaciones, particularidades de su distribución espacial y otros aspectos ecológicos relevantes como sus fuentes de néctar, sus sitios de nidificación, la estructura de sus nidos y sus fuentes naturales de fragancias.

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

fullDescriptionUnstructured:
briefDescription:

[references](#)

keywords: Eufriesea auripes

[editors](#)

institution: Universidade Estadual de Campinas
address:
publisher:
accessed:
websites:
chapter:
series:
pages: 153
issue:
year: 1999

[authors](#)

authors: Bonilla-Gómez, M. A.
title: Caracteriza\cc\~ao da estrutura espa\cco-temporal da comunidades de hábelas Euglossinas (Hymenoptera, Apidae) na Hiléia Baiana
source:
type: thesis
tags: Eufriesea auripes
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:

ruleConsidered:

laterHomonymOf

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

basionym

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:

datatype:

identifier:

source:

datatype:

identifier:

source:

datatype:

identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Gribodo, 1882)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Eufriesea auripes
rank: SPECIES
simple: Eufriesea auripes (Gribodo, 1882)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-03-10T18:37:35.123Z
created: 2016-03-10T18:37:35.123Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Eufriesea
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Eufriesea
family: Apidae
order: Hymenoptera
classHierarchy: Insecta
phylum: Arthropoda
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
reproductionUnstructured:

0

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder:
rights:
license:
modified:
created:

agent

mimeType:
dataType:
identifier:
relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedBy: TraitBank
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue: Sexual
measurementType: Tipo de Reproducción

measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataFormat:

identifier:

feedingUnstructured: Se desconocen sus fuentes naturales de néctar y de fragancias.

0

type: Heterótrofos - Herbívoros

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

behaviorUnstructured: La mayoría de las especies del género *Eufriesea* son altamente estacionales; los adultos aparecen únicamente durante la estación más lluviosa del año y pasan la mayor parte del tiempo en un periodo de inactividad en el estadio prepupal. En Colombia, los machos de esta especie han sido colectados en el mes de octubre. En otros lugares del Neotrópico aparecen entre marzo y enero. Son una especie de abejas solitarias.

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataObject:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

[endemicAtomized](#)

distribution

ancillaryData

dataObject:

additionalInformation:

reference

keywords: Eufriesea auripes,Eufriesea dressleri,Exaerete frontalis

editors

editors: Amat-G, G.

editors: Andrade-C, M. G.

editors: Amat G, E.

institution:

address:

publisher: Instituto de Ciencias Naturales-Universidad Nacional de Colombia, Conservación Internacional Colombia, Instituto Alexander von Humboldt, Ministerio de Ambiente, Vivienda y Desarrollo Territorial

accessed:

websites:

chapter:

series:

pages: 160--163

issue:

year: 2007

authors

authors: Bonilla, M A

title: Abejita estacional amazónica (Eufriesea auripes)

source: Libro Rojo de los Invertebrados Terrestres de Colombia

type: book_section

tags: Eufriesea auripes,Eufriesea dressleri,Exaerete frontalis

abstract:

identifiers

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataFormat:
identifier:
distributionUnstructured: Se distribuye en Brasil, Guyana Francesa, Guyana, Perú, Surinam, Venezuela y Colombia, en áreas entre los 580 msnm y 700 msnm. En el país En Colombia, el único registro corresponde a los bosques de galería de San Juan de Arama en el Meta (Bonilla, 2007).

distributionAtomized

stateProvince: Meta

ancillaryData

locality:
municipality:
country: Colombia
temporalCoverage
endDate:
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:
threatStatusAtomized
apendiceCITES:
locality:
municipality:
county:
stateProvince:
country: Colombia
0: IUCN
measurementValue: En peligro (EN)
relatedTo:
measurementRemarks:
measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementType: IUCN
measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

0

dataObject:

additionalInformation:

0

keywords: Eufriesea auripes

institution: Universidade Estadual de Campinas

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 153

issue:

year: 1999

0: Bonilla-Gómez, M. A.

title: Caracteriza\cc\~ao da estrutura espa\cco-temporal da comunidades de hábelas Euglossinas (Hymenoptera, Apidae) na Hiléia Baiana

source:

type: thesis

tags: Eufriesea auripes

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

directThreatsUnstructured: La principal amenaza que enfrenta esta especie es la fragmentación y destrucción de los bosques de galería cercanos a la Serranía de la Macarena, área en donde ha sido encontrada. Esta destrucción representa la destrucción misma de sus poblaciones ya que por pasar la mayor parte del año en un periodo prepupal y de inactividad no tienen la oportunidad de migrar a otros hábitats cuando sus lugares de nidificación son perturbados. Estudios sugieren que la alta estacionalidad y la distribución espacial muy restringida del género *Eufriesea* muestran una alta vulnerabilidad a la extinción local de sus especies frente a procesos de degradación de los hábitats (Bonilla, 1999).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Son abejas solitarias que habitan en bosques de galería de Sudamérica. Es una especie altamente estacional; los adultos aparecen únicamente durante la estación más lluviosa del año.

[associatedParty](#)

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomez@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Información de tipos: El holotipo está en el Museo Civico di Storia Naturale "Giacomo Doria" (MSNG), Génova, Italia.
Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaría General de la CAN y el Instituto Sinchi para el proyecto “Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC”.

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

mimeType:
dataType:
identifier:

_id
_bsontype: ObjectID
id: Vá%üfÄW DãÓ`