

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c7c24bf0106c67230e7417

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigrirostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Cistothorus platensis*, *Coccyzua pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta caerulea*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxypogon guerinii*, *Oxyura jamaicensis*, *Pachyramphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Playa cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

abstract:

last_modified:

created:

group_id:
profile_id:
id:

1

keywords: Accipiter striatus, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Anas andium, Andigena nigrirostris, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Catamenia inornata, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geranoaetus melanoleucus, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis

institution:

address:

publisher: Christopher Helm

accessed:

websites:

chapter:

series:

pages: 880

issue:

year: 2006

0: Restall, R

1: Rodner, C

2: Lentino, M

title: Birds of Northern South America: An Identification Guide, Volume 1: Species Accounts.

source:

type: book

tags: Accipiter striatus, Aglaiocercus kingii, Amblycercus holosericeus, Anas andium, Andigena nigrirostris, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Catamenia inornata, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geranoaetus melanoleucus, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:
mimeType:
dataType:
identifier:
habitatUnstructured: El hábitat de la especie está en el páramo, ya que son exclusivos de zonas altas caracterizadas por arbustos dispersos, bosques enanos y frailejonales. También hace uso de matorrales y bordes de bosques húmedos (Hilty & Brown, 1986; Restall et al., 2006).

commonNamesAtomized

ancillaryData

usedBy:
usedIn
distributionUnstructured:
endDate:
startDate:

ancillaryData

type:
language: Alemán
name: Bergameisenpitta
usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:
additionalInformation:

0

keywords: Accipiter striatus,Actitis macularius,Adelomyia melanogenys,Agelaiocercus kingii,Amblycercus holosericeus,Ampelion rubrocristatus,Anas andium,Anas clypeata,Anas discors,Andigena nigrirostris,Anisognathus igniventris,Ardea alba,Ardea herodias,Arremon assimilis,Asio flammeus,Asio stygius,Asthenes flammulata,Asthenes fuliginosa,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Butorides virescens,Cairina moschata,Calidris bairdii,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Cathartes aura,Catharus minimus,Catharus ustulatus,Chaetocercus mulsant,Chaetura pelagica,Chalcostigma heteropogon,Chordeiles minor,Chrysomus icterocephalus,Cinclodes albidiventris,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Cistothorus platensis,Coccyzua pumila,Coccyzus americanus,Coccyzus melacoryphus,Coeligena bonapartei,Colibri coruscans,Colibri thalassinus,Columba livia,Columbina talpacoti,Conirostrum rufum,Conirostrum sitticolor,Contopus cinereus,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Dendrocygna autumnalis,Dendrocygna viduata,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dolichonyx oryzivorus,Dubusia taeniata,Egretta caerulea,Egretta

thula, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus superciliaris*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxypogon guerinii*, *Oxyura jamaicensis*, *Pachyramphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Aglaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigrirostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Cistothorus platensis*, *Coccyzua pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta caerulea*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera*

ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: La especie llega a medir hasta 18 cm. Es oliva opaco o marrón encima, en la cara presenta una gran mancha loreal y anillo ocular estrecho blanco mientras que los lados de la cabeza y las partes inferiores son ocre pálido con efecto moteado blancuzco. No existe dimorfismo sexual, pero los juveniles si son fácilmente diferenciables de los adultos por presentar puntos y barras negras en la corona, la nuca y el pecho (Hilty & Brown, 1986).

briefDescription:

references

keywords: Accipiter striatus, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Anas

andium, Andigena nigristrois, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Catamenia inornata, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geranoaetus melanoleucus, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis

editors

institution:
address:
publisher: Christopher Helm
accessed:
websites:
chapter:
series:
pages: 880
issue:
year: 2006

authors

authors: Restall, R
authors: Rodner, C
authors: Lentino, M
title: Birds of Northern South America: An Identification Guide, Volume 1: Species Accounts.
source:

type: book

tags: Accipiter striatus, Aglaiocercus kingii, Amblycercus holosericeus, Anas andium, Andigena nigristrois, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Catamenia inornata, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geranoaetus melanoleucus, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

replacementNameFor

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

sanctioned

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

laterHomonymOf

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

conservedAgainst

basedOn

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: Lesson, 1844

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Grallaria quitensis
rank: SPECIES
simple: Grallaria quitensis Lesson, 1844

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-02-20T01:07:57.233Z
created: 2016-02-20T01:07:57.233Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Grallaria
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Grallaria
family: Formicariidae
order: Passeriformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:

0

keywords: Grallaria quitesis

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 143--147

issue:

volume: 19

year: 2008

0: Greeney, H F

1: Harms, I

title: Behavior of the Tawny Antpitta (*Grallaria quitensis*) in northern Ecuador.

source: Ornitología Neotropical

type: article

tags: Grallaria quitesis

abstract:
last_modified:
created:
group_id:
profile_id:
id:

1

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Aglaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigrirostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Cistothorus platensis*, *Coccyzua pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocynna autumnalis*, *Dendrocynna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta caerulea*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jadinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus superciliaris*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxypogon guerini*, *Oxyura jamaicensis*, *Pachyramphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

institution:
address:
publisher: Princeton University Press
accessed:
websites:
chapter:
series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus,Actitis macularia,Adelomyia melanogenys,Agelaiocercus kingii,Amblycercus holosericeus,Ampelion rubrocristatus,Anas andium,Anas clypeata,Anas discors,Andigena nigrirostris,Anisognathus igniventris,Ardea alba,Ardea herodias,Arremon assimilis,Asio flammeus,Asio stygius,Asthenes flammulata,Asthenes fuliginosa,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Butorides virescens,Cairina moschata,Calidris bairdii,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Cathartes aura,Catharus minimus,Catharus ustulatus,Chaetocercus mulsant,Chaetura pelagica,Chalcostigma heteropogon,Chordeiles minor,Chrysomus icterocephalus,Cinclodes albidiventris,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Cistothorus platensis,Coccyzus pumila,Coccyzus americanus,Coccyzus melacoryphus,Coeligena bonapartei,Colibri coruscans,Colibri thalassinus,Columba livia,Columbina talpacoti,Conirostrum rufum,Conirostrum sitticolor,Contopus cinereus,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Dendrocygna autumnalis,Dendrocygna viduata,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dolichonyx oryzivorus,Dubusia taeniata,Egretta caerulea,Egretta thula,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera erufa,Eriocnemis cupreiventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Gallinula melanops,Geothlypis philadelphia,Geranoaetus melanoleucus,Glaucidium jardinii,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Heliangelus amethysticollis,Heliangelus exortis,Hellmayrea gularis,Hemispingus supercilii,Hemispingus verticalis,Henicorhina leucophrys,Hirundo rustica,Icterus chrysater,Icterus icterus,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Lafresnaya lafresnaya,Leiostyris alpestris,Leptasthenura andicola,Lesbia nuna,Lesbia victoriae,Machetornis rixosa,Margarornis squamiger,Mecocerculus leucophrys,Megascops choliba,Melanerpes rubricapillus,Merganetta armata,Metallura tyrianthina,Milvago chimachima,Mimus gilvus,Mniotilta varia,Molothrus bonariensis,Muscisaxicola alpinus,Myiarchus cinerascens,Myiodynastes luteiventris,Myiotheretes fumigatus,Myiotheretes striaticollis,Myiothlypis nigrocristata,Nomonyx dominicus,Nycticorax nycticorax,Ochthoeca cinnamomeiventris,Ochthoeca diadema,Ochthoeca fumicolor,Ochthoeca rufipetoralis,Orochelidon murina,Oxypogon guerini,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus,Parkesia noveboracensis,Patagioenas fasciata,Petrochelidon pyrrhonota,Phaetusa simplex,Phalacrocorax brasilianus,Pheucticus aureoventris,Pheucticus ludovicianus,Phimosus infuscatus,Phrygilus unicolor,Phyllomyias uropygialis,Playa cayana,Picoides fumigatus,Pipraeidea melanonota,Pipreola riefferii,Piranga olivacea,Piranga rubra,Pitangus sulphuratus,Pluvialis dominica,Podilymbus podiceps,Porphyrio martinicus,Porzana carolina,Progne tapera,Protonotaria citrea,Pseudocolopteryx acutipennis,Pseudoscops clamator,Pterophanes cyanopterus,Pygochelidon cyanoleuca,Pyrocephalus rubinus,Pyrrhomyias cinnamomeus,Quiscalus lugubris,Rallus semiplumbeus,Ramphocelus dimidiatus,Ramphomicrodon microhynchus,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea,Setophaga castanea,Setophaga cerulea,Setophaga petechia,Setophaga ruticilla,Setophaga striata,Sicalis flaveola,Sicalis luteola,Spiza americana,Sporagra spinescens,Sturnella magna,Synallaxis subpudica,Systellura longirostris,Tangara vitriolina,Thraupis episcopus,Thraupis palmarum,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon,Troglodytes solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto alba,Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaidura macroura,Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

reproductionUnstructured: Su biología general y su reproducción no son bien conocidas, solo se han reportado pocas presencias de nidos y en contadas ocasiones se han descrito sus huevos (Greeney & Harms, 2008). Para Colombia se han registrado dos individuos reproductivos en el departamento del Cauca en febrero; uno en el departamento de Santander en agosto y adicionalmente se han observado en junio juveniles en el departamento de Cundinamarca (Hilty & Brown, 1986).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

0

link: <http://www.hbw.com/node/56889>

keywords: Grallaria quitesis

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:
year: 2013
0: Krabbe, N K
1: Schulenberg, T S
title: Tawny Antpitta (*Grallaria quitensis*)
source: Handbook of the Birds of the World Alive
type: book_section
tags: *Grallaria quitensis*
abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

feedingUnstructured: En análisis de contenidos estomacales se ha encontrado varios insectos como escarabajos, larvas de mariposas y zancudos; sanguijuelas, semillas de bayas y otros frutos e incluso huesos de rana. La selección del alimento la hacen en la tierra y son muy hábiles para cazar debido a sus rápidos saltos y se conoce que las partes que no pueden digerir, son regurgitadas como una pequeña bolita (Krabbe & Schulenberg, 2013).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus*

institution:
address:
publisher: Christopher Helm

accessed:

websites:

chapter:

series:

pages: 880

issue:

year: 2006

0: Restall, R

1: Rodner, C

2: Lentino, M

title: Birds of Northern South America: An Identification Guide, Volume 1: Species Accounts.

source:

type: book

tags: Accipiter striatus, Aglaiocercus kingii, Amblycercus holosericeus, Anas andium, Andigena nigrirostris, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Catamenia inornata, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geranoaetus melanoleucus, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Se desplaza corriendo rápidamente o dando saltos en el suelo en busca de matorrales o hierba, es muy huidiza, pero a diferencia de otras Grallaria usualmente es muy fácil de ver, incluso se encarama en postes de la cerca, copas de arbustos u otros sitios elevados para vocalizar. Son individuos generalmente solitarios pero en ocasiones se les observa en parejas y son muy activas vocalmente (Hilty & Brown, 1986; Restall et al., 2006).

Frecuentemente se escuchan sus vocalizaciones; su canto es un silbido fuerte y claro que dice took,tu tu, en donde la primera nota es ligeramente más alta y más fuerte. Y su llamada es un solo silbido cheew (Hilty & Brown, 1986).

ancillaryData

dataObject:

additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:
additionalInformation:

reference

link: <http://maps.iucnredlist.org/map.html?id=22703333>
keywords: Grallaria quitesis

editors

institution:
address:
publisher:
accessed:
websites:
chapter:
series:
issue:
year: 2014

authors

authors: BirdLife International
authors: NatureServe

title: Bird Species Distribution Maps of the World. 2012. Grallaria quitensis. The IUCN Red List of Threatened Species. Version 2014.2.

source:

type: miscellany

tags: Grallaria quitesis

abstract:

identifiers

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:

dataType:

identifier:

distributionUnstructured: Su distribución inicia en Colombia y desciende a lo largo de los Andes atravesando el Ecuador hasta el norte de Perú. En Colombia se distribuye en la cordillera Oriental desde Santander hasta Cundinamarca y en la cordillera Central desde Caldas hasta Nariño (BirdLife International and NatureServe, 2014; Hilty & Brown, 1986).

El Tororoi Leonado habita las tierras altas de la cordillera Oriental y Central de Colombia, esta reportado entre los 2200 y los 3700 msnm, con un promedio de rango en los 3000 msnm (Hilty & Brown, 1986).

distributionAtomized

stateProvince: Tolima

ancillaryData

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

0

link: <http://www.birdlife.org/datazone/speciesfactsheet.php?id=5098>

keywords: Grallaria quitesis

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: BirdLife International

title: Species factsheet: Grallaria quitesis

source:
type: miscellany
tags: Grallaria quitesis
abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
populationBiologyUnstructured: El tamaño de la población mundial no se ha cuantificado, pero esta especie se ha descrito como "común" (BirdLife International, 2014).

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:

measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

abstract: La especie es común en el páramo. Se encuentra en el bosque enano y en arbustos dispersos bajo los árboles, en especial cerca de lagos o zonas pantanosas. Suele posarse en las partes altas de matorrales y setos de pasto. Se avistan con frecuencia ya que se aventuran en áreas abiertas, incluso en plena luz del día (Krabbe & Schulenberg, 2013).

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:
mediaURL: <http://www.xeno-canto.org/96374>
description:

subject

source: <http://www.xeno-canto.org>

audience

bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:

agent

contentType:
dataType:
identifier:

_id

_bsontype: ObjectId

id: VÇÄKð lg#çAp