Reporte de Fichas para el usuario: edith gómez Ficha ID: 56e09de683c45700544e3cd8 Fecha de creación: 2016-03-09T22:04:27.979Z annualCycles annualCycleUnstructured: habitats 0 dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: habitatUnstructured: Esta especie habita bosques tropicales de Sudámerica. commonNamesAtomized creation\_date: 2016-03-09T22:04:27.979Z usesManagementAndConservation ancillaryData managementAndConservationUnstructured: managementAndConservationAtomized identificationKeys fullDescription dataObject: additionalInformation: geoPoint: location: thumbnailURL: description:

source:

rightsHolder:

bibliographicCitation:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: Margen frontal del pronoto encima de la fóvea, poco lamelado; clípeo fuertemente punteado; menor contraste entre las partes oscuras y claras, cada una hacia el marrón o amarillo respectivamente. Placa dorsal del mesepisterno más amplia de manera que la sutura sale de la parte anterior formando un ángulo de cerca de 90°; escutelo usualmente sin una línea central notoria; mesoescuto con margen lateral interrumpido; su coloración que puede ir desde amarillo quemado hasta negro. briefDescription:

#### references

keywords: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus.Chartergellus punctatior.Chartergus chartarius.Chartergus globiventris.Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi,Parachartergus weyrauchi,Polybia bistriata,Polybia catillifex,Polybia eberhardae,Polybia emaciata,Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea

### editors

institution: address: publisher: accessed: websites: chapter: series: pages: 585 issue: year: 1978

#### authors

authors: Richards, O. W.

title: The social wasps of the Americas excluding the Vespinae

source: type: book

tags: Agelaia angulata angulata,Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Asteloeca traili,Brachygastra albula,Brachygastra

augusti,Brachygastra lecheguana,Brachygastra propodealis,Brachygastra scutellaris,Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae.Mischocyttarus insolitus.Mischocyttarus labiatus.Mischocyttarus latissimus.Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi,Parachartergus weyrauchi,Polybia bistriata,Polybia catillifex,Polybia eberhardae,Polybia emaciata,Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea abstract:

### identifiers

last\_modified: created: group\_id: profile\_id: id:

taxonRecordName

### ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

replacementNameFor

microReference:

source:
datatype:
identifier:
source:
datatype:
identifier:
note:

sanctioned
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
laterHomonymOf
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:  conservedAgainst
basedOn
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
basionym
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: spellingCorrentionOf
typeNameEntity source: datatype: identifier:

ruleConsidered:

source: datatype: identifier: source: datatype: identifier: typificacion typeOfType:

**lectotypeMicroReferenceVoucher** 

# lectotypePublicationVoucher

# voucherReference

simple:

microReference:

year:

publishedIn

simple: source: datatype: identifier:

specialAuthorship

### authors

# year

simple:

canonicalAuthorship

# authors

# year

simple:

simple: (Spinosa, 1851)

canonicalName

infraspecificEpithet: specificEpithet: infragenericEpithet:

linkType: ref: uninomial:

simple: Angiopolybia paraensis

rank: SPECIES

simple: Angiopolybia paraensis (Spinosa, 1851)

attributes

nomenclaturalCode: isAnamorphic: id:
synonymsAtomized
ancillaryData
synonymStatus:
synonymName
anyTwo:
anyOne
providerLink: microReference:
publishedIn
source: datatype: identifier:
reletedName
source: datatype: identifier: note: ruleConsidered: microReference:
publishedIn
source: datatype: identifier:
reletedName
source: datatype: identifier: note: ruleConsidered: microReference:
publishedIn
source: datatype: identifier:
reletedName
source: datatype: identifier: note:

ruleConsidered:

microReference:
publishedIn
source: datatype: identifier:
reletedName
source: datatype: identifier: note: ruleConsidered: microReference:
publishedIn
source: datatype: identifier:
reletedName
source: datatype: identifier: note: ruleConsidered: microReference:
publishedIn
source: datatype: identifier:
reletedName
source: datatype: identifier: note: ruleConsidered:
typeNameEntity
,
source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:
source: datatype: identifier: source: datatype: identifier: source: datatype:

microReference: year: source: datatype: identifier:
combinationAuthorship
basionymAuthorship
simple:
authorship
simple: simple: Ducke, 1905
epithet
infraspecificEpithet: specificEpithet: infragenericEpithet:
genus
linkType: ref: uninomial: simple: Polybia ruficornis rank: SPECIES simple: Polybia ruficornis Ducke, 1905 nomenclaturalCode: isAnamorphic: id:
hierarchy
ancillaryData
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
source: The Catalogue of Life, 3rd January 2011
audience
bibliographicCitation: rigthsHolder: rights:

license: title:

modified: 2016-03-09T22:04:07.967Z created: 2016-03-09T22:04:07.967Z

# agent

homepage:

role: email: phone: address: position: organisation: lastName: firstName: mimeType: dataType: identifier:

parentTaxon: Angiopolybia

higherClassification: infraspecificEpithet: specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Angiopolybia family: Vespidae order: Hymenoptera classHierarchy: Insecta phylum: Arthropoda kingdom: Animalia recommended: classification:

invasiveness

invasivenessUnstructured:

reproduction

### 0

dataObject:

additionalInformation:

geoPoint: location: thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

reproductionUnstructured:
0
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
audience
rightsHolder: rights: license: modified: created:
agent
mimeType: dataType: identifier: relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy: TraitBank measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: Sexual measurementType: Tipo de Reproducción measurementID:
molecularData
molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions
interactionsUnstructured:

# feeding dataObject: additionalInformation: keywords: Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis institution: Universidad de Costa Rica address: publisher: accessed: websites: chapter: series: issue: year: 2010 0: Valverde, J.P. title: Clave taxonómica ilustrada para los géneros y las especies de avispas eusociales de Costa Rica (Hymenoptera: Vespidae, Polistinae) source: type: thesis tags: Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis abstract: last modified: created: group\_id: profile\_id: id: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Equipo Coordinador del SiB Colombia license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: feedingUnstructured: strategy: Depredadores type: Heterótrofos - Omnívoros

environmentalEnvelope

environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured: distance: structureDispersed: type: purpose:
behavior
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: behaviorUnstructured: Sus nidos presentan una cubierta ovalada y entre 2 y 4 panales sujetos independientemente al sustrato, la entrada está localizada en la parte inferior. Puede ubicarse en construcciones.
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo:

measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

# endemicAtomized

### distribution

# ancillaryData

dataObject:

additionalInformation:

#### reference

geoPoint: location: thumbnailURL:

### mediaURL

description:

# subject

source:

# audience

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

### agent

mimeType: dataType: identifier:

distributionUnstructured: Se distribuye en Sudamérica (Guyana, Guyana Francesa, Perú, Bolivia, Brasil, Venezuela y Surinam) en regiones con rangos altitudinales de hasta 600 msnm. En Colombia se ha registrado en las regiones de la Orinoquia y la Amazonia.

#### distributionAtomized

stateProvince: Amazonas

# ancillaryData

locality: municipality: country: Colombia

temporalCoverage
endDate: startDate:
distributionScope
type:
threatStatus
territory
territoryUnstructured: areaOfOccupancy: extentOfOccurrence:
populationBiology
populationBiologyUnstructured:
legislation
legislationUnstructured:
directThreats
directThreatsUnstructured:
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID: abstract: Son insectos tropicales de Sudámerica y habitan áreas que alcanzan hasta los 600 msnm. Presentan
hábitos carroñeros y eusociales (Valverde, 2010). Sus colonias se caracterizan por presentar unos pocos cientos de

individuos y las diferencias entre las castas son muy leves (Richards, 1978).

# associatedParty

role:

personnelldentifier: personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone: postalCode: country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

# ancillaryData

#### dataObject:

additionalInformation: Información de tipos: La serie tipo al parecer está depositada en Instituto e Museo di Zoología, Univ. di Torino, Turín, Italia (MIZT).

Créditos específicos: Al Instituto Ámazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

Registros biológicos: Colección Entomológica, Instituto de Ciencias Naturales, Universidad Nacional de Colombia. Bogotá, Colombia http://www.biovirtual.unal.edu.co/colecciones/datos/animalia/consultasMammalia.jsp?clase=4

#### reference

geoPoint: location:

thumbnailURL:

# mediaURL

description:

# subject

source:

#### audience

bibliographicCitation:

rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

### agent

mimeType: dataType:

identifier:

\_id

\_bsontype: ObjectID id: Và•æfÄW DãÍ€