

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d46ec03c16479905cba884

Fecha de creación: 2016-02-29T16:15:59.537Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Habita sabanas, pastizales, áreas agrícolas, zonas abiertas, ecosistemas de agua dulce y marinos (Elizondo 2000).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Español

name: Chotacabras zumbÃ³n

creation_date: 2016-02-29T16:15:59.537Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

fullDescriptionUnstructured: Presenta una banda blanca y amplia aproximadamente en la mitad del ala entre la muñeca y el extremo. La garganta y la barra alar son blancas en ambos sexos. La hembra no tiene banda caudal blanca; la garganta ocasionalmente es color ante (Hilty y Brown 2001).

briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

replacementNameFor

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:

identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (J. R. Forster, 1771)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:

ref:
uninomial:
simple: Chordeiles minor
rank: SPECIES
simple: Chordeiles minor (J. R. Forster, 1771)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

ancillaryData

synonymStatus:
synonymName

anyTwo:

anyOne

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:

source:
datatype:
identifier:

source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:
datatype:
identifier:

[combinationAuthorship](#)

basionymAuthorship

simple:

authorship

simple:

simple: J. R. Forster, 1771

epithet

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

genus

linkType:

ref:

uninomial:

simple: Caprimulgus minor

rank: SPECIES

simple: Caprimulgus minor J. R. Forster, 1771

nomenclaturalCode:

isAnamorphic:

id:

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:
rights:
license:
title:
modified: 2016-02-29T16:15:39.537Z
created: 2016-02-29T16:15:39.537Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Chordeiles
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Chordeiles
family: Caprimulgidae
order: Strigiformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:

dataType:
identifier:
reproductionUnstructured: En la época de reproducción y como despliegue de cortejo se ven individuos haciendo vuelos en picada acompañados por un fragor fuerte (Hilty y Brown 2001).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

feedingUnstructured: Insectívoro, vuelan alto y consiguen escarabajos, mariposas nocturnas y avispas (Elizondo 2000; Poulin et al. 1996).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

behaviorUnstructured: Especie de ave migratoria boreal (Salaman et al. 2009). Su vuelo es boyante con aletazos alternados con aletazos más rápidos y erráticos. Crepuscular, ocasionalmente se ve durante el día volando sobre áreas abiertas (Hilty y Brown 2001). Pínt nasal (Hilty y Brown 2001).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataObject:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

[endemicAtomized](#)

distribution

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:
distributionUnstructured: Cría desde Canadá hasta el centro de Panamá, inverna en Suramérica hacia el sur de Argentina (Hilty y Brown 2001), hasta 2600 msnm (Hilty y Brown 2001).
En todo el país (Salaman et al. 2009; Hilty y Brown 2001), en la región Andina, Amazonia, Caribe, Pacífica, Orinoquia (Hilty y Brown 2001; Salaman et al. 2009).

distributionAtomized

stateProvince: Córdoba

ancillaryData

locality:
municipality:
country: Colombia

temporalCoverage

endDate:
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES: Apéndice II

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

populationBiologyUnstructured: Según BirdLife (2012) y Poulin et al. (1996) sus poblaciones están disminuyendo. El uso a gran escala de insecticidas puede ser en parte responsable de la disminución generalizada de esta especie, ya que los insectos son su principal fuente de alimento (MNR 2009).

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Ave mediana de 24 cm con alas largas y puntiagudas, presenta una muesca en la cola y una banda blanca conspicua que atraviesa las primarias en la mitad de la distancia entre la muñeca y la punta del ala (Hilty y Brown 2001).

[associatedParty](#)

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

[ancillaryData](#)

dataObject:

additionalInformation: Registros biológicos: Especímenes depositados en la Universidad Nacional de Colombia, sede Bogotá Colección de Ornitología.
Recursos multimedia: Sonidos de esta especie están disponibles en Xeno Canto Foundation e imágenes en Animal Diversity Web.
Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaría General de la CAN y el Instituto Sinchi para el proyecto “Definición y conformación de la PIRAA, a partir del fortalecimiento del nodo subnacional de Colombia: Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIATAC”, como parte del desarrollo del programa BIOCAN.
A Thomas McNish por la fotografía de esta especie de ave.
Imagen: BIR-3015B Chordeiles minor.jpg

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder: Nancy Carolina Rojas Riaño
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id
_bsontype: ObjectID
id: VÔnÂ< G™\0^@