

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56ca35c6f0106c67230e7459

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus

multus,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dubusia taeniata,Egretta caerulea,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreiventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Gallinula melanops,Geothlypis philadelphia,Glaucidium jadinii,Grallaria ruficapilla,Hemispingus superciliosus,Hemispingus verticalis,Henicorhina leucophrys,Hirundo rustica,Icterus chrysater,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Leiothlypis peregrina,Lesbia nuna,Lesbia victoriae

abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

1

link: Neotropical Birds Online [http://neotropical.birds.cornell.edu/portal/species/overview?p{\\\_}p{\\\_}spp=270456](http://neotropical.birds.cornell.edu/portal/species/overview?p{\_}p{\_}spp=270456)

keywords: Lesbia nuna

0: Schulenberg, T S

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2011

0: Sedgwick, C W

title: Green-tailed Trainbearer (Lesbia nuna)

source:

type: miscellany

tags: Lesbia nuna

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

2

link: <http://www.hbw.com/species/green-tailed-trainbearer-lesbia-nuna>

keywords: Lesbia nuna

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:  
series:  
issue:  
year: 1999  
0: Züchner, T  
1: Boesman, P  
title: Green-tailed Trainbearer (*Lesbia nuna*)  
source: Handbook of the Birds of the World Alive  
type: book\_section  
tags: *Lesbia nuna*  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:

habitatUnstructured: Se le observa en bordes donde el bosque limita con potreros o con áreas urbanas. Es frecuente en cultivos de *Eucalyptus globulus* o *Acacia* spp. (Asociación Bogotana de Ornitología, 2000). También en bosque secundario, bosques de *Polylepis*, en bordes de bosques, áreas arbustivas, pastizales, páramos y áreas de parques con vegetación (Sedgwick, 2011; Züchner & Boesman, 1999).

## [commonNamesAtomized](#)

## [ancillaryData](#)

usedBy:  
usedIn  
distributionUnstructured:  
endDate:  
startDate:

## [ancillaryData](#)

type:  
language: Inglés  
name: Green-tailed Trainbearer

usesManagementAndConservation

## [ancillaryData](#)

dataObject:

additionalInformation:

## reference

link: <http://www.hbw.com/species/green-tailed-trainbearer-lesbia-nuna>

keywords: Lesbia nuna

## editors

editors: del Hoyo, J

editors: Elliott, A

editors: Sargatal, J

editors: Christie, D A

editors: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 1999

## authors

authors: Z"uchner, T

authors: Boesman, P

title: Green-tailed Trainbearer (Lesbia nuna)

source: Handbook of the Birds of the World Alive

type: book\_section

tags: Lesbia nuna

abstract:

## identifiers

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

mimeType:

dataType:

identifier:

managementAndConservationUnstructured: Como consecuencia de la implementación de reglamentos nacionales e internacionales, se ha controlado la importación para el comercio de aves silvestres durante las últimas décadas (Züchner & Boesman, 1999).

managementAndConservationAtomized

0

0

dataObject:

additionalInformation:

0

link: <http://www.hbw.com/species/green-tailed-trainbearer-lesbia-nuna>

keywords: Lesbia nuna

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 1999

0: Züchner, T

1: Boesman, P

title: Green-tailed Trainbearer (Lesbia nuna)

source: Handbook of the Birds of the World Alive

type: book\_section

tags: Lesbia nuna

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
useNotes: Antes usada en el comercio de aves silvestres (Züchner & Boesman, 1999).  
potential:  
properties:  
ratingPopularity:  
economics:  
useTypeAtomized:  
conservationExploitationData:  
seasonOfAvailabilityUse:  
meansOfApplicationAdministration:  
productionDetails:  
vernacularNameUseAnnotations:  
organisms:  
users:  
partUsed:  
useValue:  
sourceOfInformationText:

## references

identificationKeys

fullDescription

0

dataObject:  
additionalInformation:

0

link: <http://www.hbw.com/species/green-tailed-trainbearer-lesbia-nuna>  
keywords: Lesbia nuna

0: del Hoyo, J  
1: Elliott, A  
2: Sargatal, J  
3: Christie, D A  
4: de Juana, E

institution:  
address:  
publisher: Lynx Edicions  
accessed:

websites:

chapter:

series:

issue:

year: 1999

0: Züchner, T

1: Boesman, P

title: Green-tailed Trainbearer (Lesbia nuna)

source: Handbook of the Birds of the World Alive

type: book\_section

tags: *Lesbia nuna*

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dubusia taeniata*, *Egretta caerulea*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Glaucidium jardinii*, *Grallaria ruficapilla*, *Hemispingus superciliosus*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Leiothlypis peregrina*, *Lesbia nuna*, *Lesbia victoriae*

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus*

fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: El Cometa Coliverde mide 5.5 cm más la cola, que en machos es de 10 u 11 cm y en las hembras de 4 o 5 cm. Su pico es corto, relativamente fino y casi recto. El cuerpo es verde pastoso brillante por encima; el macho es verde brillante debajo, su gorguera verde iridiscente; la cola es brillante por debajo, angosta, profundamente bifurcada, generalmente está cerrada y es casi dos veces el largo de su cuerpo; tiene un par de plumas externas oscuras, timoneras negras con la mitad distal verde esmeralda brillante. La hembra tiene una cola mucho más corta; por debajo es blanca con manchas verde brillante. La especie más similar es Lesbia victoriae pero ésta es mucho más bronceada y su cola es más larga y negra por encima (Asociación Bogotana de Ornitología, 2000). Los inmaduros son similares a la hembra adulta, el macho inmaduro tiene grandes discos verdes en la gorguera. Por las diferencias en comportamiento y despliegue no se sugiere una relación cercana con Lesbia victoriae, podría ser más cercano a Ramphomichus (Züchner & Boesman, 1999).

briefDescription:

## references

link: <http://www.hbw.com/species/green-tailed-trainbearer-lesbia-nuna>

keywords: Lesbia nuna

## editors

editors: del Hoyo, J

editors: Elliott, A

editors: Sargatal, J

editors: Christie, D A

editors: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:



chapter:  
series:  
issue:  
year: 1999

authors

authors: Z\"uchner, T  
authors: Boesman, P  
title: Green-tailed Trainbearer (Lesbia nuna)  
source: Handbook of the Birds of the World Alive  
type: book\_section  
tags: Lesbia nuna  
abstract:

identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
  
laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

conservedAgainst

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:

datatype:  
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:

publishedIn

simple:  
source:  
datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: (Lesson, 1832)

canonicalName

infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:

uninomial:  
simple: Lesbia nuna  
rank: SPECIES  
simple: Lesbia nuna (Lesson, 1832)

attributes

nomenclaturalCode:  
isAnamorphic:  
id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:  
additionalInformation:

[reference](#)

geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:  
rightsHolder:  
rights:  
license:  
title:  
modified: 2016-02-21T22:04:27.120Z  
created: 2016-02-21T22:04:27.120Z

[agent](#)

homepage:  
role:  
email:  
phone:  
address:  
position:  
organisation:  
lastName:  
firstName:  
mimeType:  
dataType:  
identifier:  
parentTaxon: Lesbia  
higherClassification:  
infraspecificEpithet:  
specificEpithet:  
taxonRank: SPECIES  
subGenus:  
genus: Lesbia  
family: Trochilidae  
order: Apodiformes  
classHierarchy: Aves  
phylum: Chordata

kingdom: Animalia

recommended:

classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:

additionalInformation:

0

link: <http://www.hbw.com/species/green-tailed-trainbearer-lesbia-nuna>

keywords: Lesbia nuna

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 1999

0: Z"uchner, T

1: Boesman, P

title: Green-tailed Trainbearer (Lesbia nuna)

source: Handbook of the Birds of the World Alive

type: book\_section

tags: Lesbia nuna

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

reproductionUnstructured: El Cometa Coliverde se reproduce entre noviembre y abril; el nido es en forma de copa y está construido de musgo y raicillas, forrados con material vegetal suave y puesto a 4.2 m del suelo. El tamaño de la nidada es de dos huevos (Züchner & Boesman, 1999).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

0

dataObject:

additionalInformation:

0

link: <http://www.hbw.com/species/green-tailed-trainbearer-lesbia-nuna>

keywords: Lesbia nuna

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 1999

0: Züchner, T

1: Boesman, P

title: Green-tailed Trainbearer (Lesbia nuna)

source: Handbook of the Birds of the World Alive

type: book\_section

tags: Lesbia nuna

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

interactionsUnstructured: Luego de las temporadas de reproducción o floración, este colibrí suele tener algunos movimientos altitudinales. En algunas localidades se le observa competencia con *Lesbia victoriae* (Züchner & Boesman, 1999).

feeding

0

dataObject:

additionalInformation:

0

keywords: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dubusia taeniata*, *Egretta caerulea*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Glaucidium jardinii*, *Grallaria ruficapilla*, *Hemispingus superciliosus*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Leiothlypis peregrina*, *Lesbia nuna*, *Lesbia victoriae*

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:  
year: 2000  
0: Asociación Bogotana de Ornitología  
title: Aves de la Sabana de Bogotá: guía de campo  
source:  
type: book  
tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

1

link: <http://www.hbw.com/species/green-tailed-trainbearer-lesbia-nuna>

keywords: Lesbia nuna

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 1999

0: Züchner, T

1: Boesman, P

title: Green-tailed Trainbearer (Lesbia nuna)

source: Handbook of the Birds of the World Alive

type: book\_section

tags: Lesbia nuna

abstract:

last\_modified:

created:

group\_id:

profile\_id:



id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:

feedingUnstructured: Forrajean de baja a mediana altura (Züchner & Boesman, 1999), visita pequeñas flores tubulares y se agarra de las mismas para libar. Con frecuencia prefiere la flor lila del garbancillo o espino (*Duranta muticii*) y las flores pequeñas de *Fuchsia* *Fuchsia microphylla* y *F. Arborescens*; también liba de flores de árboles nativos como cedrillos (*Phyllanthus salviifolius*) y cedros (*Cedrela montana*) (Asociación Bogotana de Ornitología, 2000). Atrapa insectos en el aire mediante vuelos prolongados y visita flores de *Espeletia* para sacar de allí pequeños insectos (Züchner & Boesman, 1999).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigristrostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Cistothorus platensis*, *Coccyzua pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa*

humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta

thula, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxypogon guerinii*, *Oxyura jamaicensis*, *Pachyramphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dubusia taeniata*, *Egretta caerulea*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Glaucidium jardinii*, *Grallaria ruficapilla*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Leiothlypis peregrina*, *Lesbia nuna*, *Lesbia victoriae*

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M  
7: Valencia, I D  
8: Knapp, D  
institution:  
edition: Primera Ed  
address:  
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca  
accessed:  
websites:  
chapter:  
series:  
pages: 276  
issue:  
year: 2000

0: Asociación Bogotana de Ornitología  
title: Aves de la Sabana de Bogotá: guía de campo  
source:  
type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae

abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

2

link: <http://www.hbw.com/species/green-tailed-trainbearer-lesbia-nuna>

keywords: Lesbia nuna

0: del Hoyo, J  
1: Elliott, A  
2: Sargatal, J  
3: Christie, D A  
4: de Juana, E

institution:  
address:  
publisher: Lynx Edicions  
accessed:  
websites:  
chapter:  
series:  
issue:

year: 1999  
0: Züchner, T  
1: Boesman, P  
title: Green-tailed Trainbearer (Lesbia nuna)  
source: Handbook of the Birds of the World Alive  
type: book\_section  
tags: Lesbia nuna  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:

behaviorUnstructured: Es activo y agresivo; su vuelo es cernido como el de una abeja (Hilty & Brown, 1986). En sus despliegues de cortejo y territoriales realiza un vuelo o picadas en forma de “U”, emitiendo un sonido en el punto más bajo (Asociación Bogotana de Ornitología, 2000). Las hembras defienden pequeños territorios de alimentación (Züchner & Boesman, 1999).

El canto es como una “risa” o trino de silbidos claros y tono descendente titiititi tutu (Asociación Bogotana de Ornitología, 2000); también un repetido drrrt drrrt y variaciones de estos (Züchner & Boesman, 1999). Durante sus vuelos de despliegue en forma de “U”, cuando llega al punto más bajo produce un chasquido fuerte tak-tak-tak-tak, como una pequeña ametralladora, producido seguramente con su cola (Asociación Bogotana de Ornitología, 2000); también emite un corto y alto bzzzt (Hilty & Brown, 1986; Züchner & Boesman, 1999).

#### ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

#### measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

## endemicAtomized

## distribution

## ancillaryData

dataObject:  
additionalInformation:

## reference

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae

## editors

editors: Stiles, F G  
editors: Bohórquez, C I  
editors: Cadena, C D  
editors: De la Zerda, S  
editors: Hernández, M  
editors: Roselli, L  
editors: Kelsey, M  
editors: Valencia, I D  
editors: Knapp, D  
institution:  
edition: Primera Ed  
address:  
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca  
accessed:  
websites:  
chapter:  
series:  
pages: 276  
issue:

year: 2000

## authors

authors: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae

abstract:

## identifiers

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

## agent

contentType:

dataType:

identifier:

distributionUnstructured: Se distribuye por la cordillera de los Andes desde Venezuela hasta el norte de Bolivia (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986). En Colombia se encuentra en ambas vertientes

de la cordillera Oriental, desde Cundinamarca hasta Boyacá; al occidente en el Cauca hasta Nariño, también en el Huila (Hilty & Brown, 1986). Se reconocen seis subespecies, diferenciadas en su distribución. *L. n. gouldii* (Loddiges, 1832) está al oriente de los Andes, en Boyacá y Cundinamarca y al occidente en Cauca y Nariño, existe un registro antiguo para Mérida, Venezuela. *L. n. gracilis* (Gould, 1846) en Ecuador. *L. n. aureliae* Weller & Schuchmann, 2004 se encuentra al sur de Ecuador. *L. n. pallidiventris* (Simon, 1902) presente al norte de los Andes y Perú. *L. n. huallagae* Weller & Schuchmann, 2004 en los Andes Peruanos. *L. n. nuna* (Lesson, 1832) al sur de los Andes peruanos. *L. n. boliviana* Boucard, 1892 en la vertiente Oriental de los Andes y en el noroccidente de Bolivia (Züchner & Boesman, 1999). Se distribuye entre los 1700 y 3800 msnm, pero principalmente entre los 2200 y 2800 msnm (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986).

distributionAtomized

stateProvince: Cauca

ancillaryData

locality:  
municipality:  
country: Colombia

temporalCoverage

endDate:  
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:  
locality:  
municipality:  
county:  
stateProvince:  
country: Global  
0: IUCN  
measurementValue: Preocupación menor (LC)  
relatedTo:  
measurementRemarks:  
measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementType: IUCN  
measurementID:

territory

territoryUnstructured:



areaOfOccupancy:  
extentOfOccurrence:  
populationBiology

0

dataObject:  
additionalInformation:

0

link: <http://www.birdlife.org/datazone/species/factsheet/22687974>

keywords: Lesbia nuna

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: BirdLife International

title: Species factsheet: Lesbia nuna

source:

type: miscellany

tags: Lesbia nuna

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

populationBiologyUnstructured: El tamaño global de la población no ha sido cuantificado, sin embargo se considera una especie bastante común con una tendencia estable sin evidencia de declinaciones o cambios considerables (BirdLife International, 2014).

legislation

legislationUnstructured:

directThreats

0

dataObject:  
additionalInformation:

0

link: <http://www.hbw.com/species/green-tailed-trainbearer-lesbia-nuna>

keywords: *Lesbia nuna*

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 1999

0: Züchner, T

1: Boesman, P

title: Green-tailed Trainbearer (*Lesbia nuna*)

source: Handbook of the Birds of the World Alive

type: book\_section

tags: *Lesbia nuna*

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

link: <http://www.cites.org/esp/app/appendices.php>

keywords: *Asio flammeus*, *Asio stygius*, *Buteo platypterus*, *Buteo swainsoni*, *Chaetocercus mulsant*, *Chalcostigma heteropogon*, *Colibri coruscans*, *Elanus leucurus*, *Eupsittula pertinax*, *Falco peregrinus*, *Forpus conspicillatus*, *Glaucidium jardinii*, *Lafresnaya lafresnayi*, *Lesbia nuna*, *Lesbia victoriae*, *Megascops choliba*, *Metallura tyrianthina*, *Tyto alba*

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: CITES

title: Apéndices CITES

source:

type: miscellany

tags: *Asio flammeus*, *Asio stygius*, *Buteo platypterus*, *Buteo swainsoni*, *Chaetocercus mulsant*, *Chalcostigma heteropogon*, *Colibri coruscans*, *Elanus leucurus*, *Eupsittula pertinax*, *Falco peregrinus*, *Forpus*

conspicillatus,Glaucidium jardinii,Lafresnaya lafresnayi,Lesbia nuna,Lesbia victoriae,Megascops choliba,Metallura tyrianthina,Tyto alba

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

directThreatsUnstructured: Algunos de sus hábitats están en peligro por la deforestación de los bosques de Polylepis, la ganadería y la agricultura (Züchner & Boesman, 1999). Aunque no se considere una especie amenazada se debe controlar el tráfico y comercialización de la misma (CITES, 2014).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Las colas bifurcadas de este pequeño Cometa Coliverde son más del doble de la longitud de su cuerpo.

Machos y hembras son verde brillante y las hembras debajo son blancas con manchas verdes. Habitan en áreas con crecimiento secundario, en bordes de bosques, áreas arbustivas, pastizales, páramos y áreas de parques con vegetación. En las elevaciones mayores por lo general compite con *Lesbia victoriae*. Es activo y defensivo, vuelan alrededor de las flores a diferentes alturas, las hembras mantienen pequeños territorios florales. Durante los despliegues de cortejo, los individuos realizan sonidos como chasquidos con sus colas (Sedgwick, 2011).

## associatedParty

role:  
personnelIdentifier:  
personnelDirectory:  
homepage:  
email: enataliagomez@gmail.com  
phone:  
postalCode:  
country: Colombia  
state: Bogotá Distrito Capital  
city: Bogotá D.C.  
address:  
organisation:  
position:  
lastName: Gómez Melendro  
firstName: Edith Natalia

## ancillaryData

dataObject:  
additionalInformation:

## reference

geoPoint:  
location:  
thumbnailURL:  
mediaURL: <http://www.xeno-canto.org/79764>  
description:

## subject

source: <http://www.xeno-canto.org>

## audience

bibliographicCitation:  
rightsHolder:  
rights:  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

contentType:  
dataFormat:  
identifier:

\_id

\_bsontype: ObjectId

id: VÊ5Æð lg#çE•