

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d4b80b3c16479905cba8b6

Fecha de creación: 2016-02-29T21:28:40.926Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Cuerpo de agua. Residente relativamente común (aunque muy dispersa) en aguas dulces y salobres.

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Francés

name: Aigrette bleue

creation_date: 2016-02-29T21:28:40.926Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
fullDescriptionUnstructured: Tiene un tamaño en promedio de 51 - 64 cm. Pico gris azul, dos tercios distales negros. Patas verdosas. Azul pizarra oscura con cabeza y cuello pardo rojizo intenso. Inm: blanco puro, generalmente con ápices de rémiges negruzcos, o moteadas de negruzco en diferentes grados; pico como el adulto; patas verdoso opaco.
briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:

identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:

datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Linnaeus, 1758)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Egretta caerulea
rank: SPECIES
simple: Egretta caerulea (Linnaeus, 1758)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

ancillaryData

synonymStatus:
synonymName

anyTwo:

anyOne

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:

source:
datatype:
identifier:

source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:
datatype:
identifier:

[combinationAuthorship](#)

basionymAuthorship

simple:

authorship

simple:

simple: Linnaeus, 1758

epithet

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

genus

linkType:

ref:

uninomial:

simple: Ardea caerulea

rank: SPECIES

simple: Ardea caerulea Linnaeus, 1758

nomenclaturalCode:

isAnamorphic:

id:

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:
rights:
license:
title:
modified: 2016-02-29T21:28:20.924Z
created: 2016-02-29T21:28:20.924Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Egretta
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Egretta
family: Ardeidae
order: Pelecaniformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:

dataType:

identifier:

reproductionUnstructured: Pueden construir nidos en colonias, por lo general en manglares. Nido plataforma de palitos 1 - 3 m sobre el agua; 2 a 5 huevos azules.

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

feedingUnstructured: Principalmente pequeños vertebrados y peces.

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

behaviorUnstructured: Diurna y acuática. Generalmente sola o en grupos de 2 ó 3; a veces busca alimento activamente. No se conoce una vocalización.

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataObject:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Clara Maria Pérez Gallego
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:
distributionUnstructured: S EEUU hasta W Perú, N Chile, Uruguay y SE Brasil. Antillas Holandesas; Trinidad y Tobago. En Colombia hasta 2600 m (Sabana de Bogotá) generalmente mucho más bajo. En todo el país en las regiones Pacífica, Caribe, Amazónica, Orinoquia y Andina, hasta los 2600 msnm.

distributionAtomized

temporalCoverage

endDate:
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:
threatStatusAtomized

apendiceCITES:
locality:
municipality:
county:
stateProvince:

country: Global
0: IUCN
measurementValue: Preocupación menor (LC)
relatedTo:
measurementRemarks:
measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementType: IUCN
measurementID:

territory

territoryUnstructured:
areaOfOccupancy:
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:

measurementType:
measurementID:
abstract:

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Registros biológicos: En la Central Jaguas se registró en el área intervenida de Embalse. En la Central San Carlos se registró en el área intervenida de Embalse.

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder: Clara Maria Pérez Gallego
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id

_bsonType: ObjectID

id: VÔ,°

$\leq G^{TM\backslash 0\zeta}$