Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c22230f0106c67230e7322

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

keywords: Accipiter striatus. Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Anas andium, Andigena nigrirostris, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Geranoaetus melanoleucus, Grallaria guitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis institution:

address:

publisher: Christopher Helm

accessed: websites: chapter: series: pages: 880 issue: year: 2006 0: Restall, R 1: Rodner, C

title: Birds of Northern South America: An Identification Guide, Volume 1: Species Accounts.

source: type: book

2: Lentino, M

tags: Accipiter striatus, Aglaiocercus kingii, Amblycercus holosericeus, Anas andium, Andigena nigrirostris, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Geranoaetus melanoleucus, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis

abstract: last modified:

created: group id: profile_id:

id:

1

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus.Chaetocercus mulsant.Chaetura pelagica.Chalcostigma heteropogon.Chordeiles minor.Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina.Oxypogon querinii.Oxyura iamaicensis.Pachyramphus versicolor.Pandion haliaetus.Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens. Sturnella magna. Synallaxis subpudica. Systellura longirostris. Tangara vitriolina. Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis institution:

address:

accessed:

publisher: Princeton University Press

websites: chapter: series: pages: 836 issue: year: 1986 0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus

igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccycua pumila.Coccyzus americanus.Coccyzus melacoryphus.Coeligena bonapartei.Colibri coruscans.Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria guitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis. Turdus fuscater. Turdus ignobilis. Tyrannus melancholicus. Tyrannus savana. Tyrannus tyrannus. Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis abstract:

last_modified: created: group_id: profile_id:

id:

2

link: http://www.hbw.com/node/61997 keywords: Atlapetes schistaceus

0: del Hoyo, J 1: Elliott, A 2: Sargatal, J 3: Christie, D A

4: de Juana, E

institution: address:

publisher: Lynx Edicions

accessed: websites: chapter: series: issue: vear: 2011 0: Jaramillo. A

title: Slaty Brush-finch (Atlapetes schistaceus) source: Handbook of the Birds of the World Alive

type: book section

tags: Atlapetes schistaceus

abstract: last modified: created: group id: profile_id:

id:

geoPoint: location: thumbnailURL: description: source: bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

habitatUnstructured: La especie habita desde zonas subtropicales hasta el páramo (Restall et al., 2006). Es común en los bordes de bosque, matorrales secundarios achaparrados, bosque y matorrales desde el sotobosque hasta el límite de vegetación arbórea (Hilty & Brown, 1986). También en el bosque nublado, bosque montano secundario, páramo con frailejón (Espeletia) y en Venezuela también bosque húmedo de Polylepis; evita el bosque continuo (Jaramillo, 2011).

commonNamesAtomized

ancillaryData

usedBy:

usedIn

distributionUnstructured:

endDate: startDate:

ancillaryData

type:

language: Alemán

name: Graubrust-Buschammer

usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed: websites: chapter: series: pages: 276 issue: year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source: type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides

striata abstract: last_modified: created: group_id: profile_id:

id:

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus

holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus.Cinclodes albidiventris.Cinclus leucocephalus.Cinnvcerthia unirufa.Cistothorus platensis.Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis. Myiothlypis nigrocristata. Nomonyx dominicus. Nycticorax nycticorax. Ochthoeca cinnamomeiventris,Ochthoeca diadema,Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina, Oxypogon querinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota. Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus. Thraupis palmarum. Tringa flavipes. Tringa melanoleuca. Tringa solitaria. Troglodytes aedon. Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis institution:

address:

accessed:

publisher: Princeton University Press

websites: chapter: series: pages: 836 issue: year: 1986 0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes

flammulata, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus.Columba livia.Columbina talpacoti.Conirostrum rufum.Conirostrum sitticolor.Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis. Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba. Vanellus chilensis. Vireo olivaceus. Vultur gryphus. Zenaida auriculata. Zonotrichia capensis abstract:

 $last_modified:$

created: group_id:

profile_id:

id:

geoPoint: location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType:

identifier:

fullDescriptionUnstructured: La especie mide 18 cm. Color general gris pizarroso, más oscuro por encima, con coronilla rufa y un patrón llamativo de blanco y negruzco en la cara. Lados de la cabeza negruzcos, garganta blanca con bigoteras negras; pequeña mancha alar blanca. Juveniles: patrón de la cabeza borroso y con listado fusco indistinto por debajo (más negro encima, coronilla castaño rufo, sin espéculo alar blanco, debajo gris pizarra oscuro, más pálido en garganta en subespecies fumidus y tamae) (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986). briefDescription:

references

link: http://www.hbw.com/node/61997 keywords: Atlapetes schistaceus

editors

editors: del Hoyo, J editors: Elliott, A editors: Sargatal, J editors: Christie, D A editors: de Juana, E

institution: address:

publisher: Lynx Edicions

accessed: websites: chapter: series: issue: year: 2011

authors

authors: Jaramillo, A

title: Slaty Brush-finch (Atlapetes schistaceus) source: Handbook of the Birds of the World Alive

type: book_section

tags: Atlapetes schistaceus

abstract:

identifiers

last_modified: created: group_id: profile_id: id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:

replacementNameFor

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

sanctioned

microReference:

source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:

laterHomonymOf

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

conservedAgainst

basedOn

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

basionym microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: spellingCorrentionOf typeNameEntity source: datatype: identifier: source: datatype: identifier: source: datatype: identifier: typificacion typeOfType: **lectotypeMicroReferenceVoucher lectotypePublicationVoucher** voucherReference simple: microReference: year: publishedIn simple: source: datatype: identifier: specialAuthorship authors year simple: canonicalAuthorship authors

year

simple:

simple: (Boissonneau, 1840)

canonicalName

infraspecificEpithet: specificEpithet: infragenericEpithet:

linkType: ref: uninomial:

simple: Atlapetes schistaceus

rank: SPECIES

simple: Atlapetes schistaceus (Boissonneau, 1840)

attributes

nomenclaturalCode: isAnamorphic:

id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rigthsHolder: rights:

license:

modified: 2016-02-15T18:36:44.794Z created: 2016-02-15T18:36:44.794Z

agent

homepage:

role:

email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:

parentTaxon: Atlapetes higherClassification: infraspecificEpithet: specificEpithet: taxonRank: SPECIES

subGenus:

genus: Atlapetes family: Emberizidae order: Passeriformes classHierarchy: Aves phylum: Chordata kingdom: Animalia recommended: classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

0

dataObject: additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder: rights: license: modified:

agent

created:

mimeType: dataType: identifier: relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy: TraitBank measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: Sexual measurementType: Tipo de Reproducción measurementID: molecularData molecularDataUnstructured: migratory migratoryUnstructured: lifeForm lifeFormUnstructured: lifeCycle lifeCycleUnstructured: interactions interactionsUnstructured: feeding dataObject: additionalInformation: link: http://www.hbw.com/node/61997 keywords: Atlapetes schistaceus 0: del Hoyo, J 1: Elliott, A 2: Sargatal, J 3: Christie, D A 4: de Juana, E institution: address: publisher: Lynx Edicions accessed: websites: chapter: series: issue: year: 2011 0: Jaramillo, A title: Slaty Brush-finch (Atlapetes schistaceus)

source: Handbook of the Birds of the World Alive type: book_section tags: Atlapetes schistaceus abstract: last_modified: created: group_id: profile_id: id:
1
keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata 0: Stiles, F G 1: Bohórquez, C I 2: Cadena, C D 3: De la Zerda, S 4: Hernández, M 5: Roselli, L 6: Kelsey, M 7: Valencia, I D 8: Knapp, D institution: edition: Primera Ed
address:
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca accessed: websites: chapter:
series: pages: 276
issue:
year: 2000
0: Asociación Bogotana de Ornitología title: Aves de la Sabana de Bogotá: guía de campo
source:
type: book
tags: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata abstract:
last_modified:
created:
group_id:
profile_id:
id: geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder: rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
mimeType: dataType:
identifier:
feedingUnstructured: El Atlapetes Pizarra se alimenta de insectos y frutos, forrajea desde el suelo hasta 5 m de altura, aunque prefiere niveles altos de matorrales y subdosel. Puede ser inquieto cuando forrajea, a menudo se sube a los bejucos en busca de alimento (Asociación Bogotana de Ornitología, 2000; Jaramillo, 2011).
0
type: Heterótrofos - Omnívoros
environmentalEnvelope
environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured: distance: structureDispersed: type:
purpose:
behavior
0
dataObject: additionalInformation:
0
keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata 0: Stiles, F G 1: Bohórquez, C I 2: Cadena, C D 3: De la Zerda, S 4: Hernández, M 5: Roselli, L 6: Kelsey, M 7: Valencia, I D 8: Knapp, D
institution:
edition: Primera Ed address:
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca accessed: websites:
chapter:
series:
pages: 276

issue: year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source: type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides

striata abstract: last_modified: created: group_id: profile_id: id:

1

keywords: Accipiter striatus, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Anas andium, Andigena nigrirostris, Anisognathus igniventris, Asthenes flammulata, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Calidris bairdii, Chalcostigma heteropogon, Chordeiles minor, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coeligena bonapartei, Colibri thalassinus, Coragyps atratus, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa sittoides, Eriocnemis vestita, Falco peregrinus, Gallinago nobilis, Gallinula galeata, Geranoaetus melanoleucus, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Mecocerculus leucophrys, Mniotilta varia, Molothrus bonariensis, Picoides fumigatus, Piranga olivacea, Pyrocephalus rubinus, Rupornis magnirostris, Sicalis flaveola, Sporagra spinescens, Sturnella magna, Tringa solitaria, Zonotrichia capensis

institution: address:

publisher: Christopher Helm

accessed:
websites:
chapter:
series:
pages: 880
issue:
year: 2006
0: Restall, R
1: Rodner, C

title: Birds of Northern South America: An Identification Guide, Volume 1: Species Accounts.

source: type: book

2: Lentino, M

tags: Accipiter striatus,Aglaiocercus kingii,Amblycercus holosericeus,Anas andium,Andigena nigrirostris,Anisognathus igniventris,Asthenes flammulata,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Calidris bairdii,Chalcostigma heteropogon,Chordeiles minor,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus platensis,Coeligena bonapartei,Colibri thalassinus,Coragyps atratus,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa sittoides,Eriocnemis vestita,Falco peregrinus,Gallinago nobilis,Gallinula galeata,Geranoaetus melanoleucus,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Mecocerculus leucophrys,Mniotilta varia,Molothrus bonariensis,Picoides fumigatus,Piranga olivacea,Pyrocephalus rubinus,Rupornis magnirostris,Sicalis flaveola,Sporagra spinescens,Sturnella magna,Tringa solitaria,Zonotrichia capensis

abstract: last_modified: created: group_id: profile_id: id:

2

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis.Catamenia analis.Catamenia inornata,Cathartes aura,Catharus minimus,Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax, nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis institution:

address:

publisher: Princeton University Press

accessed: websites: chapter: series: pages: 836 issue: year: 1986

0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata. Astragalinus psaltria. Atlapetes pallidinucha. Atlapetes schistaceus. Bubulcus ibis. Buteo platypterus. Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria guitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis. Muscisaxicola alpinus. Myiarchus crinitus. Myiodynastes Iuteiventris. Myiotheretes fumigatus. Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina,Oxypogon querinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus.Ramphocelus dimidiatus.Ramphomicron microrhynchum.Riparia riparia.Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis abstract:

last modified:

created:

group_id:

profile id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

mimeType:

dataType:

identifier:

behaviorUnstructured: El canto está compuesto por una serie enérgica y completa de chillidos agudos y trinos usualmente terminada en Chui-chui-chui o t'chiu,t'chiu,t'chiu (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986). Además, cuando hace contacto con el observador utiliza un fino tzit.tzit. El reclamo de alarma es un fino krryt krryt. El canto está compuesto por frases cortas como por ejemplo weyé-chuw (Restall et al., 2006).

ancillaryData

dataObject:

additionalInformation:

geoPoint: location:

thumbnailURL:

description:

source: http://www.biologyreference.com/Ar-Bi/Behavior-Patterns.html

bibliographicCitation: Alcock, John. Animal Behavior: An Evolutionary Approach, 6th ed. Sunderland, MA: Sinauer

Associates, Inc., 1997.

rightsHolder:

rights:

license: modified:

created:

mimeType:

dataType: identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue: Vocalizaciones measurementType: Comunicación

measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:

additionalInformation:

reference

link: http://maps.iucnredlist.org/map.html?id=22731233

keywords: Atlapetes schistaceus

editors

institution: address: publisher: accessed: websites: chapter: series: issue: year: 2014

authors

authors: BirdLife International

authors: NatureServe

title: Bird Species Distribution Maps of the World. 2012. Atlapetes schistaceus. The IUCN Red List of Threatened

Species. Version 2014.2

source:

type: miscellany

tags: Atlapetes schistaceus

abstract:

identifiers

last_modified:

created:

group_id: profile_id:

id:

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

agent

mimeType: dataType: identifier:

distributionUnstructured: La especie se distribuye desde el noroccidente de Venezuela a través de la cordillera de los Andes hasta el sur de Perú. La especie en Colombia se encuentra en la Serranía de Perijá, extremo norte de la cordillera Oriental al sur de Norte de Santander, nororiente de Santander y vertiente oriental de esta cordillera al norte Boyacá; suroccidente de Boyacá y Cundinamarca. En la cordillera Central y Occidental hacia el sur hasta Nariño (BirdLife International and NatureServe, 2014; Hilty & Brown, 1986). El Atlapetes Pizarra habita entre los

2400 y 3600 msnm, pero es más numeroso por encima de los 3000 msnm (Hilty & Brown, 1986). A veces hasta 1800 msnm (Jaramillo, 2011).

distributionAtomized

stateProvince: Cauca

ancillaryData

locality: municipality: country: Colombia

temporalCoverage

endDate: startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality: municipality: county:

stateProvince: country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks: measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit: measurementAccuracy: measurementType: IUCN measurementID:

territory

territoryUnstructured: areaOfOccupancy: extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:
directThreats
directThreatsUnstructured:
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID: abstract: El Atlapetes Pizarra es un ave casi completamente gris, más oscuro por encima, con coronilla rufa y garganta blanca con bigoteras negras. Prefiere andar en parejas o grupos pequeños de hasta 5 individuos en el sotobosque medio y subdosel de los bosques altoandinos y en los niveles altos de matorrales. Acompaña regularmente las bandadas mixtas; son ruidosos y conspicuos y se alimentan de frutos e insectos (Asociación Bogotana de Ornitología, 2000).
associatedParty
role: personnelDirectory: homepage: email: enataliagomezm@gmail.com phone: postalCode: country: Colombia state: Bogotá Distrito Capital city: Bogotá D.C. address: organisation: position: lastName: Gómez Melendro

firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL: http://www.xeno-canto.org/128412

description:

subject

source: http://www.xeno-canto.org

audience

bibliographicCitation:

rightsHolder:

rights: license: modified: created:

agent

mimeType: dataType: identifier:

_id

_bsontype: ObjectID id: VÂ"0ð lg#ç2