

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c3dbbdf0106c67230e7365

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

isbn: 0963699636

keywords: *Anas clypeata*, *Anas platyrhynchos*, *Ardea herodias*, *Astragalinus psaltria*, *Bartramia longicauda*, *Buteo swainsoni*, *Cairina moschata*, *Calidris melanotos*, *Cardellina canadensis*, *Cathartes aura*, *Catharus minimus*, *Coccyzus americanus*, *Columbina talpacoti*, *Contopus cooperi*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Dendrocygna autumnalis*, *Egretta caerulea*, *Egretta thula*, *Empidonax alnorum*, *Empidonax traillii*, *Falco columbarius*, *Fulica americana*, *Geothlypis philadelphia*, *Hirundo rustica*, *Ixobrychus exilis*, *Jacana jacana*, *Mniotilta varia*, *Nycticorax nycticorax*, *Oxyura jamaicensis*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Phalacrocorax brasilianus*, *Piranga olivacea*, *Pitangus sulphuratus*, *Podilymbus podiceps*, *Porzana carolina*, *Pyrocephalus rubinus*, *Rupornis magnirostris*, *Setophaga fusca*, *Sturnella magna*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*

0: Alderfer, Jonathan

institution:

address:

publisher: Washington D.C. National Geographic Society

accessed:

websites:

chapter:

series:

pages: 664

issue:

year: 2006

0: Alderfer, J

1: Zickefoose, J

2: Littlehales, B

title: National Geographic Complete Birds of North America

source:

type: book

tags: *Anas clypeata*, *Anas platyrhynchos*, *Ardea herodias*, *Astragalinus psaltria*, *Bartramia longicauda*, *Buteo swainsoni*, *Cairina moschata*, *Calidris melanotos*, *Cardellina canadensis*, *Cathartes aura*, *Catharus minimus*, *Coccyzus americanus*, *Columbina talpacoti*, *Contopus cooperi*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Dendrocygna autumnalis*, *Egretta caerulea*, *Egretta thula*, *Empidonax alnorum*, *Empidonax traillii*, *Falco columbarius*, *Fulica americana*, *Geothlypis philadelphia*, *Hirundo rustica*, *Ixobrychus exilis*, *Jacana jacana*, *Mniotilta varia*, *Nycticorax nycticorax*, *Oxyura jamaicensis*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Phalacrocorax brasilianus*, *Piranga olivacea*, *Pitangus sulphuratus*, *Podilymbus podiceps*, *Porzana carolina*, *Pyrocephalus rubinus*, *Rupornis magnirostris*, *Setophaga fusca*, *Sturnella magna*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

habitatUnstructured: Utilizan zonas arborizadas con vegetación densa y cuerpos de agua cercanos como arroyos y pantanos. También se encuentran en áreas de bosques con matorrales y arbustos bajos. Presente a su vez en humedales, rastrojos y cercas vivas (Asociación Bogotana de Ornitología, 2000; Alderfer et al., 2006).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Francés

name: Coulicou à bec jaune

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

link: <http://eol.org/pages/915103/overview>

keywords: Coccyzus americanus

0: Parr, Cyndy

1: Hole, Robert

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: EOL

title: Coccyzus americanus

source: Encyclopedia of life

type: miscellany

tags: Coccozyus americanus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccozyus americanus

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccozyus americanus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

2

isbn: 0963699636

keywords: Anas clypeata, Anas platyrhynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccozyus americanus, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna

autumnalis, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mniotilta varia, Nycticorax nycticorax, Oxyura jamaicensis, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

0: Alderfer, Jonathan

institution:

address:

publisher: Washington D.C. National Geographic Society

accessed:

websites:

chapter:

series:

pages: 664

issue:

year: 2006

0: Alderfer, J

1: Zickefoose, J

2: Littlehales, B

title: National Geographic Complete Birds of North America

source:

type: book

tags: Anas clypeata, Anas platyrhynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccyzus americanus, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna autumnalis, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mniotilta varia, Nycticorax nycticorax, Oxyura jamaicensis, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: Su silueta general es delgada, cola larga y alas de punta redondeada, mide aproximadamente 30 cm de largo y 60 g de peso. Pecho, vientre y región anal blancos, mientras que el cuello, espalda, la coronilla y cabeza son principalmente grises. Las primarias de las alas son rojizas y las rectrices cafés machadas con franjas blancas. Anillo ocular amarillo, pico curvo y delgado con el culmen y la maxila en general

negros. La mandíbula es amarilla, con la punta negra. Los individuos inmaduros son bastante similares a los adultos pero con coberteras infracaudales anteadas. El patrón de la cola no es tan marcado y el anillo orbital es de un amarillo más opaco (Asociación Bogotana de Ornitología, 2000; Alderfer et al., 2006; EOL, 2014).

briefDescription:

## references

link: <http://eol.org/pages/915103/overview>

keywords: *Coccyzus americanus*

## editors

editors: Parr, Cyndy

editors: Hole, Robert

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

## authors

authors: EOL

title: *Coccyzus americanus*

source: Encyclopedia of life

type: miscellany

tags: *Coccyzus americanus*

abstract:

## identifiers

last\_modified:

created:

group\_id:

profile\_id:

id:

taxonRecordName

## ancillaryData

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:  
ruleConsidered:  
  
replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

conservedAgainst

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:

source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:

publishedIn

simple:  
source:  
datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: (Linnaeus, 1758)



canonicalName

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

linkType:

ref:

uninomial:

simple: Coccyzus americanus

rank: SPECIES

simple: Coccyzus americanus (Linnaeus, 1758)

attributes

nomenclaturalCode:

isAnamorphic:

id:

synonymsAtomized

ancillaryData

synonymStatus:

synonymName

anyTwo:

providerLink:

microReference:

publishedIn

source:

datatype:

identifier:

reletedName

source:

datatype:

identifier:

note:

ruleConsidered:

microReference:

publishedIn

source:

datatype:

identifier:

note:

ruleConsidered:

microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:

typeNameEntity

source:

datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typeVoucherEntity

typeOfType:  
simple:  
microReference:  
year:  
source:  
datatype:  
identifier:

combinationAuthorship

basionymAuthorship

simple:  
authorship  
simple:  
simple: (Linnaeus, 1758)

epithet  
infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:

genus  
linkType:  
ref:  
uninomial:  
simple: Coccyzus americanus americanus  
rank: SUBSPECIES  
simple: Coccyzus americanus americanus (Linnaeus, 1758)  
nomenclaturalCode:  
isAnamorphic:  
id:

hierarchy

ancillaryData

dataObject:  
additionalInformation:

reference

geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source: The Catalogue of Life, 3rd January 2011

## audience

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-02-17T02:26:06.533Z

created: 2016-02-17T02:26:06.533Z

## agent

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataType:

identifier:

parentTaxon: Coccyzus

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Coccyzus

family: Cuculidae

order: Cuculiformes

classHierarchy: Aves

phylum: Chordata

kingdom: Animalia

recommended:

classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:

additionalInformation:

0

link: <http://www.allaboutbirds.org/Page.aspx?pid=1189>  
keywords: Bubulcus ibis,Cardellina canadensis,Coccyzus americanus  
institution:  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
issue:  
year: 2014  
0: Cornell University  
title: The Cornell Lab of Ornithology  
source:  
type: miscellany  
tags: Bubulcus ibis,Cardellina canadensis,Coccyzus americanus  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

1

link: <http://eol.org/pages/915103/overview>  
keywords: Coccyzus americanus  
0: Parr, Cyndy  
1: Hole, Robert  
institution:  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
issue:  
year: 2014  
0: EOL  
title: Coccyzus americanus  
source: Encyclopedia of life  
type: miscellany  
tags: Coccyzus americanus  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:

rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

reproductionUnstructured: El macho corteja a la hembra apartándose de la rama de percha y agachando sus hombros hacia ella; durante la cópula ambos están agarrados a la rama de la percha. Posteriormente tanto el macho y la hembra construyen juntos un nido de forma plana y oblicua de aproximadamente 20 centímetros de diámetro usando ramas que encuentran en el suelo o sueltas en árboles. Localizan los nidos en árboles de altura media que les pueden servir también de lugar de forrajeo. El tamaño de la nidada oscila entre 1 a 5 huevos con un periodo de incubación de aproximadamente 10 días. La cría nace a los 8 días de la puesta del huevo, cubierta de plumón y con los ojos abiertos. Los huevos son lisos, verde azulados sin marca. Cuando la hembra está receptiva se percha en una rama con la cabeza arriba (Cornell University, 2014; EOL, 2014).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

0

keywords: *Actitis macularia*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:  
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

isbn: 0963699636

keywords: Anas clypeata, Anas platyrhynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccyzus americanus, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna autumnalis, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mniotilta varia, Nycticorax nycticorax, Oxyura jamaicensis, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

0: Alderfer, Jonathan

institution:

address:

publisher: Washington D.C. National Geographic Society

accessed:

websites:

chapter:

series:

pages: 664

issue:

year: 2006

0: Alderfer, J

1: Zickefoose, J

2: Littlehales, B

title: National Geographic Complete Birds of North America

source:

type: book

tags: Anas clypeata, Anas platyrhynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccyzus

americanus, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna autumnalis, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mniotilta varia, Nycticorax nycticorax, Oxyura jamaicensis, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

2

link: <http://www.allaboutbirds.org/Page.aspx?pid=1189>  
keywords: Bubulcus ibis, Cardellina canadensis, Coccyzus americanus  
institution:  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
issue:  
year: 2014  
0: Cornell University  
title: The Cornell Lab of Ornithology  
source:  
type: miscellany  
tags: Bubulcus ibis, Cardellina canadensis, Coccyzus americanus  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

3

link: [http://awsassets.panda.org/downloads/migratorias{\\\_}aves{\\\_}42{\\\_}final.pdf](http://awsassets.panda.org/downloads/migratorias{\_}aves{\_}42{\_}final.pdf)  
keywords: Coccyzus americanus  
0: Naranjo, L G  
1: Amaya, J D  
2: Eusse-González, D  
3: Cifuentes-Sarmiento, Y  
institution:  
address:  
publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia  
accessed:  
websites:  
chapter:  
series:  
pages: 708  
issue:  
volume: 1



year: 2012  
0: Morales-Rozo, Andrea  
title: Coccozyus americanus  
source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves  
type: book\_section  
tags: Coccozyus americanus

abstract:  
last\_modified:

created:  
group\_id:  
profile\_id:  
id:

geoPoint:  
location:  
thumbnailURL:

description:  
source:

bibliographicCitation:  
rightsHolder:

rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:  
created:

contentType:  
dataType:

identifier:

feedingUnstructured: En general se alimenta de insectos pero tiene preferencia por las orugas, aunque también come otros gusanos. Es una de las pocas especies de aves que puede comer orugas peludas venenosas. También come polillas, escarabajos, hormigas, grillos y arañas. Hay registros de verlos cazando pequeñas ranas o lagartijas y en invierno pueden comer bayas u otros tipos de frutos y semillas (Asociación Bogotana de Ornitología, 2000; Alderfer et al., 2006; Cornell University, 2014; Morales-Rozo, 2012).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:  
structureDispersed:

type:  
purpose:

behavior

0

dataObject:  
additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinuca, Atlapetes schistaceus, Bartramia

longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

link: <http://www.allaboutbirds.org/Page.aspx?pid=1189>

keywords: Bubulcus ibis,Cardellina canadensis,Coccyzus americanus

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: Cornell University

title: The Cornell Lab of Ornithology

source:

type: miscellany

tags: Bubulcus ibis,Cardellina canadensis,Coccyzus americanus

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Es común verlo en bosques densos, donde forrajea principalmente perchedo y observando en busca de alimento, especialmente orugas. Atrapa sus presas con un salto o vuelo sobre ellas. Suele tomar una postura en la que encorva sus hombros para tratar de ocultar su pecho blanco. En la Sabana de Bogotá es común observar aves cansadas del viaje de migración, por lo que es frecuente registrar choques contra ventanas y vidrios (Asociación Bogotana de Ornitología, 2000; Cornell University, 2014).

En Norteamérica los machos vocalizan una serie de sonidos huecos pausados en sílabas “ka-ka-ka-ka-ka-kow-kow-kowlp-kowlp-kowlp-kowlp”. Tanto machos como hembras emiten sonidos durante el cortejo, que son suaves a manera de arrullo. En época de migración no emite vocalizaciones, por lo que en su estadía en Colombia y el resto de sus zonas de invernada es generalmente silenciosa (Asociación Bogotana de Ornitología, 2000; Cornell University, 2014).

## ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

## measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

## endemicAtomized

## distribution

## ancillaryData

dataObject:  
additionalInformation:

## reference

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus

## editors

editors: Stiles, F G  
editors: Bohórquez, C I  
editors: Cadena, C D  
editors: De la Zerda, S  
editors: Hernández, M  
editors: Roselli, L  
editors: Kelsey, M  
editors: Valencia, I D  
editors: Knapp, D  
institution:  
edition: Primera Ed  
address:  
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca  
accessed:  
websites:  
chapter:  
series:  
pages: 276  
issue:  
year: 2000

## authors

authors: Asociación Bogotana de Ornitología  
title: Aves de la Sabana de Bogotá: guía de campo  
source:  
type: book  
tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus

abstract:

## identifiers

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

## agent

contentType:

dataType:

identifier:

distributionUnstructured: Esta especie de cuco se distribuye en el oriente de Norteamérica sin embargo existen algunos registros ocasionales en el oeste. Durante la migración se distribuyen en toda Centroamérica incluyendo algunas islas del Caribe y en Sudamérica hasta el oriente de Perú y el Norte de Argentina. En Colombia se distribuye por todo el territorio nacional en época de invernada (Asociación Bogotana de Ornitología, 2000; Alderfer et al, 2006; Hilty & Brown, 1986). Tiene una distribución altitudinal bastante amplia, se puede encontrar desde el nivel del mar hasta los 2600 msnm (Asociación Bogotana de Ornitología, 2000; Alderfer et al., 2006; Hilty & Brown, 1986; Morales-Rozo, 2012).

## distributionAtomized

stateProvince: Córdoba

## ancillaryData

locality:

municipality:

country: Colombia

## temporalCoverage

endDate:

startDate:

## distributionScope

type:

## threatStatus

## ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

## measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

0

link: <http://www.iucnredlist.org/details/22684331/0>

keywords: Coccozys americanus

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: BirdLife International

title: Species factsheet: Coccozys americanus

source:

type: miscellany

tags: Coccozys americanus

abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

1

isbn: 0963699636  
keywords: Anas clypeata, Anas platyrhynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccyzus americanus, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna autumnalis, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mniotilta varia, Nycticorax nycticorax, Oxyura jamaicensis, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

0: Alderfer, Jonathan  
institution:  
address:  
publisher: Washington D.C. National Geographic Society

accessed:  
websites:

chapter:  
series:  
pages: 664  
issue:

year: 2006  
0: Alderfer, J  
1: Zickefoose, J  
2: Littlehales, B

title: National Geographic Complete Birds of North America

source:  
type: book

tags: Anas clypeata, Anas platyrhynchos, Ardea herodias, Astragalinus psaltria, Bartramia longicauda, Buteo swainsoni, Cairina moschata, Calidris melanotos, Cardellina canadensis, Cathartes aura, Catharus minimus, Coccyzus americanus, Columbina talpacoti, Contopus cooperi, Contopus virens, Coragyps atratus, Crotophaga ani, Dendrocygna autumnalis, Egretta caerulea, Egretta thula, Empidonax alnorum, Empidonax traillii, Falco columbarius, Fulica americana, Geothlypis philadelphia, Hirundo rustica, Ixobrychus exilis, Jacana jacana, Mniotilta varia, Nycticorax nycticorax, Oxyura jamaicensis, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Phalacrocorax brasilianus, Piranga olivacea, Pitangus sulphuratus, Podilymbus podiceps, Porzana carolina, Pyrocephalus rubinus, Rupornis magnirostris, Setophaga fusca, Sturnella magna, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon

abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:

id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

populationBiologyUnstructured: Las poblaciones en el oeste de Estados Unidos están disminuyendo rápidamente y en general la tendencia de la población es a disminuir (Alderfer et al., 2006; BirdLife International, 2014).

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Migratorio boreal común en buena parte de Colombia, especialmente en la zona andina. Es café grisáceo por encima y blanco por debajo, normalmente tímido pero fácil de ver, tiene una figura delgada y cola larga, el pico curvo, con la maxila negra y la mandíbula amarilla muy característico (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986; Morales-Rozo, 2012). En la cara, un anillo ocular amarillo muy útil para su identificación. Es una especie que se reproduce en Norteamérica y empieza su migración a mediados de julio, para llegar a Sudamérica en los meses de septiembre y octubre. Consume principalmente insectos con movimientos poco vistosos y aparentemente con baja actividad (Alderfer et al., 2006; Morales-Rozo, 2012).

[associatedParty](#)



role:  
personnelIdentifier:  
personnelDirectory:  
homepage:  
email: enataliagomez@gmail.com  
phone:  
postalCode:  
country: Colombia  
state: Bogotá Distrito Capital  
city: Bogotá D.C.  
address:  
organisation:  
position:  
lastName: Gómez Melendro  
firstName: Edith Natalia

## ancillaryData

dataObject:  
additionalInformation:

## reference

geoPoint:  
location:  
thumbnailURL:  
mediaURL: <https://www.flickr.com/photos/puttefin/6199547802/in/photolist-8Kxxdz-hevpiU-e4T7Eg-vkPhsE-arQizd-vCoyiR-bWumtX-vCLsXT-vA61iA-eRTGYq-2jrDko-dCzFog-abcRJu-ae7d7e-cck2p1-a7hk6u-bnPJRL-e3j1qc-ukfz7p-axej71-dHcMAi-sdhkBb-sdqtNr-ry4R34-sdhkCo-czydjC-vxduAr-vfh6es-xP5sQ3-xP5tam-DviJVp-sdhkDf-auFQA2-suSVin-caUEBU-x2BJzE-rWcHej-oF3ZnB-crmRi3-p7uj7A-d2Rb6N-8FRdt7-8JgjjP-8JjnLw-8Jgj7H-3rAN4g-asUYEL-b2PZhgf-f4cz5i-f4cAmB>  
description:

## subject

source: [https://farm7.staticflickr.com/6176/6199547802\\_0061a12fef.jpg](https://farm7.staticflickr.com/6176/6199547802_0061a12fef.jpg)

## audience

bibliographicCitation: <a href="http://www.allaboutbirds.org/guide/Yellow-billed\_Cuckoo/lifehistory" rel="nofollow">www.allaboutbirds.org/guide/Yellow-billed\_Cuckoo/lifehistory</a>:  
The entire period from egg laying to fledgling leaving the nest lasts only 17 days. On day six or seven after hatching, the feathers of the young burst out of their sheaths, allowing the nestling to become fully feathered in two hours.

Birds! Awesome. Who'd want to shoot anything else.

<b>Member of the <a href="http://www.flickr.com/groups/1390236@N25/"> <b>Flickr Bird Brigade</b> </a> </b>  
<i>Activists for birds and wildlife</i> </i>  
rightsHolder:  
rights:  
license: <http://creativecommons.org/licenses/by-nd/2.0/>  
modified:  
created:

## agent

mimeType:

dataType:  
identifier:

\_id

\_bsontype: ObjectID  
id: VÄU½ð lg#ç6P