

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c372e8f0106c67230e734b

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

keywords: Accipiter striatus,Actitis macularius,Adelomyia melanogenys,Aglaiocercus kingii,Amblycercus holosericeus,Ampelion rubrocristatus,Anas andium,Anas clypeata,Anas discors,Andigena nigrirostris,Anisognathus igniventris,Ardea alba,Ardea herodias,Arremon assimilis,Asio flammeus,Asio stygius,Asthenes flammulata,Asthenes fuliginosa,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Butorides virescens,Cairina moschata,Calidris bairdii,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Cathartes aura,Catharus minimus,Catharus ustulatus,Chaetocercus mulsant,Chaetura pelagica,Chalcostigma heteropogon,Chordeiles minor,Chrysomus icterocephalus,Cinclodes albidiventris,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus platensis,Coccyzus pumila,Coccyzus americanus,Coccyzus melacoryphus,Coeligena bonapartei,Colibri coruscans,Colibri thalassinus,Columba livia,Columbina talpacoti,Conirostrum rufum,Conirostrum sitticolor,Contopus cinereus,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Dendrocygna autumnalis,Dendrocygna viduata,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dolichonyx oryzivorus,Dubusia taeniata,Egretta thula,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreovertris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Geothlypis philadelphia,Geranoaetus melanoleucus,Glaucidium jardinii,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Heliangelus amethysticollis,Heliangelus exortis,Hellmayrea gularis,Hemispingus superciliosus,Hemispingus verticalis,Henicorhina leucophrys,Hirundo rustica,Icterus chrysater,Icterus icterus,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Lafresnaya lafresnayi,Leiothlypis peregrina,Leptasthenura andicola,Lesbia nuna,Lesbia victoriae,Machetornis rixosa,Margarornis squamiger,Mecocerculus leucophrys,Megascops choliba,Melanerpes rubricapillus,Merganetta armata,Metallura tyrianthina,Milvago chimachima,Mimus gilvus,Mniotilta varia,Molothrus bonariensis,Muscisaxicola alpinus,Myiarchus crinitus,Myiodynastes luteiventris,Myiotheretes fumigatus,Myiotheretes striaticollis,Myiothlypis nigrocristata,Nomonyx dominicus,Nycticorax nycticorax,Ochthoeca cinnamomeiventris,Ochthoeca diadema,Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina,Oxyopogon guerini,Oxyura jamaicensis,Pachyrhamphus versicolor,Pandion haliaetus,Pardirallus maculatus,Parkesia noveboracensis,Patagioenas fasciata,Petrochelidon pyrrhonota,Phaetusa simplex,Phalacrocorax brasilianus,Pheucticus aureoventris,Pheucticus ludovicianus,Phimosus infuscatus,Phrygilus unicolor,Phyllomyias uropygialis,Playa cayana,Picoides fumigatus,Pipraeidea melanonota,Pipreola riefferii,Piranga olivacea,Piranga rubra,Pitangus sulphuratus,Pluvialis dominica,Podilymbus podiceps,Porphyrion martinicus,Porzana carolina,Progne tapera,Protonotaria citrea,Pseudocolopteryx acutipennis,Pseudoscops clamator,Pterophanes cyanopterus,Pygochelidon cyanoleuca,Pyrocephalus rubinus,Pyrrhomyias cinnamomeus,Quiscalus lugubris,Rallus semiplumbeus,Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea,Setophaga castanea,Setophaga cerulea,Setophaga petechia,Setophaga ruticilla,Setophaga striata,Sicalis flaveola,Sicalis luteola,Spiza americana,Sporagra spinescens,Sturnella magna,Synallaxis subpudica,Systellura longirostris,Tangara vitriolina,Thraupis episcopus,Thraupis palmarum,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon,Troglodytes solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto alba,Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaidura macroura,Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigristrostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccyzua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiostyris alpestris, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: Asio flammeus,Asio stygius,Chaetocercus mulsant,Megascops choliba

institution:

address:

publisher: Centro de Ornitología y Biodiversidad CORBIDI

accessed:

websites:

chapter:

series:

pages: 645

issue:

year: 2010

0: Schulenberg, T S

1: Stotz, Douglas F

2: Lane, Daniel F

3: O'Neill, John P

4: Parker III, A Theodore

title: Aves de Perú

source:

type: book

tags: Asio flammeus,Asio stygius,Chaetocercus mulsant,Megascops choliba

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

habitatUnstructured: Es poco común y local en los bordes de selva húmeda y pastizales y áreas cultivadas con árboles dispersos y setos. Principalmente en terrenos abiertos, no están en selva (Hilty & Brown, 1986). Vive principalmente en las flores de los árboles, en zonas semiabiertas y jardines; rara vez se encuentra en los bordes del bosque húmedo montano (Hilty & Brown, 1986; Schulenberg et al., 2010).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:  
endDate:  
startDate:

ancillaryData

type:  
language: Inglés  
name: White-bellied Woodstar

usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:  
additionalInformation:

0

keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant

0: Stiles, F G  
1: Bohórquez, C I  
2: Cadena, C D  
3: De la Zerda, S  
4: Hernández, M  
5: Roselli, L  
6: Kelsey, M  
7: Valencia, I D  
8: Knapp, D  
institution:  
edition: Primera Ed  
address:  
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca  
accessed:  
websites:  
chapter:  
series:  
pages: 276  
issue:  
year: 2000  
0: Asociación Bogotana de Ornitología  
title: Aves de la Sabana de Bogotá: guía de campo  
source:  
type: book

tags: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Aglaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigristrostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus platensis*, *Coccyzus pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreovertris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus superciliaris*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxyphaps squamiger*, *Oxyura jamaicensis*, *Pachyrhamphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccyzua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

2

link: <http://www.museum.lsu.edu/~Remsen/SACCprop231.htm>  
keywords: Chaetocercus mulsant  
institution:  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
issue:  
year: 2006  
0: Stiles, F G  
title: Proposal (231) to South American Classification Committee  
source: South American Classification Committee  
type: miscellany  
tags: Chaetocercus mulsant  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

3

link: <http://www.hbw.com/species/white-bellied-woodstar-chaetocercus-mulsant>  
keywords: Chaetocercus mulsant  
0: del Hoyo, J  
1: Elliott, A  
2: Sargatal, J  
3: Christie, D A  
4: de Juana, E  
institution:  
address:  
publisher: Lynx Edicions  
accessed:  
websites:  
chapter:  
series:  
issue:  
year: 1999  
0: Z"uchner, T  
1: Boesman, P  
title: White-bellied Woodstar (Chaetocercus mulsant)  
source: Handbook of the Birds of the World Alive  
type: book\_section  
tags: Chaetocercus mulsant  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: La Estrellita Vientriblanca presenta dimorfismo sexual; el macho mide aproximadamente 7 cm de longitud, mientras que la hembra alcanza 6.4 cm. El pico tiene unos 18 mm de largo. El macho es verde brillante en la parte superior; presenta una mancha blanca detrás de cada ojo. La garganta es rosada a púrpura brillante. Las partes inferiores, faja pectoral y centro del abdomen, son predominantemente blancas contrastando con los flancos verde oscuro, parece tener un chaleco abierto. La cola es relativamente corta y puntiaguda, con las plumas exteriores negras. La hembra presenta las partes superiores de verde bronce, con la mancha postocular alargada y más notoria que la del macho. Las partes inferiores son blancas con los flancos rojizos. Su cola es más corta que la del macho, y en general, la cola corta de esta especie la hace parecer de las más pequeñas (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986). Anteriormente clasificado en el género *Acestrura* junto con *C. bombus*, *C. heliodor*, *C. astreans* y *C. berlepschi*, pero no hay evidencia en la morfología externa que justifique un tratamiento en un género separado de *C. jourdanii*; cambió con la propuesta 231 realizada al South American Classification Committee. (Stiles, 2006; Züchner & Boesman, 1999).

briefDescription:

## references

link: <http://maps.iucnredlist.org/map.html?id=22688251>

keywords: *Chaetocercus mulsant*

## editors

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

## authors

authors: BirdLife International

authors: NatureServe

title: Bird Species Distribution Maps of the World. 2012. *Chaetocercus mulsant*. The IUCN Red List of Threatened Species. Version 2014.2

source:

type: miscellany

tags: *Chaetocercus mulsant*

abstract:

## identifiers

last\_modified:

created:



group\_id:  
profile\_id:  
id:

taxonRecordName

## ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:

source:  
datatype:  
identifier:

source:  
datatype:  
identifier:

note:  
ruleConsidered:

replacementNameFor

microReference:

source:  
datatype:  
identifier:

source:  
datatype:  
identifier:

note:  
ruleConsidered:

sanctioned

microReference:

source:  
datatype:  
identifier:

source:  
datatype:  
identifier:

note:  
ruleConsidered:

laterHomonymOf

microReference:

source:  
datatype:  
identifier:

source:  
datatype:  
identifier:

note:

ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

basionym

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:

datatype:

identifier:

source:

datatype:

identifier:

source:

datatype:

identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:

microReference:

year:

publishedIn

simple:

source:  
datatype:  
identifier:  
  
specialAuthorship

authors

year  
  
simple:  
  
canonicalAuthorship

authors

year  
  
simple:  
simple: (Bourcier, 1842)  
  
canonicalName  
  
infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: Chaetocercus mulsant  
rank: SPECIES  
simple: Chaetocercus mulsant (Bourcier, 1842)

attributes  
  
nomenclaturalCode:  
isAnamorphic:  
id:

synonymsAtomized

ancillaryData

synonymStatus:  
  
synonymName  
  
anyTwo:

anyOne

providerLink:  
microReference:  
  
publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typeVoucherEntity

typeOfType:  
simple:  
microReference:  
year:  
source:  
datatype:  
identifier: (Bourcier, 1842)

[combinationAuthorship](#)

basionymAuthorship

simple:

authorship

simple:  
simple:

epithet

infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:

genus

linkType:  
ref:  
uninomial:  
simple: Acestrura mulsant  
rank:  
simple:  
nomenclaturalCode:  
isAnamorphic:  
id:

## [hierarchy](#)

## [ancillaryData](#)

dataObject:  
additionalInformation:

## [reference](#)

geoPoint:  
location:  
thumbnailURL:

## [mediaURL](#)

description:

## [subject](#)

source: The Clements Checklist

## [audience](#)

bibliographicCitation:  
rightsHolder:  
rights:  
license:  
title:  
modified: 2016-02-16T18:51:29.180Z  
created: 2016-02-16T18:51:29.180Z

## [agent](#)

homepage:  
role:  
email:  
phone:  
address:  
position:  
organisation:  
lastName:  
firstName:  
mimeType:  
dataType:  
identifier:  
parentTaxon: Chaetocercus  
higherClassification:  
infraspecificEpithet:  
specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Chaetocercus

family: Trochilidae

order: Apodiformes

classHierarchy: Aves

phylum: Chordata

kingdom: Animalia

recommended:

classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:

additionalInformation:

0

link: <http://www.hbw.com/species/white-bellied-woodstar-chaetocercus-mulsant>

keywords: Chaetocercus mulsant

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 1999

0: Z"uchner, T

1: Boesman, P

title: White-bellied Woodstar (Chaetocercus mulsant)

source: Handbook of the Birds of the World Alive

type: book\_section

tags: Chaetocercus mulsant

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
reproductionUnstructured: Solo se registró un nido a 10 m de altura en un árbol en abril en Ecuador, no se conoce más información sobre aspectos reproductivos para este colibrí (Züchner & Boesman, 1999).

0

dataObject:  
additionalInformation:

## reference

geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

## audience

rightsHolder:  
rights:  
license:  
modified:  
created:

## agent

mimeType:  
dataType:  
identifier:  
relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedBy: TraitBank  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue: Sexual  
measurementType: Tipo de Reproducción  
measurementID:

## molecularData

molecularDataUnstructured:

## migratory

migratoryUnstructured:



lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

0

dataObject:

additionalInformation:

0

link: Neotropical Birds Online [http://neotropical.birds.cornell.edu/portal/species/overview?p{\\\_}p{\\\_}spp=279736](http://neotropical.birds.cornell.edu/portal/species/overview?p{\_}p{\_}spp=279736)

keywords: Chaetocercus mulsant

0: Schulenberg, Thomas S

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2011

0: Sedgwick, C W

title: White-bellied Woodstar (Chaetocercus mulsant)

source:

type: miscellany

tags: Chaetocercus mulsant

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

interactionsUnstructured: La Estrellita Vientriblanca es una especie de colibrí principalmente nectarívoro, por su pequeño tamaño y su vuelo lento y sostenido como un abejorro, logra entrar en territorios de alimentación de otros colibríes pasando desapercibido, lo que le confiere ventajas a la hora de encontrar su alimento (Sedgwick, 2011).

feeding

0

dataObject:

additionalInformation:

0

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Aglaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigrirostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus platensis*, *Coccyzus pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus superciliaris*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxypogon guerinii*, *Oxyura jamaicensis*, *Pachyrhamphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigristrois, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus platensis, Coccyzua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreovertris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Mactornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striatocollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyrhamphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

1

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia

longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

feedingUnstructured: La Estrellita Vientriblanca se alimenta principalmente de ericáceas como Cavendishia o rubiáceas como Palicourea. Visita en muchas ocasiones flores pequeñas y compuestas y suele detenerse por momentos en el aire. Pasa largos ratos cazando insectos mientras revolotea continuamente por encima de la copa de un árbol (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986).

type: Heterótrofos - Omnívoros

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant

abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

1

link: <http://www.hbw.com/species/white-bellied-woodstar-chaetocercus-mulsant>

keywords: Chaetocercus mulsant

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 1999

0: Z"uchner, T

1: Boesman, P

title: White-bellied Woodstar (Chaetocercus mulsant)

source: Handbook of the Birds of the World Alive

type: book\_section

tags: Chaetocercus mulsant

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

2

keywords: Accipiter striatus,Actitis macularius,Adelomyia melanogenys,Aglaiocercus kingii,Amblycercus holosericeus,Ampelion rubrocristatus,Anas andium,Anas clypeata,Anas discors,Andigena nigrirostris,Anisognathus igniventris,Ardea alba,Ardea herodias,Arremon assimilis,Asio flammeus,Asio stygius,Asthenes flammulata,Asthenes fuliginosa,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Butorides virescens,Cairina moschata,Calidris bairdii,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Cathartes aura,Catharus minimus,Catharus ustulatus,Chaetocercus mulsant,Chaetura pelagica,Chalcostigma heteropogon,Chordeiles minor,Chrysomus icterocephalus,Cinclodes albidiventris,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus platensis,Coccyzus pumila,Coccyzus americanus,Coccyzus melacoryphus,Coeligena bonapartei,Colibri coruscans,Colibri thalassinus,Columba livia,Columbina talpacoti,Conirostrum rufum,Conirostrum sitticolor,Contopus cinereus,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Dendrocygna autumnalis,Dendrocygna viduata,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dolichonyx oryzivorus,Dubusia taeniata,Egretta thula,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreiventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Geothlypis philadelphia,Geranoaetus melanoleucus,Glaucidium jardinii,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza

tags: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigristrois*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus platensis*, *Coccyzus pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alhorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus*

exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Se desplaza lentamente con la cola levantada o meneándose de arriba hacia abajo rápidamente, el vuelo es bastante pausado, como suspendido en el aire (Asociación Bogotana de Ornitología, 2000). Usualmente este es un colibrí solitario en estratos medio-alto. Se posa con su pico dirigido hacia arriba. Revolotea al libar, manteniendo el pico verticalmente hacia arriba (Hilty & Brown, 1986; Züchner & Boesman, 1999). Emite una serie de chirridos muy débiles y agudos, chiri-chiri, chiri, chiri, chiri-chiri. A veces produce un zumbido al volar muy parecido al de un abejorro (Asociación Bogotana de Ornitología, 2000) o un traqueteo más agudo (Züchner & Boesman, 1999).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:



source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:  
additionalInformation:

[reference](#)

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant

[editors](#)

editors: Stiles, F G  
editors: Bohórquez, C I  
editors: Cadena, C D  
editors: De la Zerda, S  
editors: Hernández, M  
editors: Roselli, L  
editors: Kelsey, M  
editors: Valencia, I D  
editors: Knapp, D  
institution:  
edition: Primera Ed  
address:  
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca  
accessed:  
websites:  
chapter:

series:  
pages: 276  
issue:  
year: 2000

## authors

authors: Asociación Bogotana de Ornitología  
title: Aves de la Sabana de Bogotá: guía de campo  
source:  
type: book  
tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant  
abstract:

## identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

contentType:  
dataType:  
identifier:  
distributionUnstructured: La Estrellita Vientriblanca se distribuye por los Andes desde Colombia hasta Bolivia (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986). Está dispersa irregularmente en el centro y el este de los Andes de Colombia a Ecuador y Perú (Züchner & Boesman, 1999). En Colombia se encuentra en la cordillera Central y en ambas vertientes de la cordillera Oriental, se extiende desde Cundinamarca hacia el sur hasta el sureste de Nariño (Hilty & Brown, 1986).  
Se encuentra entre los 1500 y los 2800 msnm, probablemente más alto. En la mayoría de regiones está por encima de los 2200 msnm (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986).

## distributionAtomized

stateProvince: Tolima

## ancillaryData

locality:  
municipality:  
country: Colombia

temporalCoverage

endDate:  
startDate:

distributionScope

type:

## threatStatus

## ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES: Apéndice II

locality:  
municipality:  
county:  
stateProvince:  
country: Colombia  
0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:  
measurementRemarks:  
measurementMethod:

## measurementDeterminedBy

measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementType: IUCN  
measurementID:

territory

territoryUnstructured:  
areaOfOccupancy:  
extentOfOccurrence:

populationBiology

0

dataObject:  
additionalInformation:

0

link: <http://www.birdlife.org/datazone/speciesfactsheet.php?id=2109>

keywords: Chaetocercus mulsant

institution:  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
issue:  
year: 2014  
0: BirdLife International  
title: Species factsheet: Chaetocercus mulsant  
source:  
type: miscellany  
tags: Chaetocercus mulsant  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
populationBiologyUnstructured: El tamaño global de la población no ha sido cuantificado, sin embargo se considera una especie bastante común con una tendencia estable sin evidencia de declinaciones o cambios considerables (BirdLife International, 2014).

## legislation

legislationUnstructured:

## directThreats

0

dataObject:

additionalInformation:

0

link: <http://www.cites.org/esp/app/appendices.php>

keywords: Asio flammeus,Asio stygius,Buteo platypterus,Buteo swainsoni,Chaetocercus mulsant,Chalcostigma heteropogon,Colibri coruscans,Elanus leucurus,Eupsittula pertinax,Falco peregrinus,Forpus conspicillatus,Glaucidium jardinii,Lafresnaya lafresnayi,Lesbia nuna,Lesbia victoriae,Megascops choliba,Metallura tyrianthina,Tyto alba

institution:

address:

**publisher:**  
**accessed:**  
**websites:**  
**chapter:**  
**series:**  
**issue:**  
**year:** 2014  
**0:** CITES  
**title:** Apéndices CITES  
**source:**  
**type:** miscellany  
**tags:** Asio flammeus,Asio stygius,Buteo platypterus,Buteo swainsoni,Chaetocercus mulsant,Chalcostigma heteropogon,Colibri coruscans,Elanus leucurus,Eupsittula pertinax,Falco peregrinus,Forpus conspicillatus,Glaucidium jardinii,Lafresnaya lafresnayi,Lesbia nuna,Lesbia victoriae,Megascops choliba,Metallura tyrianthina,Tyto alba  
**abstract:**  
**last\_modified:**  
**created:**  
**group\_id:**  
**profile\_id:**  
**id:**  
**geoPoint:**  
**location:**  
**thumbnailURL:**  
**description:**  
**source:**  
**bibliographicCitation:**  
**rightsHolder:**  
**rights:** Asociación Bogotana de Ornitología  
**license:** Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
**modified:**  
**created:**  
**contentType:**  
**dataType:**  
**identifier:**  
**directThreatsUnstructured:** Aunque no se considera una especie amenazada se debe controlar su tráfico y comercialización (CITES, 2014).

#### ancillaryData

**dataObject:**  
**additionalInformation:**  
**geoPoint:**  
**location:**  
**thumbnailURL:**  
**description:**  
**source:**  
**bibliographicCitation:**  
**rightsHolder:**  
**rights:**  
**license:**  
**modified:**  
**created:**  
**contentType:**  
**dataType:**  
**identifier:**

## measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: La Estrellita Vientriblanca es una de las especies más pequeñas de colibríes; vuela lentamente como un abejorro, en consecuencia, puede entrar al territorio de alimentación de otros colibríes. Se encuentra principalmente en bordes de bosques y pastizales, áreas cultivadas húmedas con árboles son el hogar de estos pequeños colibríes. Buscan flores en todos los niveles y con frecuencia se posan en una percha alta a la intemperie. Macho y hembra son diferentes, ambos sexos tienen grandes manchas blancas de flanco y postocular. Los machos tienen una banda blanca en el pecho ancho, mientras que las hembras son más claras debajo (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986; Schulenberg et al., 2010; Sedgwick, 2011; Züchner & Boesman, 1999).

## associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

## ancillaryData

dataObject:

additionalInformation:

## reference

geoPoint:

location:

thumbnailURL:

mediaURL: <http://www.xeno-canto.org/130052>

description:

## subject

source: <http://www.xeno-canto.org>

## audience

bibliographicCitation:

rightsHolder:

rights:

license:  
modified:  
created:

agent

contentType:  
dataType:  
identifier:

\_id

\_bsonType: ObjectId  
id: VÃrèð lg#ç4°