Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e48a1e83c45700544e3e42

Fecha de creación: 2016-03-12T21:29:08.900Z

creation\_date: 2016-03-12T21:29:08.900Z usesManagementAndConservation

## ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: En estas avispas el escutelo se ve, de perfil, curvado regularmente formando con el metanoto una única inclinación o convexidad y nunca se encuentra elevado y proyectado sobre el metanoto. El margen posterior del metanoto se prolonga hacia el área medio-basal del propodeo como un lóbulo o ángulo agudo. El segmento abdominal I es a menudo peciolado.

briefDescription:

#### references

keywords: Agelaia angulata angulata,Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Asteloeca traili,Brachygastra albula,Brachygastra augusti,Brachygastra lecheguana,Brachygastra propodealis,Brachygastra scutellaris,Chartergellus amazonicus,Chartergellus punctatior,Chartergus chartarius,Chartergus globiventris,Chartergus metanotalis,Clypearia sulcata,Clypearia weyrauchi,Epipona tatua,Leipomeles dorsata,Metapolybia acincta,Metapolybia docilis,Metapolybia nigra,Metapolybia suffusa,Mischocyttarus adolphi,Mischocyttarus alfkenii excrucians,Mischocyttarus alfkenii trinitatis,Mischocyttarus carbonarius carbonarius,Mischocyttarus flavicans,Mischocyttarus foveatus,Mischocyttarus injucundus tingomariae,Mischocyttarus insolitus,Mischocyttarus labiatus,Mischocyttarus latissimus,Mischocyttarus metoecus,Mischocyttarus petiolatus,Mischocyttarus pseudomimeticus,Mischocyttarus punctatus,Mischocyttarus tomentosus,Mischocyttarus vaqueroi,Parachartergus amazonensis,Parachartergus colobopterus,Parachartergus fasciipennis,Parachartergus fraternus,Parachartergus fulgidipennis,Parachartergus griseus,Parachartergus richardsi,Parachartergus weyrauchi,Polybia bistriata,Polybia catillifex,Polybia eberhardae,Polybia emaciata,Polybia fastidiosuscula,Polybia jurinei,Polybia liliacea,Polybia micans,Polybia occidentalis,Polybia platycephala

sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea

#### editors

institution: address: publisher: accessed: websites: chapter: series: pages: 585 issue: year: 1978

#### authors

authors: Richards, O. W.

title: The social wasps of the Americas excluding the Vespinae

source: type: book

tags: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta. Metapolybia docilis. Metapolybia nigra. Metapolybia suffusa. Mischocyttarus adolphi. Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis.Polybia sericea.Polybia striata.Polybia velutina.Protopolybia acutiscutis.Protopolybia amarella.Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea abstract:

#### identifiers

last\_modified: created: group\_id: profile\_id:

id:

taxonRecordName

# ancillaryData

# providerSpecificData anyTwo: providerLink: publicationStatus microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: replacementNameFor microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: sanctioned microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: laterHomonymOf microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: conservedAgainst

basedOn

#### authors

## year

simple:

canonicalAuthorship

#### authors

## year

simple:

#### canonicalName

infraspecificEpithet: specificEpithet: infragenericEpithet:

linkType: ref:

uninomial:

simple: Protopolybia exigua exigua

rank: SUBSPECIES

simple: Protopolybia exigua subsp. exigua

#### attributes

nomenclaturalCode: isAnamorphic:

i4·

# synonymsAtomized

## hierarchy

# ancillaryData

dataObject:

additionalInformation:

#### reference

geoPoint: location: thumbnailURL:

## mediaURL

description:

# subject

source: NUB Generator [autonym]

#### audience

bibliographicCitation:

agent	
homepa role: email: phone: address	

firstName: mimeType: dataType:

identifier:

position: organisation: lastName:

parentTaxon: Protopolybia exigua

higherClassification: infraspecificEpithet: specificEpithet:

taxonRank: SUBSPECIES

subGenus:

genus: Protopolybia family: Vespidae order: Hymenoptera classHierarchy: Insecta phylum: Arthropoda kingdom: Animalia recommended: classification:

#### commonNameAtomized

# ancillaryData

dataObject:

additionalInformation:

#### reference

geoPoint: location:

thumbnailURL:

#### mediaURL

description:

## subject

source:

#### audience

bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
agent
mimeType: dataType: identifier: usedBy:
usedIn
distributionUnstructured: endDate: startDate:
ancillaryData
type: language: Español name: Avispa
invasiveness
invasivenessUnstructured:
reproduction
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder:
rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured: 0
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured: 0 dataObject:

location: thumbnailURL:
mediaURL
description:
subject
audience
rightsHolder: rights: license: modified: created:
agent
mimeType: dataType: identifier: relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy: TraitBank measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: Sexual measurementType: Tipo de Reproducción measurementID:
molecularData
molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions
interactionsUnstructured:
feeding
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL:

description: source: bibliographicCitation:
rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
mimeType: dataType: identifier: feedingUnstructured:
0
0
strategy: Depredadores type:
environmentalEnvelope
environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured: distance: structureDispersed: type: purpose:
behavior
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source:
bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified:
created: mimeType: dataType: identifier: behaviorUnstructured: No es activa en la defensa del nido. Suelen tomar agua de pequeños pozos y forrajean principalmente al nivel del piso.
ancillaryData

dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier: measurementOrFact relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID: annualCycle annualCycleUnstructured: habitat dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: habitatUnstructured: Habita en sabanas secas de Sudamérica. endemicAtomized

#### distribution

## ancillaryData

dataObject:

additionalInformation:

#### reference

geoPoint: location: thumbnailURL:

#### mediaURL

description:

## subject

source:

#### audience

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

#### agent

mimeType: dataType: identifier:

distributionUnstructured: Esta especie se distribuye en Colombia, Venezuela, Guyana, Guyana Francesa, Trinidad y Tobago, Ecuador, Perú, Brasil y Bolivia. Se ha registrado en zonas con altitudes que alcanzan hasta los 1000 msnm.

## distributionAtomized

stateProvince: Arauca

## ancillaryData

locality: municipality: country: Colombia

temporalCoverage

endDate: startDate:

distributionScope

type:

## threatStatus

#### territory

territoryUnstructured: areaOfOccupancy: extentOfOccurrence:

populationBiology
populationBiologyUnstructured:
legislation
legislationUnstructured:
directThreats
directThreatsUnstructured:
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID: abstract: Protopolybia exigua exigua se distribuye en Sudamérica y habita sabanas secas. En Colombia se ha reportado en las regiones de la Amazonia, Orinoquía y Pacífica. Los adultos presentan la cara posterior del propodeo redondeada; lóbulos medianos del edeago truncados; dos bandas amarillas en el mesoescuto y dos manchas o bandas anteriores en el segundo tergo abdominal.
associatedParty
role: personnelIdentifier: personnelDirectory: homepage: email: enataliagomezm@gmail.com phone: postalCode: country: Colombia state: Bogotá Distrito Capital city: Bogotá D.C. address:

organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

#### ancillaryData

#### dataObject:

additionalInformation: Información de tipos: Holotipo: hembra coleccionado en Brasil y está en el Musée National d'histoire Naturelle, Paris, Francia (MNHN).

Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

#### reference

geoPoint: location: thumbnailURL:

#### mediaURL

description:

#### subject

source:

#### audience

bibliographicCitation: rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

#### agent

\_id

mimeType: dataType: identifier:

\_bsontype: ObjectID id: VäŠ fÄW Dãä