

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d6f24c3c16479905cba915

Fecha de creación: 2016-03-02T14:01:51.067Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Relativamente común en los llanos en dehesas más secas, áreas cultivadas con árboles y en pueblos, ocasionalmente en bordes de bosque de galería (Hilty & Brown 1986). En La Guajira habita en matorral o rastrojo y pastizales (Cerrejón 2009-2010).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Inglés

name: Carib Grackle

creation\_date: 2016-03-02T14:01:51.067Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: Se reconocen ocho subespecies:

*Quiscalus lugubris contrusus* (J. L. Peters, 1925)

*Quiscalus lugubris fortirostris* Lawrence, 1868

*Quiscalus lugubris guadeloupensis* Lawrence, 1879

*Quiscalus lugubris inflexirostris* Swainson, 1838

*Quiscalus lugubris insularis* Richmond, 1896

*Quiscalus lugubris lugubris* Swainson, 1838

*Quiscalus lugubris luminosus* (Lawrence, 1878)

*Quiscalus lugubris orquillensis* (Cory, 1909)

briefDescription:

## references

taxonRecordName

## ancillaryData

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

replacementNameFor

microReference:

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:

publishedIn

simple:  
source:  
datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: Swainson, 1838

canonicalName

infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:

ref:  
uninomial:  
simple: Quiscalus lugubris  
rank: SPECIES  
simple: Quiscalus lugubris Swainson, 1838

## attributes

nomenclaturalCode:  
isAnamorphic:  
id:

## synonymsAtomized

## hierarchy

## ancillaryData

dataObject:  
additionalInformation:

## reference

geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source: The Catalogue of Life, 3rd January 2011

## audience

bibliographicCitation:  
rightsHolder:  
rights:  
license:  
title:  
modified: 2016-03-02T14:01:31.067Z  
created: 2016-03-02T14:01:31.067Z

## agent

homepage:  
role:  
email:  
phone:  
address:  
position:  
organisation:  
lastName:  
firstName:  
mimeType:  
dataType:  
identifier:

parentTaxon: Quiscalus  
higherClassification:  
infraspecificEpithet:  
specificEpithet:  
taxonRank: SPECIES  
subGenus:  
genus: Quiscalus  
family: Icteridae  
order: Passeriformes  
classHierarchy: Aves  
phylum: Chordata  
kingdom: Animalia  
recommended:  
classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:

identifier:

feedingUnstructured: En La Guajira se observó que los individuos forrajean entre el suelo y el dosel y su dieta es omnívora (Cerrejón 2009-2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Quiscalus lugubris es una especie gregaria. Pernocita en grandes y ruidosas bandadas en árboles, a menudo en parques urbanos (Hilty & Brown 1986).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:  
identifier:

measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:  
additionalInformation:

[reference](#)

geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

[agent](#)

contentType:  
dataType:  
identifier:  
distributionUnstructured: En bases de datos, literatura y monitoreos de fauna se encontró que: Esta especie se distribuye en: Guyana, Venezuela (Arctos 2012); Barbados, Dominica, Granada, Guadalupe, República Dominicana, San Vicente y Las Granadinas, Venezuela (Collections MNH SI 2012)  
Las siguientes subespecies se distribuyen en Colombia:  
Quiscalus lugubris contrusus San Vicente y Las Granadinas, (Collections MNH SI 2012)  
Quiscalus lugubris fortirostris Antigua y Barbuda (Arctos 2012); Antigua y Barbados, Barbados (Collections MNH SI 2012)  
Quiscalus lugubris guadeloupensis Dominica, Guadalupe, Martinica, Montserrat (Collections MNH SI 2012)



Quiscalus lugubris inflexirostris Santa Lucia (Collections MNH SI 2012)  
Quiscalus lugubris insularis Venezuela (Collections MNH SI 2012)  
Quiscalus lugubris lugubris Trinidad y Tobago (Arctos 2012); Guyana, Trinidad y Tobago, Venezuela (Collections MNH SI 2012)  
Quiscalus lugubris luminosus Granada, San Vicente y Las Granadinas (Collections MNH SI 2012)  
En bases de datos, literatura y monitoreos de fauna se encontró que: Esta especie se distribuye en: La Guajira (Cerrejón 2009-2010). La subespecie Quiscalus lugubris lugubris se distribuye en: Arauca, Casanare, Meta (ICN 2004).  
Se encuentra en las regiones Caribe y Orinoquía hasta los 600 m (Hilty & Brown 1986).

distributionAtomized

temporalCoverage

endDate:  
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:  
locality:  
municipality:  
county:  
stateProvince:  
country: Global  
0: IUCN  
measurementValue: Preocupación menor (LC)  
relatedTo:  
measurementRemarks:  
measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementType: IUCN  
measurementID:

territory

territoryUnstructured:  
areaOfOccupancy:  
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Los machos miden unos 27 cm mientras que las hembras miden 22 cm aproximadamente. Tienen los ojos amarillo pálido, pico negro, algo delgado y decurvado; la cola es cuneiforme. Los machos son enteramente negros con un ligero lustre púrpura. Las hembras de color pardo hollín oscuro, la garganta más pálida (Hilty & Brown 1986).

## associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

## ancillaryData

dataObject:  
additionalInformation: Hilty S. L., Brown W. L. (1986) A Guide to the Birds of Colombia Princeton, New Jersey. U. S. A. | ARCTOS COLLECTION MANAGEMENT INFORMATION SYSTEM Specimen Search | CERREJÓN Monitoreo de fauna en el Valle del Cerrejón. ASESORÍAS Y ESTUDIOS AMBIENTALES J.D.B. S.A.S. Contrato CON 00272009 | SMITHSONIAN NATIONAL MUSEUM OF NATURAL HISTORY (2012) Search Museum Collections Records | INSTITUTO DE CIENCIAS NATURALES, Facultad de Ciencias, Universidad Nacional de Colombia Colecciones en Línea | ORRELL T. (custodio) (2012) ITIS Global: The Integrated Taxonomic Information System (version Sept 12). En: Species 2000 & ITIS Catalogue of Life, 20th November 2012 | BIRDLIFE INTERNATIONAL (2012) Quiscalus lugubris. En: IUCN 2012. IUCN Red List of Threatened Species. Versión 2012.2

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

agent

contentType:  
dataType:  
identifier:

\_id  
\_bsontype: ObjectID  
id: VÖòL< G™\0'P