

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d5ea6c3c16479905cba901

Fecha de creación: 2016-03-01T19:15:58.800Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

habitatUnstructured: Relativamente común a común en terreno seco y semiabierto, sabana y potreros con árboles y arbustos dispersos, ocasionalmente en playas arenosas (Hilty & Brown 1986). En La Guajira habita en pastizales y matorrales o rastrojos (Cerrejón 2009-2010).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Inglés

name: Cattle Tyrant

creation\_date: 2016-03-01T19:15:58.800Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
fullDescriptionUnstructured: Se reconocen tres subespecies:  
Machetornis rixosa flavigularis Todd, 1912  
Machetornis rixosa obscurodorsalis Phelps & W. H. Phelps Jr, 1948  
Machetornis rixosa rixosa (Vieillot, 1819)  
briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:

identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

conservedAgainst

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:

datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion  
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:  
  
publishedIn

simple:  
source:  
datatype:  
identifier:  
  
specialAuthorship

authors

year  
  
simple:  
canonicalAuthorship

authors

year  
  
simple:  
simple: (Vieillot, 1819)  
  
canonicalName

infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: Machetornis rixosa  
rank: SPECIES  
simple: Machetornis rixosa (Vieillot, 1819)

attributes

nomenclaturalCode:  
isAnamorphic:  
id:

synonymsAtomized

ancillaryData

synonymStatus:  
synonymName

anyTwo:

anyOne

providerLink:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:

note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:

note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:

note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:

note:  
ruleConsidered:

typeNameEntity

source:  
datatype:  
identifier:

source:  
datatype:  
identifier:

source:  
datatype:  
identifier:

typeVoucherEntity

typeOfType:  
simple:  
microReference:  
year:  
source:  
datatype:  
identifier:

## [combinationAuthorship](#)

basionymAuthorship

simple:

authorship

simple:

simple: (Vieillot, 1819)

epithet

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

genus

linkType:

ref:

uninomial:

simple: Machetornis rixosus

rank: SPECIES

simple: Machetornis rixosus (Vieillot, 1819)

nomenclaturalCode:

isAnamorphic:

id:

## [hierarchy](#)

## [ancillaryData](#)

dataObject:

additionalInformation:

## [reference](#)

geoPoint:

location:

thumbnailURL:

## [mediaURL](#)

description:

## [subject](#)

source: The Catalogue of Life, 3rd January 2011

## [audience](#)

bibliographicCitation:

rightsHolder:  
rights:  
license:  
title:  
modified: 2016-03-01T19:15:38.796Z  
created: 2016-03-01T19:15:38.796Z

## agent

homepage:  
role:  
email:  
phone:  
address:  
position:  
organisation:  
lastName:  
firstName:  
mimeType:  
dataType:  
identifier:  
parentTaxon: Machetornis  
higherClassification:  
infraspecificEpithet:  
specificEpithet:  
taxonRank: SPECIES  
subGenus:  
genus: Machetornis  
family: Tyrannidae  
order: Passeriformes  
classHierarchy: Aves  
phylum: Chordata  
kingdom: Animalia  
recommended:  
classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions



interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataFormat:

identifier:

feedingUnstructured: Forrajea principalmente en el suelo; ocasionalmente vuela corto en pos de insectos (Hilty & Brown 1986). En La Guajira forrajea en el suelo y se alimenta de invertebrados (Cerrejón 2009-2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Se observan en parejas o pequeños grupos familiares; se posa en la copa de arbustos bajos o ramas más bajas de árboles. Corre bien, sigue al ganado y se le posa en el lomo (Hilty & Brown 1986). Genera un chillido agudo. Al amanecer vocaliza un trino silbado suiii sii dii (Wetmore 1926 en Hilty & Brown 1986).

## ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

## measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

## endemicAtomized

## distribution

## ancillaryData

dataObject:

additionalInformation:

## reference

geoPoint:

location:

thumbnailURL:

## mediaURL

description:

## subject

source:

audience

bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

agent

mimeType:  
dataType:  
identifier:  
distributionUnstructured: En bases de datos se encontró que:  
Esta especie se distribuye en: Argentina, Brasil, Uruguay (Collections MNH SI 2012).  
Las siguientes subespecies se distribuyen en:  
Machetornis rixosa flavigularis Colombia, Venezuela (Collections MNH SI 2012).  
Machetornis rixosa rixosa Argentina, Brasil, Paraguay (Collections MNH SI 2012). En bases de datos y monitoreos de fauna se encontró que:  
Esta especie se distribuye en: La Guajira (Cerrejón 2009-2010).  
La subespecie Machetornis rixosa flavigularis se distribuye en: Antioquia, Arauca, Bolívar, Cesar, Córdoba, La Guajira, Magdalena, Meta, Santander (ICN 2004); Antioquia, Atlántico, Bolívar, Cesar, Córdoba, La Guajira, Magdalena (Collections MNH SI 2012), hasta los 800 m (Hilty & Brown 1986).

distributionAtomized

stateProvince: Córdoba

ancillaryData

locality:  
municipality:  
country: Colombia  
  
temporalCoverage  
  
endDate:  
startDate:  
  
distributionScope  
  
type:

threatStatus

ancillaryData

threatStatusUnstructured:  
  
threatStatusAtomized  
  
apendiceCITES:  
locality:  
municipality:  
county:  
stateProvince:  
country: Global  
0: IUCN  
measurementValue: Preocupación menor (LC)

relatedTo:  
measurementRemarks:  
measurementMethod:

## measurementDeterminedBy

measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementType: IUCN  
measurementID:

## territory

territoryUnstructured:  
areaOfOccupancy:  
extentOfOccurrence:

## populationBiology

populationBiologyUnstructured:

## legislation

legislationUnstructured:

## directThreats

directThreatsUnstructured:

## ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

## measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

abstract: Mide unos 19 cm. Tiene los ojos rojos y las patas largas. Principalmente marrón oliva pálido en las partes

dorsales, con la coronilla de color gris y una cresta naranja encendido usualmente oculta (únicamente en los machos); presenta una línea ocular estrecha de color negruzco, alas y cola parduscas, la cola con márgenes pálidos; las partes ventrales amarillo brillante incluyendo las cobertoras alares internas (Hilty & Brown 1986).

## associatedParty

role:  
personnelIdentifier:  
personnelDirectory:  
homepage:  
email: enataliagomez@gmail.com  
phone:  
postalCode:  
country: Colombia  
state: Bogotá Distrito Capital  
city: Bogotá D.C.  
address:  
organisation:  
position:  
lastName: Gómez Melendro  
firstName: Edith Natalia

## ancillaryData

dataObject:  
additionalInformation: Registros biológicos: En el portal de Global Biodiversity Information Facility se reporta que para Colombia hay 290 registros. GBIF  
Metadatos: GBIF  
Créditos específicos: Ficha elaborada con el apoyo de Carbones del Cerrejón Limited.  
Otros recursos en Internet: Arctos, Avibase, Colecciones en Línea, EoL, iSpecies, IUCN Red List, SI NMNH, xenocanto

## reference

geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:  
rightsHolder:  
rights: Equipo Coordinador del SiB Colombia  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

mimeType:  
dataType:  
identifier:

\_id

\_bsontype: ObjectId

id:  $V \in G^{\text{TM}} \setminus 0^i$