Reporte de Fichas para el usuario: edith gómez Ficha ID: 56c7b472f0106c67230e7404 Fecha de creación: undefined annualCycles annualCycleUnstructured: habitats 0 dataObject: additionalInformation: link: http://www.hbw.com/node/56983 keywords: Myornis senilis 0: del Hoyo, J 1: Elliott, A 2: Sargatal, J 3: Christie, D A 4: de Juana, E institution: address: publisher: Lynx Edicions accessed: websites: chapter: series: issue: year: 2003 0: Krabbe, N K 1: Schulenberg, T S title: Ash-coloured Tapaculo (Myornis senilis) source: Handbook of the Birds of the World Alive type: book\_section tags: Myornis senilis abstract: last\_modified: created: group\_id: profile\_id: geoPoint: location: thumbnailURL: description:

source:

modified: created: mimeType: dataType: identifier:

rightsHolder:

bibliographicCitation:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

habitatUnstructured: La especie habita en densos matorrales de bambú Chusquea o caña Neurolepis en las fronteras forestales y bosques montanos húmedos (Krabbe & Schulenberg, 2003).

#### commonNamesAtomized

#### ancillaryData

usedBy:

usedIn

distributionUnstructured:

endDate: startDate:

# ancillaryData

type:

language: Alemán

name: Aschgrauer Tapaculo

usesManagementAndConservation

# ancillaryData

managementAndConservationUnstructured:

# managementAndConservationAtomized

identificationKeys

fullDescription

O

dataObject: additionalInformation:

0

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus

superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Myornis senilis.Nomonyx dominicus.Nycticorax nycticorax.Ochthoeca cinnamomeiyentris.Ochthoeca diadema.Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus. Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution: address:

accessed:

publisher: Princeton University Press

websites: chapter: series: pages: 836 issue: year: 1986 0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus

nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Myornis senilis, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor.Ochthoeca rufipectoralis.Orochelidon murina.Oxypogon querinii.Oxyura jamaicensis.Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: Mide 14.5 cm. Pico y patas oscuras. El macho es gris oscuro, ligeramente pálido por debajo; cola notablemente larga (6 cm) y generalmente permanece erguida hacia arriba a la altura del pico. La hembra es similar pero las plumas de la espalda y partes inferiores cafés. Los juveniles café canela por encima, leonado ocre por debajo con barras subterminales negras sobre las plumas terciarias, algunas veces también en las coberteras (Hilty & Brown, 1986).

briefDescription:

#### references

link: http://www.birdlife.org/datazone/species/factsheet/22703483

keywords: Myornis senilis

# editors

institution: address:

publisher: accessed: websites: chapter: series: issue: year: 2014

# authors

authors: BirdLife International

title: Species factsheet: Myornis senilis

source:

type: miscellany tags: Myornis senilis

abstract:

# identifiers

last\_modified: created: group\_id: profile\_id: id:

taxonRecordName

# ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

replacementNameFor

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

sanctioned

identifier: source: datatype: identifier: note: ruleConsidered: laterHomonymOf microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: conservedAgainst basedOn microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: basionym microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: spellingCorrentionOf typeNameEntity source: datatype: identifier: source: datatype: identifier:

microReference:

source: datatype:

source: datatype: identifier:
typificacion
typeOfType:
lectotypeMicroReferenceVoucher
lectotypePublicationVoucher
voucherReference
simple: microReference: year:
publishedIn
simple: source: datatype: identifier:
specialAuthorship
authors
year
simple:
canonicalAuthorship
authors
year
simple: simple: (Lafresnaye, 1840)
canonicalName
infraspecificEpithet: specificEpithet: infragenericEpithet: linkType: ref:
uninomial: simple: Myornis senilis rank: SPECIES simple: Myornis senilis (Lafresnaye, 1840)
attributes
nomenclaturalCode: isAnamorphic: id:

# synonymsAtomized

# ancillaryData

dataObject:

additionalInformation:

#### reference

geoPoint: location:

thumbnailURL:

# mediaURL

description:

# subject

source:

# audience

bibliographicCitation:

rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

# agent

mimeType: dataType:

identifier:

synonymStatus:

synonymName

anyTwo:

# anyOne

providerLink: microReference:

publishedIn

source: datatype: identifier:

#### reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

publishedIn

source: datatype: identifier: reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

# publishedIn

source: datatype: identifier:

#### reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

# publishedIn

source: datatype: identifier:

# reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

# publishedIn

source: datatype: identifier:

# reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

# publishedIn

source: datatype:

i do ostifi o u
identifier:
reletedName
source: datatype: identifier: note: ruleConsidered:
typeNameEntity
source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:
typeVoucherEntity
typeOfType: simple: microReference: year: source: datatype: identifier: (Lafresnaye, 1840)
combinationAuthorship
basionymAuthorship
simple:
authorship
simple: simple:
epithet
infraspecificEpithet: specificEpithet: infragenericEpithet:
genus
linkType: ref: uninomial: simple: Scytalopus senilis rank: simple: nomenclaturalCode: isAnamorphic: id:

# hierarchy

# ancillaryData

dataObject:

additionalInformation:

#### reference

geoPoint: location:

thumbnailURL:

# mediaURL

description:

# subject

source: The Catalogue of Life, 3rd January 2011

# audience

bibliographicCitation:

rigthsHolder: rights:

license:

title:

modified: 2016-02-22T18:38:16.177Z created: 2016-02-22T18:38:16.177Z

# agent

homepage:

role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:

parentTaxon: Myornis higherClassification: infraspecificEpithet: specificEpithet: taxonRank: SPECIES

subGenus: genus: Myornis

genus: Myornis
family: Rhinocryptidae
order: Passeriformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:

classification: invasiveness invasivenessUnstructured: reproduction dataObject: additionalInformation: link: http://www.hbw.com/node/56983 keywords: Myornis senilis 0: del Hoyo, J 1: Elliott, A 2: Sargatal, J 3: Christie, D A 4: de Juana, E institution: address: publisher: Lynx Edicions accessed: websites: chapter: series: issue: year: 2003 0: Krabbe, N K 1: Schulenberg, T S title: Ash-coloured Tapaculo (Myornis senilis) source: Handbook of the Birds of the World Alive type: book\_section tags: Myornis senilis abstract: last\_modified: created: group\_id: profile\_id: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured: Dentro de los datos del periodo de reproducción se reportan polluelos en junio al sureste

de Ecuador y en agosto en Perú (La Libertad); breve período de actividad de canto sugiere que puede ser un solo periodo de reproducción en el año (Krabbe & Schulenberg, 2003). molecularData molecularDataUnstructured: migratory migratoryUnstructured: lifeForm lifeFormUnstructured: lifeCycle lifeCycleUnstructured: interactions interactionsUnstructured: feeding feedingUnstructured: environmentalEnvelope environmental Envelope Unstructured:ecologicalSignificance ecologicalSignificanceUnstructured: dispersal dispersalUnstructured: distance: structureDispersed: type: purpose: behavior 0 dataObject: additionalInformation: link: http://www.hbw.com/node/56983 keywords: Myornis senilis 0: del Hoyo, J 1: Elliott, A 2: Sargatal, J 3: Christie, D A 4: de Juana, E institution: address: publisher: Lynx Edicions

accessed: websites:

chapter: series: issue: year: 2003 0: Krabbe, N K 1: Schulenberg, T S

title: Ash-coloured Tapaculo (Myornis senilis) source: Handbook of the Birds of the World Alive

type: book\_section tags: Myornis senilis

abstract: last\_modified: created: group\_id: profile\_id: id:

1

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis.Catamenia analis.Catamenia inornata.Cathartes aura.Catharus minimus.Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Myornis senilis, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon querinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa

solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution: address:

publisher: Princeton University Press

accessed: websites: chapter: series: pages: 836 issue: year: 1986 0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus.Cinclodes albidiventris.Cinclus leucocephalus.Cinnycerthia unirufa.Cistothorus apolinari.Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnavii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria guitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Myornis senilis, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris. Tangara vitriolina. Thraupis episcopus. Thraupis palmarum. Tringa flavipes. Tringa melanoleuca. Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus

melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis abstract: last_modified: created: group_id: profile_id: id: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: behaviorUnstructured: Brinca y se desplaza entre la densa maraña de bambú o caña, generalmente desde muy cerca al suelo hasta 4 m de altura; rara vez en el suelo entre las hojas de bambú muertas (Krabbe & Schulenberg, 2003). En Perú emite notas bajas kep y trinos cortos y bajos (Hilty & Brown, 1986).
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID:

#### endemicAtomized

# distribution

# ancillaryData

dataObject:

additionalInformation:

#### reference

link: http://maps.iucnredlist.org/map.html?id=22703483

keywords: Myornis senilis

# editors

institution: address: publisher: accessed: websites: chapter: series: issue: year: 2014

# authors

authors: BirdLife International and NatureServe

title: Bird Species Distribution Maps of the World. 2012. Myornis senilis. The IUCN Red List of Threatened Species.

Version 2014.3

source:

type: miscellany tags: Myornis senilis

abstract:

# identifiers

last\_modified:

created:

group\_id: profile\_id:

:4.

geoPoint:

location:

thumbnailURL:

# mediaURL

description:

# subject

source:

# audience

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

#### agent

mimeType: dataType: identifier:

distributionUnstructured: La especie se distribuye principalmente en Sudamérica desde Colombia hasta Perú. En Colombia se registra en la cordillera Occidental en el páramo del Sol, municipio de Urrao, al occidente de la cordillera Central en Caldas (Nevado del Ruiz), en la cordillera Oriental en Norte de Santander (Páramo de Tamá) y en Cundinamarca (BirdLife International and NatureServe, 2014; Flórez et al, 2004; Hilty & Brown, 1986). La especie se encuentra principalmente entre los 2400 y los 3100 msnm (Hilty & Brown, 1986).

#### distributionAtomized

stateProvince: Tolima

# ancillaryData

locality: municipality: country: Colombia

temporalCoverage

endDate: startDate:

distributionScope

type:

threatStatus

# ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality: municipality: county: stateProvince: country: Global 0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks: measurementMethod:

#### measurementDeterminedBy

measurementDeterminedDate:

measurementUnit: measurementAccuracy: measurementType: IUCN measurementID: territory territoryUnstructured: areaOfOccupancy: extentOfOccurrence: populationBiology dataObject: additionalInformation: link: http://www.birdlife.org/datazone/species/factsheet/22703483 keywords: Myornis senilis institution: address: publisher: accessed: websites: chapter: series: issue: year: 2014 0: BirdLife International title: Species factsheet: Myornis senilis source: type: miscellany tags: Myornis senilis abstract: last\_modified: created: group\_id: profile\_id: id: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: populationBiologyUnstructured: El tamaño de la población mundial no se ha cuantificado, pero esta especie es descrita como 'rara' (BirdLife International, 2014). legislation

legislationUnstructured:

# directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint: location:

thumbnailURL: description:

source:

bibliographicCitation:

rightsHolder:

rights:
license:
modified:
created:
mimeType:
dataType:

measurementOrFact

relatedTo:

identifier:

measurementRemarks: measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: El Tapaculo Ceniza es un ave oscura y de tamaño mediano con una cola larga que generalmente permanece erguida hacia arriba a la altura del pico. Se encuentra en densos matorrales montanos húmedos de zonas altas y relativamente inexploradas. En consecuencia, se sabe muy poco acerca de la ecología de forrajeo y reproducción de la especie. En ocasiones se ubica en el género Scytalopus, y todavía hay un debate activo en cuanto a que género pertenece esta especie. Aunque es poco común a lo largo de su rango de distribución, el Tapaculo Ceniza aún no ha sido incluido en las listas de especies amenazadas (The Cornell Lab of Ornithology, 2010).

# associatedParty

role:

personnelldentifier: personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone: postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro

firstName: Edith Natalia ancillaryData dataObject: additionalInformation: reference geoPoint: location: thumbnailURL: mediaURL: http://www.xeno-canto.org/128345 description: subject source: http://www.xeno-canto.org audience bibliographicCitation: rightsHolder: rights: license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: agent mimeType:

mimeType: dataType: identifier:

\_id

\_bsontype: ObjectID id: VÇ´rð lg#çD