

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e44e3883c45700544e3e2a

Fecha de creación: 2016-03-12T17:13:35.249Z

creation_date: 2016-03-12T17:13:35.249Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

fullDescriptionUnstructured: Ojos claramente peludos; espacio malar tan largo como el ancho de la fosa antenal; tórax y propodeo con dorada pubescencia erecta; mesoescuto sin punteaduras; especie negra.

briefDescription:

[references](#)

keywords: Phylloderma stenops, Polybia sericea

[editors](#)

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 624--625

issue:

volume: 51

year: 1970

[authors](#)

authors: Jeanne, R. L.
title: Note on a bat (*Phylloderma stenops*) preying upon the brood of a social wasp
source: Journal of Mammalogy
type: article
tags: *Phylloderma stenops*,*Polybia sericea*
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Olivier, 1792)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Polybia sericea
rank: SPECIES
simple: Polybia sericea (Olivier, 1792)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

ancillaryData

synonymStatus:
synonymName
anyTwo:

anyOne

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:

ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:
datatype:
identifier:

combinationAuthorship

basionymAuthorship

simple:
authorship

simple:
simple: de Saussure, 1853

epithet

infraspecificEpithet:
specificEpithet:
infragenericEpithet:

genus

linkType:
ref:
uninomial:
simple: Apoica cubitalis
rank: SPECIES
simple: Apoica cubitalis de Saussure, 1853
nomenclaturalCode:
isAnamorphic:
id:

[hierarchy](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-03-12T17:13:15.249Z
created: 2016-03-12T17:13:15.249Z

[agent](#)

homepage:
role:
email:
phone:
address:
position:

organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Polybia
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Polybia
family: Eumenidae
order: Hymenoptera
classHierarchy: Insecta
phylum: Arthropoda
kingdom: Animalia
recommended:
classification:

commonNameAtomized

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

mimeType:
dataType:
identifier:
usedBy:

usedIn

distributionUnstructured: Meta - Colombia
endDate:
startDate:

ancillaryData

type:
language: Español
name: Vaquera

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:

0

keywords: Agelaia angulata angulata,Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Asteloea traili,Brachygastra albula,Brachygastra augusti,Brachygastra lecheguana,Brachygastra propodealis,Brachygastra scutellaris,Brachymenes wagnerianus,Chartergellus amazonicus,Chartergellus punctatior,Chartergus chartarius,Chartergus globiventris,Chartergus metanotalis,Clypearia sulcata,Clypearia weyrauchi,Epipona guerini,Epipona tatua,Eumenes versicolor filiformis,Leipomeles dorsata,Metapolybia acincta,Metapolybia docilis,Metapolybia nigra,Metapolybia suffusa,Mischocyttarus adolphi,Mischocyttarus alfkeni,Mischocyttarus alfkenii excrucians,Mischocyttarus alfkenii trinitatis,Mischocyttarus carbonarius carbonarius,Mischocyttarus flavicans,Mischocyttarus foveatus,Mischocyttarus injucundus tingomariae,Mischocyttarus insolitus,Mischocyttarus labiatus,Mischocyttarus latissimus,Mischocyttarus metoecus,Mischocyttarus petiolatus,Mischocyttarus pseudomimeticus,Mischocyttarus punctatus,Mischocyttarus tomentosus,Mischocyttarus vaqueroi,Omicron aequale nigrum,Omicron conclamatum,Omicron criticum pulchripictum,Omicron foxi,Omicron globicolle,Omicron graculum atripes,Omicron lustratum,Omicron nanum incarum,Omicron paranympus,Omicron ypsilon,Pachymenes ghilianii olivaceus,Pachymenes orellanae orellanae,Pachymenes orellanae vardi,Parachartergus amazonensis,Parachartergus colobopterus,Parachartergus fasciipennis,Parachartergus fraternus,Parachartergus fulgidipennis,Parachartergus griseus,Parachartergus richardsi,Parachartergus smithii,Parachartergus weyrauchi,Pararhaphidoglossa clypeolaris,Pararhaphidoglossa confluenta confluenta,Pararhaphidoglossa cressoniana,Pararhaphidoglossa proxima,Polybia bistriata,Polybia catillifex,Polybia eberhardae,Polybia emaciata,Polybia fastidiosuscula,Polybia jurinei,Polybia liliacea,Polybia micans,Polybia occidentalis,Polybia platycephala sylvestris,Polybia quadricincta,Polybia raii raii,Polybia rejecta,Polybia scrobalis pronotalis,Polybia scrobalis scrobalis,Polybia sericea,Polybia striata,Polybia velutina,Protopolybia acutiscutis,Protopolybia amarella,Protopolybia chartergoides chartergoides,Protopolybia exigua binominata,Protopolybia exigua exigua,Protopolybia fuscatus,Protopolybia rugulosa,Protopolybia sedula,Pseudopolybia compressa,Pseudopolybia vespiceps,Santamenes peregrinus,Synoeca septentrionalis,Synoeca surinama,Synoeca virginea,Synoecoides depressus,Zethus brasiliensis fuscatus,Zethus mexicanus lugubris,Zethus mexicanus mexicanus
institution: Universidad Nacional de Colombia
address:
publisher:
accessed:
websites:
chapter:
series:
issue:
year: 1997

0: Sarmiento, C. E.

title: Véspidos de Colombia (Hymenoptera: Vespidae)

source:

type: thesis

tags: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca trilli, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Brachymenes wagnerianus, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona guerini, Epipona tatua, Eumenes versicolor filiformis, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Omicron aequale nigrum, Omicron conclamatum, Omicron criticum pulchripictum, Omicron foxi, Omicron globicollis, Omicron graculum atripes, Omicron lustratum, Omicron nanum incarum, Omicron paranympus, Omicron ypsilon, Pachymenes ghilianii olivaceus, Pachymenes orellanae orellanae, Pachymenes orellanae vardyi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus smithii, Parachartergus weyrauchi, Pararhaphidoglossa clypeolaris, Pararhaphidoglossa confluenta confluenta, Pararhaphidoglossa cressoniana, Pararhaphidoglossa proxima, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raii raii, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Santamenes peregrinus, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea, Synoecoides depressus, Zethus brasiliensis fuscatus, Zethus mexicanus lugubris, Zethus mexicanus mexicanus

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

reproductionUnstructured: El nido tiene forma globosa y es de color marrón rojizo o marrón oscuro que puede medir 36 cm de largo por 11 cm de ancho, presenta 11 panales en su interior; una colonia puede contener 900 individuos. En las llanuras es posible encontrar nidos sujetos a restos de troncos secos cubiertos por pastizales (Sarmiento, 1997).

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:
subject

audience

rightsHolder:
rights:
license:
modified:
created:

agent

mimeType:
dataType:
identifier:
relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedBy: TraitBank
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue: Sexual
measurementType: Tipo de Reproducción
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

0

dataObject:
additionalInformation:

keywords: *Agelaia angulata angulata*, *Agelaia angulicollis*, *Agelaia brevistigma*, *Agelaia cajennensis*, *Agelaia fulvofasciata*, *Agelaia hamiltoni*, *Agelaia testacea*, *Angiopolybia obidensis*, *Angiopolybia pallens*, *Angiopolybia paraensis*, *Angiopolybia zischkai*, *Apoica thoracica*, *Asteloeca trilli*, *Brachygastra albula*, *Brachygastra augusti*, *Brachygastra lecheguana*, *Brachygastra propodealis*, *Brachygastra scutellaris*, *Chartergellus amazonicus*, *Chartergellus punctatior*, *Chartergus chartarius*, *Chartergus globiventris*, *Chartergus metanotalis*, *Clypearia sulcata*, *Clypearia weyrauchi*, *Epipona tatua*, *Leipomeles dorsata*, *Metapolybia acincta*, *Metapolybia docilis*, *Metapolybia nigra*, *Metapolybia suffusa*, *Mischocyttarus adolphi*, *Mischocyttarus alfkeni*, *Mischocyttarus alfkenii excrucians*, *Mischocyttarus alfkenii trinitatis*, *Mischocyttarus carbonarius carbonarius*, *Mischocyttarus flavicans*, *Mischocyttarus foveatus*, *Mischocyttarus injucundus tingomariae*, *Mischocyttarus insolitus*, *Mischocyttarus labiatus*, *Mischocyttarus latissimus*, *Mischocyttarus metoecus*, *Mischocyttarus petiolatus*, *Mischocyttarus pseudomimeticus*, *Mischocyttarus punctatus*, *Mischocyttarus tomentosus*, *Mischocyttarus vaqueroi*, *Parachartergus amazonensis*, *Parachartergus colobopterus*, *Parachartergus fasciipennis*, *Parachartergus fraternus*, *Parachartergus fulgidipennis*, *Parachartergus griseus*, *Parachartergus richardsi*, *Parachartergus weyrauchi*, *Polybia bistriata*, *Polybia catillifex*, *Polybia eberhardae*, *Polybia emaciata*, *Polybia fastidiosuscula*, *Polybia jurinei*, *Polybia liliacea*, *Polybia micans*, *Polybia occidentalis*, *Polybia platycephala sylvestris*, *Polybia quadricincta*, *Polybia raii raii*, *Polybia rejecta*, *Polybia scrobalis pronotalis*, *Polybia scrobalis scrobalis*, *Polybia sericea*, *Polybia striata*, *Polybia velutina*, *Protopolybia acutiscutis*, *Protopolybia amarella*, *Protopolybia chartergoides chartergoides*, *Protopolybia exigua binominata*, *Protopolybia exigua exigua*, *Protopolybia fuscatus*, *Protopolybia rugulosa*, *Protopolybia sedula*, *Pseudopolybia compressa*, *Pseudopolybia vespiceps*, *Synoeca septentrionalis*, *Synoeca surinama*, *Synoeca virginea*

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 585

issue:

year: 1978

O: Richards, O. W.

title: The social wasps of the Americas excluding the Vespinae

source:

type: book

tags: *Agelaia angulata angulata*, *Agelaia angulicollis*, *Agelaia brevistigma*, *Agelaia cajennensis*, *Agelaia fulvofasciata*, *Agelaia hamiltoni*, *Agelaia testacea*, *Angiopolybia obidensis*, *Angiopolybia pallens*, *Angiopolybia paraensis*, *Angiopolybia zischkai*, *Apoica thoracica*, *Asteloeca trilli*, *Brachygastra albula*, *Brachygastra augusti*, *Brachygastra lecheguana*, *Brachygastra propodealis*, *Brachygastra scutellaris*, *Chartergellus amazonicus*, *Chartergellus punctatior*, *Chartergus chartarius*, *Chartergus globiventris*, *Chartergus metanotalis*, *Clypearia sulcata*, *Clypearia weyrauchi*, *Epipona tatua*, *Leipomeles dorsata*, *Metapolybia acincta*, *Metapolybia docilis*, *Metapolybia nigra*, *Metapolybia suffusa*, *Mischocyttarus adolphi*, *Mischocyttarus alfkeni*, *Mischocyttarus alfkenii excrucians*, *Mischocyttarus alfkenii trinitatis*, *Mischocyttarus carbonarius carbonarius*, *Mischocyttarus flavicans*, *Mischocyttarus foveatus*, *Mischocyttarus injucundus tingomariae*, *Mischocyttarus insolitus*, *Mischocyttarus labiatus*, *Mischocyttarus latissimus*, *Mischocyttarus metoecus*, *Mischocyttarus petiolatus*, *Mischocyttarus pseudomimeticus*, *Mischocyttarus punctatus*, *Mischocyttarus tomentosus*, *Mischocyttarus vaqueroi*, *Parachartergus amazonensis*, *Parachartergus colobopterus*, *Parachartergus fasciipennis*, *Parachartergus fraternus*, *Parachartergus fulgidipennis*, *Parachartergus griseus*, *Parachartergus richardsi*, *Parachartergus weyrauchi*, *Polybia bistriata*, *Polybia catillifex*, *Polybia eberhardae*, *Polybia emaciata*, *Polybia fastidiosuscula*, *Polybia jurinei*, *Polybia liliacea*, *Polybia micans*, *Polybia occidentalis*, *Polybia platycephala sylvestris*, *Polybia quadricincta*, *Polybia raii raii*, *Polybia rejecta*, *Polybia scrobalis pronotalis*, *Polybia scrobalis scrobalis*, *Polybia sericea*, *Polybia striata*, *Polybia velutina*, *Protopolybia acutiscutis*, *Protopolybia amarella*, *Protopolybia chartergoides chartergoides*, *Protopolybia exigua binominata*, *Protopolybia exigua exigua*, *Protopolybia fuscatus*, *Protopolybia rugulosa*, *Protopolybia sedula*, *Pseudopolybia compressa*, *Pseudopolybia vespiceps*, *Synoeca septentrionalis*, *Synoeca surinama*, *Synoeca virginea*

abstract:
last_modified:
created:
group_id:
profile_id:
id:

1

keywords: Phylloderma stenops,Polybia sericea

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 624--625

issue:

volume: 51

year: 1970

O: Jeanne, R. L.

title: Note on a bat (Phylloderma stenops) preying upon the brood of a social wasp

source: Journal of Mammalogy

type: article

tags: Phylloderma stenops,Polybia sericea

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

interactionsUnstructured: Sus nidos se encuentran suspendidos de ramas de cítricos (Citrus sp.) a alturas que varían desde 1 m hasta 12 m (Richards, 1978).

Se han encontrado obreras parasitadas por Strepsiptera (Braxisenos brasiliensis). El murciélago Phylloderma stenops se alimenta de larvas y pupas, ataca el nido varias noches consecutivas hasta destruir completamente la colonia (Jeanne, 1970). La especie presenta varios véspidos miméticos: Polybia chrysothorax, Mischocyttarus marginatus, Polistes subsericeus, Montezumia nigriceps, Montezumia spinolae, Montezumia sp. y Pachymenes sp. (Richards, 1978).

feeding

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured:

0

0

strategy: Depredadores

type:

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

0

keywords: Phylloderma stenops, Polybia sericea

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

pages: 624--625

issue:

volume: 51
year: 1970
0: Jeanne, R. L.
title: Note on a bat (*Phylloderma stenops*) preying upon the brood of a social wasp
source: Journal of Mammalogy
type: article
tags: *Phylloderma stenops*, *Polybia sericea*
abstract:
last_modified:
created:
group_id:
profile_id:
id:

1

keywords: *Agelaia angulata angulata*, *Agelaia angulicollis*, *Agelaia brevistigma*, *Agelaia cajennensis*, *Agelaia fulvofasciata*, *Agelaia hamiltoni*, *Agelaia testacea*, *Angiopolybia obidensis*, *Angiopolybia pallens*, *Angiopolybia paraensis*, *Angiopolybia zischkai*, *Apoica thoracica*, *Asteloeca trilli*, *Brachygastra albula*, *Brachygastra augusti*, *Brachygastra lecheguana*, *Brachygastra propodealis*, *Brachygastra scutellaris*, *Chartergellus amazonicus*, *Chartergellus punctatior*, *Chartergus chartarius*, *Chartergus globiventris*, *Chartergus metanotalis*, *Clypearia sulcata*, *Clypearia weyrauchi*, *Epipona tatua*, *Leipomeles dorsata*, *Metapolybia acincta*, *Metapolybia docilis*, *Metapolybia nigra*, *Metapolybia suffusa*, *Mischocyttarus adolphi*, *Mischocyttarus alfkeni*, *Mischocyttarus alfkenii excrucians*, *Mischocyttarus alfkenii trinitatis*, *Mischocyttarus carbonarius carbonarius*, *Mischocyttarus flavicans*, *Mischocyttarus foveatus*, *Mischocyttarus injucundus tingomariae*, *Mischocyttarus insolitus*, *Mischocyttarus labiatus*, *Mischocyttarus latissimus*, *Mischocyttarus metoecus*, *Mischocyttarus petiolatus*, *Mischocyttarus pseudomimeticus*, *Mischocyttarus punctatus*, *Mischocyttarus tomentosus*, *Mischocyttarus vaqueroi*, *Parachartergus amazonensis*, *Parachartergus colobopterus*, *Parachartergus fasciipennis*, *Parachartergus fraternus*, *Parachartergus fulgidipennis*, *Parachartergus griseus*, *Parachartergus richardsi*, *Parachartergus weyrauchi*, *Polybia bistriata*, *Polybia catillifex*, *Polybia eberhardae*, *Polybia emaciata*, *Polybia fastidiosuscula*, *Polybia jurinei*, *Polybia liliacea*, *Polybia micans*, *Polybia occidentalis*, *Polybia platycephala sylvestris*, *Polybia quadricincta*, *Polybia raii raii*, *Polybia rejecta*, *Polybia scrobalis pronotalis*, *Polybia scrobalis scrobalis*, *Polybia sericea*, *Polybia striata*, *Polybia velutina*, *Protopolybia acutiscutis*, *Protopolybia amarella*, *Protopolybia chartergoides chartergoides*, *Protopolybia exigua binominata*, *Protopolybia exigua exigua*, *Protopolybia fuscatus*, *Protopolybia rugulosa*, *Protopolybia sedula*, *Pseudopolybia compressa*, *Pseudopolybia vespiceps*, *Synoeca septentrionalis*, *Synoeca surinama*, *Synoeca virginea*

institution:
address:
publisher:
accessed:
websites:
chapter:
series:
pages: 585
issue:
year: 1978
0: Richards, O. W.
title: The social wasps of the Americas excluding the Vespinae
source:
type: book
tags: *Agelaia angulata angulata*, *Agelaia angulicollis*, *Agelaia brevistigma*, *Agelaia cajennensis*, *Agelaia fulvofasciata*, *Agelaia hamiltoni*, *Agelaia testacea*, *Angiopolybia obidensis*, *Angiopolybia pallens*, *Angiopolybia paraensis*, *Angiopolybia zischkai*, *Apoica thoracica*, *Asteloeca trilli*, *Brachygastra albula*, *Brachygastra augusti*, *Brachygastra lecheguana*, *Brachygastra propodealis*, *Brachygastra scutellaris*, *Chartergellus amazonicus*, *Chartergellus punctatior*, *Chartergus chartarius*, *Chartergus globiventris*, *Chartergus metanotalis*, *Clypearia sulcata*, *Clypearia weyrauchi*, *Epipona tatua*, *Leipomeles dorsata*, *Metapolybia*

acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistrata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raii raii, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Según Richards (1978) almacenan pequeñas cantidades de néctar en sus nidos. Forrajea en dos picos horarios, 7-13 y 16-17 horas, cuando capturan una larva pequeña la malaxan y la llevan al nido, pero si es grande, cortan la parte anterior y la malaxan y luego ensartan el resto en una rama o pasto elevado del piso, llevan al nido la primer parte y regresan poco tiempo después para recolectar lo demás; ésta estrategia de dejar el alimento en un parte elevada reduce la probabilidad de que las hormigas la encuentren mientras retornan por él; el retorno es posible mediante la localización de puntos destacados del lugar; la captura de la presa depende de estímulos olfativos y visuales (Jeanne, 1970).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

annualCycle

annualCycleUnstructured:

habitat

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
habitatUnstructured: Esta especie habita pastizales y bosques secos de Sudamérica.

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:

dataType:

identifier:

distributionUnstructured: Esta especie se distribuye en Perú, Colombia, Venezuela, Guyana, Surinám, Guyana Francesa, Brasil, Paraguay, Uruguay, Argentina y Bolivia. Ha sido registrada en zonas con altitudes que alcanzan hasta los 700 msnm.

distributionAtomized

stateProvince: Arauca

ancillaryData

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Polybia sericea pertenece a la familia Eumenidae y habita pastizales y bosques secos de Sudamérica. Presenta hábitos depredadores y comportamiento social. Vive en colonias y estas pueden contener hasta 900 individuos. Se alimenta de néctar y presas.

associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation: Información de tipos: El tipo está extraviado.

Registros biológicos: Colección Entomológica, Instituto de Ciencias Naturales, Universidad Nacional de Colombia. Bogotá, Colombia <http://www.biovirtual.unal.edu.co/coleccion/datos/animalia/consultasMammalia.jsp?clase=4>
Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto “Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC”.

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

mimeType:
dataType:
identifier:

_id

_bsontype: ObjectID
id: VãN8fÄW Dãã