

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c7c381f0106c67230e7418

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus

multis,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dubusia taeniata,Egretta caerulea,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreiventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Gallinula melanops,Geothlypis philadelphia,Glaucidium jardinii,Grallaria ruficapilla,Hemispingus superciliosus,Hemispingus verticalis,Henicorhina leucophrys,Hirundo rustica,Icterus chrysater,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Leiothlypis peregrina,Lesbia nuna,Lesbia victoriae,Leucophaeus atricilla

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

link: http://awsassets.panda.org/downloads/migratorias{_}aves{_}42{_}final.pdf

keywords: Leucophaeus atricilla

0: Naranjo, L G

1: Amaya, J D

2: Eusse-González, D

3: Cifuentes-Sarmiento, Y

institution:

address:

publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia

accessed:

websites:

chapter:

series:

pages: 708

issue:

volume: 1

year: 2012

0: Zamudio, Jeisson

title: Leucophaeus atricilla

source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves

type: book_section

tags: Leucophaeus atricilla

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotá de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:
created:
mimeType:
dataType:
identifier:
habitatUnstructured: Esta especie habita en playas arenosas, bajos areno-fangosos, zonas de manglar, puertos y pueblos costeros (Zamudio, 2012); también usa áreas de agua abierta en lagunas o humedales, sobre postes, puentes o tocones en el agua (Asociación Bogotana de Ornitología, 2000).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:
usedIn
distributionUnstructured:
endDate:
startDate:

[ancillaryData](#)

type:
language: Español
name: Gaviota reidora
usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:
additionalInformation:

0

link: http://awsassets.panda.org/downloads/migratorias{_}aves{_}42{_}final.pdf
keywords: Leucophaeus atricilla
0: Naranjo, L G
1: Amaya, J D
2: Eusse-González, D
3: Cifuentes-Sarmiento, Y
institution:
address:
publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia
accessed:
websites:
chapter:

series:
pages: 708
issue:
volume: 1
year: 2012
0: Zamudio, Jeisson
title: Leucophaeus atricilla
source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves
type: book_section
tags: Leucophaeus atricilla
abstract:
last_modified:
created:
group_id:
profile_id:
id:

1

link: <http://www.museum.lsu.edu/~Rensen/SACCprop250.html>
keywords: Leucophaeus atricilla
institution:
address:
publisher:
accessed:
websites:
chapter:
series:
issue:
year: 2006
0: Jaramillo, Alvaro
title: Proposal (250) to South American Classification Committee
source: South American Classification Committee
type: miscellany
tags: Leucophaeus atricilla
abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
fullDescriptionUnstructured: La Gaviota Reidora mide aproximadamente 40 cm; en plumaje reproductivo, cabeza

enteramente negra con una estrecha mancha negra atrás del ojo. Manto gris oscuro, primarias exteriores enteramente negras sin manchas subterminales, cola blanca, pico rojizo y patas rojizas negruzcas. Pecho blanco. En plumaje básico cabeza blanca con manchas negruzcas desde nuca, auriculares y frente al ojo (Zamudio, 2012). Anteriormente *Larus atricilla* (Linnaeus, 1758). Cambió de acuerdo con la propuesta (250) del South American Classification Committee (Jaramillo, 2006).

briefDescription:

references

keywords: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chrysomus icterocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Coccyzus americanus*, *Coeligena bonapartei*, *Colibri coruscans*, *Columba livia*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cooperi*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dubusia taeniata*, *Egretta caerulea*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Glaucidium jardinii*, *Grallaria ruficapilla*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Leiothlypis peregrina*, *Lesbia nuna*, *Lesbia victoriae*, *Leucophaeus atricilla*

editors

editors: Stiles, F G
editors: Bohórquez, C I
editors: Cadena, C D
editors: De la Zerda, S
editors: Hernández, M
editors: Roselli, L
editors: Kelsey, M
editors: Valencia, I D
editors: Knapp, D
institution:
edition: Primera Ed
address:
publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca
accessed:
websites:
chapter:
series:
pages: 276
issue:
year: 2000

authors

authors: Asociación Bogotana de Ornitología
title: Aves de la Sabana de Bogotá: guía de campo
source:
type: book
tags: *Actitis macularius*, *Ampelion rubrocristatus*, *Anas discors*, *Anisognathus igniventris*, *Ardea alba*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bartramia longicauda*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Catharus fuscescens*, *Catharus ustulatus*, *Chaetocercus*

mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dubusia taeniata,Egretta caerulea,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreiventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Gallinula melanops,Geothlypis philadelphia,Glaucidium jardinii,Grallaria ruficapilla,Hemispingus superciliaris,Hemispingus verticalis,Henicorhina leucophrys,Hirundo rustica,Icterus chrysater,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Leiothlypis peregrina,Lesbia nuna,Lesbia victoriae,Leucophaeus atricilla

identifiers

last_modified:
 created:
 group_id:
 profile_id:
 id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
 providerLink:

publicationStatus

microReference:
 source:
 datatype:
 identifier:
 source:
 datatype:
 identifier:
 note:
 ruleConsidered:

replacementNameFor

microReference:
 source:
 datatype:
 identifier:
 source:
 datatype:
 identifier:
 note:
 ruleConsidered:

sanctioned

microReference:
 source:
 datatype:
 identifier:

source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:
specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Linnaeus, 1758)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Leucophaeus atricilla
rank: SPECIES
simple: Leucophaeus atricilla (Linnaeus, 1758)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

ancillaryData

synonymStatus:
synonymName
anyTwo:

anyOne

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:
datatype:
identifier:

[combinationAuthorship](#)

basionymAuthorship

simple:

authorship

simple:

simple: Bruch, 1855

epithet

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

genus

linkType:

ref:

uninomial:

simple: *Atricilla megalopterus*

rank: SPECIES

simple: *Atricilla megalopterus* Bruch, 1855

nomenclaturalCode:

isAnamorphic:

id:

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-02-21T23:03:59.175Z

created: 2016-02-21T23:03:59.175Z

[agent](#)

homepage:

role:

email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Leucophaeus
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Leucophaeus
family: Laridae
order: Charadriiformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:

0

link: <http://eol.org/pages/1049596/overview>

keywords: Leucophaeus atricilla

0: Frankis, Michael

1: Parr, Cyndy

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

0: EOL

title: Leucophaeus atricilla

source: Encyclopedia of life

type: miscellany

tags: Leucophaeus atricilla

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

reproductionUnstructured: La Gaviota Reidora se reproduce en grandes colonias; construye un gran nido principalmente de pastos, que coloca sobre el suelo; incuba durante tres semanas, entre tres y cuatro huevos verdosos (EOL, n.d.).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis

philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidnucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:
created:
mimeType:
dataType:
identifier:

feedingUnstructured: Esta especie se alimenta de carroña, peces pequeños e insectos, los cuales a veces captura en el aire (Asociación Bogotana de Ornitología, 2000).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:
structureDispersed:
type:
purpose:

behavior

0

dataObject:
additionalInformation:

0

keywords: Accipiter striatus,Actitis macularius,Adelomyia melanogenys,Aglaiocercus kingii,Amblycercus holosericeus,Ampelion rubrocristatus,Anas andium,Anas clypeata,Anas discors,Andigena nigrirostris,Anisognathus igniventris,Ardea alba,Ardea herodias,Arremon assimilis,Asio flammeus,Asio stygius,Asthenes flammulata,Asthenes fuliginosa,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Butorides virescens,Cairina moschata,Calidris bairdii,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Cathartes aura,Catharus minimus,Catharus ustulatus,Chaetocercus mulsant,Chaetura pelagica,Chalcostigma heteropogon,Chordeiles minor,Chrysomus icterocephalus,Cinclodes albidiventris,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Cistothorus platensis,Coccyzua pumila,Coccyzus americanus,Coccyzus melacoryphus,Coeligena bonapartei,Colibri coruscans,Colibri thalassinus,Columba livia,Columbina talpacoti,Conirostrum rufum,Conirostrum sitticolor,Contopus cinereus,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Dendrocygna autumnalis,Dendrocygna viduata,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dolichonyx oryzivorus,Dubusia taeniata,Egretta caerulea,Egretta thula,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreiventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Gallinula melanops,Geothlypis philadelphia,Geranoaetus melanoleucus,Glaucidium jardinii,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Heliangelus amethysticollis,Heliangelus exortis,Hellmayrea gularis,Hemispingus superciliosus,Hemispingus verticalis,Henricorhina leucophrys,Hirundo rustica,Icterus chrysater,Icterus icterus,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Lafresnaya lafresnayi,Leiothlypis peregrina,Leptasthenura andicola,Lesbia nuna,Lesbia victoriae,Leucophaeus atricilla,Machetornis rixosa,Margarornis squamiger,Mecocerculus leucophrys,Megascops choliba,Melanerpes rubricapillus,Merganetta armata,Metallura tyrianthina,Milvago chimachima,Mimus gilvus,Mniotilta varia,Molothrus bonariensis,Muscisaxicola alpinus,Myiarchus crinitus,Myiodynastes luteiventris,Myiotheretes fumigatus,Myiotheretes striaticollis,Myiothlypis nigrocristata,Nomonyx dominicus,Nycticorax nycticorax,Ochthoeca cinnamomeiventris,Ochthoeca diadema,Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus

versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Agelaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccyzus pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta caerulea, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnaya, Leiostyris alpestris, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla, Mactornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes luteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon guerinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon

pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus ludovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

abstract:

last_modified:

created:

group_id:

profile_id:

id:

1

keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus supercilialis, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:

websites:

chapter:

series:

pages: 276

issue:

year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source:

type: book

tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dubusia taeniata, Egretta caerulea, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreiventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Gallinula melanops, Geothlypis philadelphia, Glaucidium jardinii, Grallaria ruficapilla, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Leiothlypis peregrina, Lesbia nuna, Lesbia victoriae, Leucophaeus atricilla

abstract:

last_modified:

created:

group_id:

profile_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Es la gaviota más común en invierno boreal; a menudo en grandes bandadas; es una especie agresiva y oportunista en forrajeo (Hilty & Brown, 1986). Descansa sobre el agua lejos de la orilla (Asociación Bogotana de Ornitología, 2000).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

link: http://awsassets.panda.org/downloads/migratorias{_}aves{_}42{_}final.pdf
keywords: Leucophaeus atricilla

[editors](#)

editors: Naranjo, L G
editors: Amaya, J D
editors: Eusse-González, D
editors: Cifuentes-Sarmiento, Y
institution:
address:
publisher: Ministerio de Ambiente y Desarrollo Sostenible / WWF Colombia
accessed:
websites:
chapter:
series:
pages: 708
issue:
volume: 1
year: 2012

[authors](#)

authors: Zamudio, Jeisson
title: Leucophaeus atricilla
source: Guía de las Especies Migratorias de la Biodiversidad en Colombia Aves
type: book_section
tags: Leucophaeus atricilla
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:
distributionUnstructured: Esta especie se distribuye a lo largo de la Costa Atlántica de Norteamérica desde Maine central hasta el sur de Georgia, alrededor de las costas de Florida y a lo largo de la Costa del Golfo a sur de Texas. En México se encuentra desde las islas de los estados de Campeche y Yucatán hasta las costas del norte del golfo de California y sur de Colima. También en las Antillas, desde las islas Bahamas hacia el sur a través de las Antillas Menores, así como en Aruba, Bonaire y en islas costeras de Venezuela; migra a ambas costas de Centro y Sudamérica hasta Chile (Zamudio, 2012). En Colombia está presente en ambas costas; unos pocos registros en el interior: Valle del Cauca en Cartago y la Laguna de Sonso, Boyacá (Laguna de Tota) y Antioquia (Medellín), Cauca (Popayán), bocas de Aracataca y Buritaca (Magdalena) (Zamudio, 2012). Se distribuye desde el nivel del mar hasta los 3000 msnm (Hilty & Brown, 1986).

distributionAtomized

stateProvince: Córdoba

ancillaryData

locality:
municipality:
country: Colombia

temporalCoverage

endDate:
startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

0

link: <http://maps.iucnredlist.org/map.html?id=22694455>

keywords: Leucophaeus atricilla

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: BirdLife International

1: NatureServe

title: Bird Species Distribution Maps of the World. 2012. Larus atricilla. The IUCN Red List of Threatened Species. Version 2014.3

source:

type: miscellany
tags: Leucophaeus atricilla
abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
populationBiologyUnstructured: La Gaviota Reidora tiene un amplio rango de distribución; la tendencia de la población está incrementando, por lo cual se encuentra en categoría de Preocupación Menor (BirdLife International and NatureServe, 2014).

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:

measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

abstract: La Gaviota Reidora mide aproximadamente 40 cm; ambos sexos son similares. Los adultos (plumaje de invierno): cabeza, cuello, rabadilla, cola, región ventral y bordes posteriores de las alas blancos; los lados de la cabeza y la nuca moteados indistintamente con fusco; manto gris oscuro, las puntas de las alas negras. Pico y patas negruzcos. El plumaje de cría difiere de tener la cabeza completamente negra, el pico y las patas rojos oscuros. Los juveniles son similares a los adultos en plumaje invernal pero con las remeras fuscas, contrastando con el manto y con una banda subterminal negra en la cola (Asociación Bogotana de Ornitología, 2000; Dunn & Aldefer, 2006). Es una especie transeúnte y residente de invierno; común en ambas costas (Hilty & Brown, 1986); generalmente habita en áreas de agua abierta en lagunas o humedales, descansando sobre el agua lejos de la orilla o sobre postes (Asociación Bogotana de Ornitología, 2000). Se alimenta de carroña, peces pequeños e insectos, los cuales a veces captura en el aire (Asociación Bogotana de Ornitología, 2000). La Gaviota Reidora tiene un amplio rango de distribución; la tendencia de la población está incrementando, por lo cual se encuentra en categoría de Preocupación Menor (BirdLife International and NatureServe, 2014).

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomez@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:
mediaURL: <http://avibase.bsc-eoc.org/species.jsp?lang=ES&avibaseid=6CDC3C9C8B50EA71>
description:

subject

source: <http://avibase.bsc-eoc.org>

audience

bibliographicCitation:
rightsHolder:
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:
created:

agent

contentType:
dataType:
identifier:

_id
_bsonType: ObjectId
id: VÇÃ•ö lg#çA€