

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e0f3c383c45700544e3cfb

Fecha de creación: 2016-03-10T04:10:49.061Z

annualCycles

annualCycleUnstructured:

habitats

habitatUnstructured:

[commonNamesAtomized](#)

creation\_date: 2016-03-10T04:10:49.061Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: Mesoescuto y número densamente punteado; mandíbulas con un fuerte anillo basal; ojos hirsutos; cara con pequeñas manchas rojizas en la parte superior de los orificios antenales; tergo abdominal I con una banda preapical completa o parcial.

briefDescription:

[references](#)

keywords: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus

alfkeni, *Mischocyttarus alfkenii excrucians*, *Mischocyttarus alfkenii trinitatis*, *Mischocyttarus carbonarius carbonarius*, *Mischocyttarus flavicans*, *Mischocyttarus foveatus*, *Mischocyttarus injucundus injucundus*, *Mischocyttarus insolitus*, *Mischocyttarus labiatus*, *Mischocyttarus latissimus*, *Mischocyttarus metoecus*, *Mischocyttarus petiolatus*, *Mischocyttarus pseudomimeticus*, *Mischocyttarus punctatus*, *Mischocyttarus tomentosus*, *Mischocyttarus vaqueroi*, *Parachartergus amazonensis*, *Parachartergus colobopterus*, *Parachartergus fasciipennis*, *Parachartergus fraternus*, *Parachartergus fulgidipennis*, *Parachartergus griseus*, *Parachartergus richardsi*, *Parachartergus weyrauchi*, *Polybia bistrata*, *Polybia catillifex*, *Polybia eberhardae*, *Polybia emaciata*, *Polybia fastidiosuscula*, *Polybia jurinei*, *Polybia liliacea*, *Polybia micans*, *Polybia occidentalis*, *Polybia platycephala platycephala*, *Polybia quadricincta*, *Polybia raii*, *Polybia rejecta*, *Polybia scrobalis pronotalis*, *Polybia scrobalis scrobalis*, *Polybia sericea*, *Polybia striata*, *Polybia velutina*, *Protopolybia acutiscutis*, *Protopolybia amarella*, *Protopolybia chartergoides chartergoides*, *Protopolybia exigua binominata*, *Protopolybia exigua exigua*, *Protopolybia fuscatus*, *Protopolybia rugulosa*, *Protopolybia sedula*, *Pseudopolybia compressa*, *Pseudopolybia vespiceps*, *Synoeca septentrionalis*, *Synoeca surinama*, *Synoeca virginea*

## editors

institution:  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
pages: 585  
issue:  
year: 1978

## authors

authors: Richards, O. W.  
title: The social wasps of the Americas excluding the Vespinae  
source:  
type: book

tags: *Agelaia angulata angulata*, *Agelaia angulicollis*, *Agelaia brevistigma*, *Agelaia cajennensis*, *Agelaia fulvofasciata*, *Agelaia hamiltoni*, *Agelaia testacea*, *Angiopolybia obidensis*, *Angiopolybia pallens*, *Angiopolybia paraensis*, *Angiopolybia zischkai*, *Apoica thoracica*, *Asteloeca trilli*, *Brachygastra albula*, *Brachygastra augusti*, *Brachygastra lecheguana*, *Brachygastra propodealis*, *Brachygastra scutellaris*, *Chartergellus amazonicus*, *Chartergellus punctator*, *Chartergus chartarius*, *Chartergus globiventris*, *Chartergus metanotalis*, *Clypearia sulcata*, *Clypearia weyrauchi*, *Epipona tatua*, *Leipomeles dorsata*, *Metapolybia acincta*, *Metapolybia docilis*, *Metapolybia nigra*, *Metapolybia suffusa*, *Mischocyttarus adolphi*, *Mischocyttarus alfkeni*, *Mischocyttarus alfkenii excrucians*, *Mischocyttarus alfkenii trinitatis*, *Mischocyttarus carbonarius carbonarius*, *Mischocyttarus flavicans*, *Mischocyttarus foveatus*, *Mischocyttarus injucundus injucundus*, *Mischocyttarus insolitus*, *Mischocyttarus labiatus*, *Mischocyttarus latissimus*, *Mischocyttarus metoecus*, *Mischocyttarus petiolatus*, *Mischocyttarus pseudomimeticus*, *Mischocyttarus punctatus*, *Mischocyttarus tomentosus*, *Mischocyttarus vaqueroi*, *Parachartergus amazonensis*, *Parachartergus colobopterus*, *Parachartergus fasciipennis*, *Parachartergus fraternus*, *Parachartergus fulgidipennis*, *Parachartergus griseus*, *Parachartergus richardsi*, *Parachartergus weyrauchi*, *Polybia bistrata*, *Polybia catillifex*, *Polybia eberhardae*, *Polybia emaciata*, *Polybia fastidiosuscula*, *Polybia jurinei*, *Polybia liliacea*, *Polybia micans*, *Polybia occidentalis*, *Polybia platycephala platycephala*, *Polybia quadricincta*, *Polybia raii*, *Polybia rejecta*, *Polybia scrobalis pronotalis*, *Polybia scrobalis scrobalis*, *Polybia sericea*, *Polybia striata*, *Polybia velutina*, *Protopolybia acutiscutis*, *Protopolybia amarella*, *Protopolybia chartergoides chartergoides*, *Protopolybia exigua binominata*, *Protopolybia exigua exigua*, *Protopolybia fuscatus*, *Protopolybia rugulosa*, *Protopolybia sedula*, *Pseudopolybia compressa*, *Pseudopolybia vespiceps*, *Synoeca septentrionalis*, *Synoeca surinama*, *Synoeca virginea*

abstract:

## identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:

identifier:  
note:  
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:  
microReference:  
year:

publishedIn

simple:  
source:  
datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: Richards, 1978  
canonicalName  
infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: Chartergellus punctatior  
rank: SPECIES  
simple: Chartergellus punctatior Richards, 1978

attributes

nomenclaturalCode:  
isAnamorphic:  
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:  
additionalInformation:

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-10T04:10:29.059Z

created: 2016-03-10T04:10:29.059Z

[agent](#)

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataType:

identifier:

parentTaxon: Chartergellus

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Chartergellus

family: Vespidae

order: Hymenoptera

classHierarchy: Insecta

phylum: Arthropoda

kingdom: Animalia

recommended:

classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
reproductionUnstructured:

0

dataObject:  
additionalInformation:

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder:  
rights:  
license:  
modified:  
created:

agent

mimeType:  
dataType:  
identifier:  
relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedBy: TraitBank  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue: Sexual  
measurementType: Tipo de Reproducción  
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

feedingUnstructured:

0

0

strategy: Depredadores

type: Heterótrofos - Omnívoros

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:



additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
behaviorUnstructured: Son insectos depredadores con hábitos eusociales.

#### ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

#### measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

#### endemicAtomized

#### distribution

#### ancillaryData

dataObject:  
additionalInformation:

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

agent

mimeType:  
dataType:  
identifier:  
distributionUnstructured: Esta especie se distribuye en Colombia, Perú, Brasil, Guyana Francesa y Surinam, en zonas que alcanzan los 1600 msnm. En el país esta reportada para las regiones Pacífica, Orinoquía y Amazónica.

distributionAtomized

stateProvince: Meta

ancillaryData

locality:  
municipality:  
country: Colombia

temporalCoverage

endDate:  
startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured:  
areaOfOccupancy:  
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:  
legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Presenta una carena pronotal muy separada de la fovea; protuberancia frente de la fovea, aguda y baja; metanoto con superficie dorsal estrecha; espiráculo del propodeo estrecho, 4 veces más ancho que alto.

## associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

## ancillaryData

dataObject:  
additionalInformation: Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto “Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC”.  
Información de tipos: Holotipo capturado en Perú y depositado en el British Museum of Natural History, Londres, Inglaterra (BMNH).

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:  
rightsHolder:  
rights:  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

agent

contentType:  
dataType:  
identifier:

\_id  
\_bsonType: ObjectID  
id: VòóÄfÄW Dãï°