

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e2e02e83c45700544e3d85

Fecha de creación: 2016-03-11T15:11:48.569Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Habita bosques húmedos tropicales de Sudamérica.

[commonNamesAtomized](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:  
dataFormat:  
identifier:  
usedBy:

usedIn

distributionUnstructured:  
endDate:  
startDate:

## ancillaryData

type:  
language: Español  
name: Libélula  
creation\_date: 2016-03-11T15:11:48.569Z

usesManagementAndConservation

## ancillaryData

managementAndConservationUnstructured:

## managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
contentType:  
dataFormat:  
identifier:

fullDescriptionUnstructured: Base del ala del macho hialina o con una mancha blancuzca, venación alar basal negra. Patrón de coloración de ambos sexos similar, áreas cobrizas metálicas acompañadas de áreas pálidas limitadas a finas líneas en las suturas torácicas. Lóbulo anterior del protórax ligeramente curvo, sin lóbulo desarrollado en el margen anterior. Paraproctos del macho vestigiales. Lóbulo medio de los cercos del macho suavemente redondeado en un semicírculo. Tercio posterior de los cercos fuertemente arqueados.

briefDescription:

## references

keywords: Mnesarete fulgida,Mnesarete metallica

editors

institution:  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
pages: 2--84  
issue:  
volume: 506  
year: 2006

authors

authors: Garrison, R. W.  
title: A synopsis of the genera Mnesarete Cowley, Bryoplathanon gen. nov., and Ormenoplebia gen. nov. (Odonata: Calopterygidae)  
source: Contributions in Science (Natural History Museum of Los Angeles County)  
type: article  
tags: Mnesarete fulgida,Mnesarete metallica  
abstract:

identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
  
taxonRecordName

ancillaryData

providerSpecificData  
  
anyTwo:  
providerLink:  
  
publicationStatus  
  
microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
  
replacementNameFor  
  
microReference:  
source:  
datatype:  
identifier:

source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:

publishedIn

simple:  
source:  
datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: Selys, 1879

canonicalName

infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:

simple: Mnesarete fulgida  
rank: SPECIES  
simple: Mnesarete fulgida Selys, 1879

attributes

nomenclaturalCode:  
isAnamorphic:  
id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:  
additionalInformation:

[reference](#)

geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:  
rightsHolder:  
rights:  
license:  
title:  
modified: 2016-03-11T15:11:28.566Z  
created: 2016-03-11T15:11:28.566Z

[agent](#)

homepage:  
role:  
email:  
phone:  
address:  
position:  
organisation:  
lastName:  
firstName:  
mimeType:  
dataType:  
identifier:  
parentTaxon: Mnesarete  
higherClassification:

infraspecificEpithet:  
specificEpithet:  
taxonRank: SPECIES  
subGenus:  
genus: Mnesarete  
family: Calopterygidae  
order: Odonata  
classHierarchy: Insecta  
phylum: Arthropoda  
kingdom: Animalia  
recommended:  
classification:  
  
invasiveness  
  
invasivenessUnstructured:  
  
reproduction

0  
  
dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
reproductionUnstructured:

0  
  
dataObject:  
additionalInformation:

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder:

rights:  
license:  
modified:  
created:

## agent

contentType:  
dataType:  
identifier:  
relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedBy: TraitBank  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue: Sexual  
measurementType: Tipo de Reproducción  
measurementID:

## molecularData

molecularDataUnstructured:

## migratory

migratoryUnstructured:

## lifeForm

lifeFormUnstructured:

## lifeCycle

lifeCycleUnstructured:

## interactions

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
contentType:  
dataType:  
identifier:  
interactionsUnstructured: Se han encontrado en compañía de individuos de los géneros Protoneura, Perissolestes, Philogenia, Mecistogaster, y Heteragrion.



feeding

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
feedingUnstructured:

0

0

strategy: Depredadores  
type:

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:  
distance:  
structureDispersed:  
type:  
purpose:

behavior

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Vuelan lentamente y se congregan cerca de claros soleados del bosque a grandes alturas, alrededor de 2.6 m. o en zonas sombreadas cerca de quebradas. Estas quebradas en el bosque húmedo son claras pero tienen profundos fondos fangosos.

## ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

## measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

## endemicAtomized

## distribution

## ancillaryData

dataObject:

additionalInformation:

## reference

geoPoint:

location:

thumbnailURL:

## mediaURL

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:

dataType:

identifier:

distributionUnstructured: Se ha registrado en zonas que van desde los 90 msnm hasta los 400 msnm de altitud. En Colombia se distribuye en la región de la Amazonia.

[distributionAtomized](#)

[ancillaryData](#)

locality:

municipality:

country: Ecuador

temporalCoverage

endDate:

startDate:

distributionScope

type:

[threatStatus](#)

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

[ancillaryData](#)

dataObject:

additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

#### measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

abstract: Mnesarete fulgida es un insecto depredador que forraje en áreas sombreadas y habita en bosques húmedos tropicales de Sudamérica. El patrón de coloración de ambos sexos es similar, presentan áreas cobrizas metálicas acompañadas de áreas pálidas limitadas a finas líneas en las suturas torácicas. En el macho la base del alas es hialina o con una mancha blancuzca y la venación basal es negra.

#### associatedParty

role:  
personnelIdentifier:  
personnelDirectory:  
homepage:  
email: enataliagomezm@gmail.com  
phone:  
postalCode:  
country: Colombia  
state: Bogotá Distrito Capital  
city: Bogotá D.C.  
address:  
organisation:  
position:  
lastName: Gómez Melendro  
firstName: Edith Natalia

#### ancillaryData

dataObject:  
additionalInformation: Información de tipos: El lectotipo de esta especie está depositado en el British Museum of Natural History (BMNH), Londres, Inglaterra.  
Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaría General de la CAN y el Instituto Sinchi para el

proyecto “Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC”.

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:  
rightsHolder:  
rights:  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

agent

mimeType:  
dataType:  
identifier:

\_id  
\_bsontype: ObjectID  
id: Vââ.fÅW DãØP