

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e4eab183c45700544e3e63

Fecha de creación: 2016-03-13T04:21:12.552Z

creation_date: 2016-03-13T04:21:12.552Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataFormat:

identifier:

fullDescriptionUnstructured: El cuerpo del macho es gris azulado oscuro. Las alas tienen una mancha negra en el ápice. La hembra es similar al macho pero tiene el abdomen negro. Machos y hembras inmaduras tienen el cuerpo café claro y líneas transversales café oscuro en el dorso del tórax, y el abdomen presenta bandas intercaladas café claro y café oscuro. Tamaño del macho 2.9 a 3.6 cm, y de la hembra 3.3 a 3.6 cm.

briefDescription:

[references](#)

keywords: Argia cupraurea, Argia pulla Hagen in, Cora chiribiquete, Cora inca, Erythemis vesiculosa, Erythrodiplax kimminsi, Erythrodiplax umbrata, Erythrodiplax unimaculata, Mecistogaster linearis, Orthemis discolor, Perissolestes magdalenae, Perithemis lais, Perithemis mooma, Polythore beata, Polythore concinna, Polythore derivata, Polythore mutata, Telebasis corallina, Uracis fastigiata, Uracis imbuta

[editors](#)

institution:

address:

publisher: Instituto Nacional de Biodiversidad, Ministerio del Ambiente y Energía, Universidad Nacional de Costa Rica, Banco Mundial, GEF

accessed:

websites:

chapter:

series:

issue:

year: 2006

authors

authors: Esquivel, C
title: Libélulas de Mesoamérica y el Caribe. Dragonflies and damselflies of Middle America and the Caribbean
source:
type: book
tags: Argia cupraurea,Argia pulla Hagen in,Cora chiribiquete,Cora inca,Erythemis vesiculosa,Erythrodiplax kimminsi,Erythrodiplax umbrata,Erythrodiplax unimaculata,Mecistogaster linearis,Orthemis discolor,Perissolestes magdalenae,Perithemis lais,Perithemis mooma,Polythore beata,Polythore concinna,Polythore derivata,Polythore mutata,Telebasis corallina,Uracis fastigiata,Uracis imbuta
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:

datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:

identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:

microReference:

year:

publishedIn

simple:

source:

datatype:

identifier:

specialAuthorship

[authors](#)

[year](#)

simple:

canonicalAuthorship

[authors](#)

[year](#)

simple:

simple: Burmeister, 1839

canonicalName

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

linkType:

ref:

uninomial:

simple: Uracis imbuta

rank: SPECIES

simple: Uracis imbuta Burmeister, 1839

attributes

nomenclaturalCode:

isAnamorphic:

id:

[synonymsAtomized](#)

ancillaryData

synonymStatus:
synonymName
anyTwo:

anyOne

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:
datatype:
identifier:

[combinationAuthorship](#)

basionymAuthorship

simple:

authorship

simple:

simple: Rambur, 1842

epithet

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

genus

linkType:

ref:

uninomial:

simple: Uracis quadra

rank: SPECIES

simple: Uracis quadra Rambur, 1842

nomenclaturalCode:

isAnamorphic:

id:

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-13T04:20:52.549Z

created: 2016-03-13T04:20:52.549Z

[agent](#)

homepage:

role:

email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Uracis
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Uracis
family: Libellulidae
order: Odonata
classHierarchy: Insecta
phylum: Arthropoda
kingdom: Animalia
recommended:
classification:

commonNameAtomized

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

mimeType:
dataType:

identifier:

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

ancillaryData

type:

language: Español

name: Libélula

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

reproductionUnstructured: Los hábitos reproductivos de esta especie han sido poco estudiados. Se han encontrado hembras desovando sobre el barro húmedo de un sendero de bosque lluvioso, permaneciendo en vuelo suspendido cerca del suelo y cada cierto tiempo bajaba repentinamente el abdomen hasta tocar el suelo con la punta de éste, aparentemente poniendo los huevos en el barro.

0

dataObject:

additionalInformation:

reference

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder:
rights:
license:
modified:
created:

agent

mimeType:
dataType:
identifier:
relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedBy: TraitBank
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue: Sexual
measurementType: Tipo de Reproducción
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

contentType:
dataType:
identifier:
feedingUnstructured:

0

0

strategy: Depredadores
type:

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

behaviorUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

annualCycle

annualCycleUnstructured:

habitat

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

habitatUnstructured: Habita lagunas con vegetación emergente en el bosque.

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:
dataType:
identifier:
distributionUnstructured: Se distribuye en Belice, El Salvador, México, Guatemala, Honduras, Nicaragua, Costa Rica, Panamá, Colombia, Ecuador, Perú, Bolivia, Venezuela, Trinidad y Tobago, Guyana, Guyana Francesa, Surinám, Brasil y Argentina. En el país esta especie se ha registrado en las regiones de la Amazonia, Caribe, Pacífica, Andina y Orinoquía.

distributionAtomized

stateProvince: Amazonas

ancillaryData

locality:
municipality:
country: Colombia

temporalCoverage

endDate:
startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured:
areaOfOccupancy:
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:

modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

abstract: Uracis imbuta es una especie depredadora que habita en bosques de Sudamérica y Centroamérica. Los machos son gris azulado oscuro y las alas tienen una mancha negra en el ápice. La hembra es similar al macho pero tiene el abdomen negro.

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:
mediaURL: <https://www.flickr.com/photos/dmills727/12315026643/in/photolist-fQAEBg-9jWFhd-jLguzw-jLeJnz-jLeUNz-9jWFc7-jLgkSS-bwhAWk-boiPAi-bnAuXJ-bnAuWy>
description:

subject

source: https://farm4.staticflickr.com/3790/12315026643_e06687e528.jpg

audience

bibliographicCitation:
rightsHolder: dmills727
rights:
license: <http://creativecommons.org/licenses/by-nc-nd/2.0/>
modified:
created:

agent

mimeType:
dataType:
identifier:

_id
_bsontype: ObjectID
id: Vãê±f ÅW Dãæ0