Reporte de Fichas para el usuario: edith gómez
Ficha ID: 56e0b3e283c45700544e3ce4 Fecha de creación: 2016-03-09T23:38:15.888Z
annualCycles
annualCycleUnstructured:
habitats
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: habitatUnstructured: Esta especie habita en áreas perturbadas de tierras bajas.
commonNamesAtomized
ancillaryData
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
source:
audience
bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
agent

mime i ype:	
dataType:	
identifier:	
usedBy:	
usedIn	
distributionUns	structured:

endDate: startDate:

## ancillaryData

type:

language: Español name: Ciega

creation\_date: 2016-03-09T23:38:15.888Z usesManagementAndConservation

## ancillaryData

managementAndConservationUnstructured:

# managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

keywords: Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica

institution: Universidad de Costa Rica

publisher: accessed: websites: chapter: series: issue: year: 2010 0: Valverde, J.P.

address:

title: Clave taxonómica ilustrada para los géneros y las especies de avispas eusociales de Costa Rica

(Hymenoptera: Vespidae, Polistinae)

source: type: thesis

tags: Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica abstract:

last\_modified:

created	d:
group_	_id:
profile	id:

id:

geoPoint: location:

thumbnailURL: description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: Ocelos agrandados, separados de los ojos compuestos por menos de un diámetro ocelar; ala posterior con el lóbulo yugal reducido (Valverde, 2010). Tergo abdominal II marrón; ápice del tergo abdominal I y base del II sin manchas amarillas; metanoto y propodeo con marcas pálidas; metanoto con un tubérculo central suave.

briefDescription:

### references

keywords: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus.Mischocyttarus vaqueroi,Parachartergus amazonensis,Parachartergus colobopterus,Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea

### editors

institution: address: publisher: accessed: websites: chapter: series: pages: 585 issue: year: 1978

### authors

authors: Richards, O. W.

title: The social wasps of the Americas excluding the Vespinae

source: type: book

tags: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae.Mischocyttarus insolitus.Mischocyttarus labiatus.Mischocyttarus latissimus.Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi,Parachartergus weyrauchi,Polybia bistriata,Polybia catillifex,Polybia eberhardae,Polybia emaciata,Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea abstract:

## identifiers

last\_modified: created: group\_id: profile\_id: id:

taxonRecordName

## ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype: identifier: note:

rule(	Considered:
repla	acementNameFor
sour data iden sour data iden note	type: tifier: ce: type: tifier:
sand	ctioned
sour data iden sour data iden note	type: tifier: ce: type: tifier:
later	HomonymOf
sour data iden sour data iden note	type: tifier: ce: type: tifier:
con	servedAgainst
base	edOn
sour data ident sour data ident note	type: tifier: ce: type: tifier:
basi	onym
sour data	type: tifier:

datatype: identifier: note: ruleConsidered:
spellingCorrentionOf
typeNameEntity
source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:
typificacion
typeOfType:
lectotypeMicroReferenceVoucher
lectotypePublicationVoucher
voucherReference
simple: microReference: year:
publishedIn
simple: source: datatype: identifier:
specialAuthorship
authors
year
simple:
canonicalAuthorship
authors
year
simple: simple: Buysson, 1906

canonicalName

infraspecificEpithet: specificEpithet: infragenericEpithet: linkType: ref: uninomial: simple: Apoica thoracica rank: SPECIES simple: Apoica thoracica Buysson, 1906 attributes nomenclaturalCode: isAnamorphic: synonymsAtomized hierarchy ancillaryData dataObject: additionalInformation: reference geoPoint: location: thumbnailURL: mediaURL description: subject source: The Catalogue of Life, 3rd January 2011 audience bibliographicCitation: rigthsHolder: rights: license: title: modified: 2016-03-09T23:37:55.880Z created: 2016-03-09T23:37:55.880Z agent homepage: role: email:

role:
email:
phone:
address:
position:
organisation:
lastName:

firstName: mimeType: dataType: identifier: parentTaxon

parentTaxon: Apoica higherClassification: infraspecificEpithet: specificEpithet:

taxonRank: SPECIES

subGenus: genus: Apoica family: Vespidae order: Hymenoptera classHierarchy: Insecta phylum: Arthropoda kingdom: Animalia recommended: classification:

invasiveness

invasivenessUnstructured:

reproduction

Λ

dataObject:

additionalInformation:

0

keywords: Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica

institution: Universidad de Costa Rica

address: publisher: accessed: websites: chapter: series: issue: year: 2010

0: Valverde, J.P.

title: Clave taxonómica ilustrada para los géneros y las especies de avispas eusociales de Costa Rica

(Hymenoptera: Vespidae, Polistinae)

source: type: thesis

tags: Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica

abstract: last\_modified:

created: group\_id: profile\_id: id:

geoPoint:

location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Equipo Coordinador del SiB Colombia license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured: Sus nidos son panales planos con forma de disco, sésil en una rama y sin envoltura (Valverde, 2010).
0
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
audience
rightsHolder: rights: license: modified: created:
agent
mimeType: dataType: identifier: relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy: TraitBank measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: Sexual measurementType: Tipo de Reproducción measurementID:

molecularData

molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions
interactionsUnstructured:
feeding
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: feedingUnstructured:
0
0
strategy: Depredadores type: Heterótrofos - Omnívoros
environmentalEnvelope
environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured: distance: structureDispersed: type: purpose:

# behavior dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: behaviorUnstructured: Son insectos eusociales con habitos predadores. Activas principalmente en la noche, durante el día permanecen sobre el nido y lo defienden activamente; los machos parecen volar junto con las hembras en los enjambres. ancillaryData dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier: measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

## endemicAtomized

### distribution

# ancillaryData

dataObject:

additionalInformation:

#### reference

geoPoint: location:

thumbnailURL:

## mediaURL

description:

# subject

source:

## audience

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

## agent

mimeType: dataType: identifier:

distributionUnstructured: Se distribuye en Centroamérica y Sudamérica (Guatemala, Honduras, Costa Rica, Panamá, Colombia, Venezuela, Ecuador, Perú, Brasil y Bolivia) en áreas con altitudes desde los 0 hasta los 1594 msnm. En el país se extiende ampliamente a excepción de la Costa Atlántica y el norte del Chocó.

### distributionAtomized

stateProvince: Cauca

# ancillaryData

locality: municipality: country: Colombia

temporalCoverage

endDate: startDate:

distributionScope

type:

### threatStatus

territory

territoryUnstructured: areaOfOccupancy: extentOfOccurrence:
populationBiology
populationBiologyUnstructured:
legislation
legislationUnstructured:
directThreats
directThreatsUnstructured:
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementType: measurementID: abstract: Son avispas con los ocelos ensanchados y el mesepisterno dividido por una sutura en forma de T o Y invertida.
associatedParty
role: personnelIdentifier: personnelDirectory: homepage: email: enataliagomezm@gmail.com phone: postalCode: country: Colombia state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

# ancillaryData

### dataObject:

additionalInformation: Información de tipos: El lectotipo proviene de Espirito Santo, Brasil, y se encuentra en el Universitetes Zoologiske Museum, Copenhague, Dinamarca (UZMC).

Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

### reference

geoPoint: location: thumbnailURL:

### mediaURL

description:

## subject

source:

### audience

bibliographicCitation:

rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

# agent

mimeType: dataType: identifier:

\_id

\_bsontype: ObjectID id: Và³âfÄW DãÎ@