Reporte de Fichas para el usuario: edith gómez
Ficha ID: 56e20d7783c45700544e3d5d Fecha de creación: 2016-03-11T00:12:46.271Z
annualCycles
annualCycleUnstructured:
habitats
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: habitatUnstructured: Estos insectos habitan en bosques húmedos tropicales.
commonNamesAtomized
ancillaryData
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
source:
audience
bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
agent

mimeType:
dataType:
identifier:
usedBy:
usedIn

distributionUnstructured:

endDate: startDate:

ancillaryData

type:

language: Español name: Avispa

creation_date: 2016-03-11T00:12:46.271Z usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint: location: thumbnailURL: description: source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: Cuerpo amarillo brillante. Frente y mesoscutum con esculturación finamente granulada, la mayor parte del dorso del clípeo débilmente reticulada. Quilla pronotal muy débil. Tergito gastral 1 relativamente corto, pero fuertemente expandido posteriormente.

briefDescription:

references

keywords: Agelaia angulata angulata,Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Asteloeca traili,Brachygastra albula,Brachygastra

augusti,Brachygastra lecheguana,Brachygastra propodealis,Brachygastra scutellaris,Brachymenes wagnerianus, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona guerini, Epipona tatua, Eumenes versicolor filiformis, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus iniucundus tingomariae. Mischocyttarus insolitus. Mischocyttarus labiatus. Mischocyttarus latissimus. Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Omicron aequale nigritum, Omicron conclamatum, Omicron criticum pulchripictum, Omicron foxi, Omicron globicolle, Omicron graculum atripes, Omicron lustratum, Omicron nanum incarum, Omicron paranymphus, Omicron ypsilon, Pachymenes ghilianii olivaceus, Pachymenes orellanae orellanae, Pachymenes orellanae vardyi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi,Parachartergus smithii,Parachartergus weyrauchi,Pararhaphidoglossa clypeolaris,Pararhaphidoglossa confluenta confluenta. Pararhaphidoglossa cressoniana, Pararhaphidoglossa proxima, Polybia bistriata, Polybia catillifex,Polybia eberhardae,Polybia emaciata,Polybia fastidiosuscula,Polybia jurinei,Polybia liliacea,Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Santamenes peregrinus, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea, Synoecoides depressus, Zethus brasiliensis fuscatus, Zethus mexicanus lugubris, Zethus mexicanus mexicanus

editors

institution: Universidad Nacional de Colombia

address:
publisher:
accessed:
websites:
chapter:
series:
issue:
year: 1997

authors

authors: Sarmiento, C. E.

title: Véspidos de Colombia (Hymenoptera: Vespidae)

source: type: thesis

tags: Agelaia angulata angulata,Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia paraensis,Angiopolybia zischkai,Apoica thoracica,Asteloeca traili,Brachygastra albula,Brachygastra augusti,Brachygastra lecheguana,Brachygastra propodealis,Brachygastra scutellaris,Brachymenes wagnerianus,Chartergellus amazonicus,Chartergellus punctatior,Chartergus chartarius,Chartergus globiventris,Chartergus metanotalis,Clypearia sulcata,Clypearia weyrauchi,Epipona guerini,Epipona tatua,Eumenes versicolor filiformis,Leipomeles dorsata,Metapolybia acincta,Metapolybia docilis,Metapolybia nigra,Metapolybia suffusa,Mischocyttarus adolphi,Mischocyttarus alfkeni,Mischocyttarus alfkenii excrucians,Mischocyttarus alfkenii trinitatis,Mischocyttarus carbonarius carbonarius,Mischocyttarus flavicans,Mischocyttarus foveatus,Mischocyttarus injucundus tingomariae,Mischocyttarus insolitus,Mischocyttarus labiatus,Mischocyttarus latissimus,Mischocyttarus metoecus,Mischocyttarus petiolatus,Mischocyttarus pseudomimeticus,Mischocyttarus punctatus,Mischocyttarus tomentosus,Mischocyttarus vaqueroi,Omicron aequale nigritum,Omicron conclamatum,Omicron criticum pulchripictum,Omicron foxi,Omicron globicolle,Omicron graculum atripes,Omicron lustratum,Omicron nanum incarum,Omicron paranymphus,Omicron ypsilon,Pachymenes ghilianii olivaceus,Pachymenes orellanae orellanae vardyi,Parachartergus amazonensis,Parachartergus colobopterus,Parachartergus

fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus smithii, Parachartergus weyrauchi, Pararhaphidoglossa clypeolaris, Pararhaphidoglossa confluenta confluenta, Pararhaphidoglossa cressoniana, Pararhaphidoglossa proxima, Polybia bistriata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raui raui, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina. Protopolybia acutiscutis. Protopolybia amarella. Protopolybia chartergoides chartergoides. Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Santamenes peregrinus, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea, Synoecoides depressus, Zethus brasiliensis fuscatus, Zethus mexicanus lugubris, Zethus mexicanus mexicanus abstract:

identifiers

last modified: created: group id: profile id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:

replacementNameFor

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

sanctioned

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

laterHomonymOf

microReference:

source:
datatype:
identifier:
source:
datatype:
identifier:
note:

ruleConsidered:

conservedAgainst

basedOn

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

basionym

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

spelling Corrention Of

typeNameEntity

source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:

typificacion typeOfType: lectotypeM

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:

microReference:

year:

publishedIn

simple: source: datatype: identifier:

specialAuthorship

authors

year

simple:

canonicalAuthorship

authors

year

simple:

simple: Zikan, 1949

canonicalName

infraspecificEpithet: specificEpithet: infragenericEpithet:

linkType: ref: uninomial:

simple: Mischocyttarus insolitus

rank: SPECIES

simple: Mischocyttarus insolitus Zikan, 1949

attributes

nomenclaturalCode: isAnamorphic:

id.

synonymsAtomized

hierarchy

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location:

thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rigthsHolder: rights:

license:

title:

modified: 2016-03-11T00:12:26.269Z created: 2016-03-11T00:12:26.269Z

agent

homepage:

role: email: phone: address: position: organisation: lastName: firstName: mimeType: dataType: identifier:

parentTaxon: Mischocyttarus

higherClassification: infraspecificEpithet: specificEpithet: taxonRank: SPECIES

subGenus:

genus: Mischocyttarus family: Vespidae order: Hymenoptera classHierarchy: Insecta phylum: Arthropoda kingdom: Animalia recommended:

classification:
invasiveness
invasivenessUnstructured:
reproduction
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured: El nido de esta especie es muy extraño ya que consiste en una serie de grupos de 1 a 4 celdas independientes, dispuestas paralelamente a la hoja gracias al ángulo recto que forma el pedicelo con cada celda.
0
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
audience
rightsHolder: rights: license: modified: created:
agent
mimeType: dataType: identifier:

relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy: TraitBank measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: Sexual measurementType: Tipo de Reproducción measurementID:
molecularData
molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions
interactionsUnstructured:
feeding
0
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: feedingUnstructured: 0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: feedingUnstructured:
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: feedingUnstructured: 0

environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured: distance: structureDispersed: type: purpose:
behavior
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Nancy Carolina Rojas Riaño license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: behaviorUnstructured: Son insectos sociales con hábitos predadores.
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks:

measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy:

measurementValue: measurementType: measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

agent

mimeType: dataType: identifier:

distributionUnstructured: Se conocen poblaciones de este insecto en Brasil y Colombia. En el país se registra en la región de la Amazonia.

distributionAtomized

stateProvince: Amazonas

ancillaryData

locality: municipality: country: Colombia

temporalCoverage

endDate: startDate:
distributionScope
type:
threatStatus
territory
territoryUnstructured: areaOfOccupancy: extentOfOccurrence:
populationBiology
populationBiologyUnstructured:
legislation
legislationUnstructured:
directThreats
directThreatsUnstructured:
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID: abstract: Mischocyttarus insolitus habita bosques húmedos tropicales de Brasil y Colombia. Es una especie depredadora con comportamiento eusocial y presenta una coloración amarillo brillante en su cuerpo. Sus nidos
consisten en una serie de grupos de 1 a 4 celdas independientes dispuestas paralelamente a la hoja.

associatedParty

role:

personnelldentifier: personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone: postalCode: country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation: Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

Información de tipos: El holotipo se encuentra en el Departamento de Entomología de Instituto Oswaldo Cruz (FIOC), Sao Paulo, Brasil.

reference

geoPoint: location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

agent

mimeType: dataType: identifier:

_id

_bsontype: ObjectID id: VâĐ wĀW DãÕĐ