Reporte de Fichas para el usuario: edith gómez Ficha ID: 56c4e738f0106c67230e738e Fecha de creación: undefined annualCycles annualCycleUnstructured: habitats 0 dataObject: additionalInformation: keywords: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus 0: Stiles, F G 1: Bohórquez, C I 2: Cadena, C D 3: De la Zerda, S 4: Hernández, M 5: Roselli, L 6: Kelsey, M 7: Valencia, I D 8: Knapp, D institution: edition: Primera Ed address: publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca accessed: websites: chapter: series: pages: 276 issue: year: 2000 0: Asociación Bogotana de Ornitología title: Aves de la Sabana de Bogotá: guía de campo source: type: book tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus

mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Coccyzus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus

fumigatus abstract: last_modified: created: group_id: profile_id:

isbn: 978 987 9132272

keywords: Bartramia longicauda, Buteo swainsoni, Calidris melanotos, Catamenia analis, Coccyzus melacoryphus, Columbina talpacoti, Contopus cinereus, Contopus fumigatus, Crotophaga ani, Crotophaga major, Dendrocygna viduata, Egretta caerulea, Hirundo rustica, Icterus icterus, Jacana jacana, Machetornis rixosa, Milvago chimachima, Nycticorax nycticorax, Pandion haliaetus, Pardirallus maculatus, Patagioenas fasciata, Phaetusa simplex, Pheucticus aureoventris, Phimosus infuscatus, Pipraeidea melanonota, Progne tapera, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pygochelidon cyanoleuca, Sayornis nigricans 0: Editores, Vásques Mazzini

institution: address: publisher: accessed: websites: chapter: series: pages: 432 issue: year: 2010

title: Aves de Argentina y Uruguay-Birds of Argentina and Uruguay: guía de identificación edición total- a field guide

total edition source: type: book

0: Narosky, T 1: Yzurieta, D

tags: Bartramia longicauda, Buteo swainsoni, Calidris melanotos, Catamenia analis, Coccyzus melacoryphus, Columbina talpacoti, Contopus cinereus, Contopus fumigatus, Crotophaga ani, Crotophaga major, Dendrocygna viduata, Egretta caerulea, Hirundo rustica, Icterus icterus, Jacana jacana, Machetornis rixosa, Milvago chimachima, Nycticorax nycticorax, Pandion haliaetus, Pardirallus maculatus, Patagioenas fasciata, Phaetusa simplex, Pheucticus aureoventris, Phimosus infuscatus, Pipraeidea melanonota, Progne tapera. Pseudocolopteryx acutipennis. Pseudoscops clamator, Pygochelidon cyanoleuca. Sayornis nigricans abstract: Narosky, T., & Yzurieta, D. (2010). Aves de Argentina y Uruguay-Birds of Argentina and Uruguay: guía de identificación edición total- a field guide total edition. (V. M. Editores, Ed.) (p. 432). Buenos Aires, Argentina.

last modified: created: group id: profile id:

geoPoint: location:

thumbnailURL: description: source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

habitatUnstructured: El Pibí Ahumado habita en árboles secos (Narosky & Yzurieta, 2010); bordes de bosques primarios o secundarios, bosques abiertos de alisos o Eucaliptos; casi siempre se observa perchado en ramas más o menos altas y expuestas (Asociación Bogotana de Ornitología, 2000).

commonNamesAtomized

ancillaryData

usedBy:

usedIn

distributionUnstructured:

endDate: startDate:

ancillaryData

type:

language: Alemán name: Schiefertyrann

usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus fumigatus

- 0: Stiles, F G
- 1: Bohórquez, C I
- 2: Cadena, C D
- 3: De la Zerda, S
- 4: Hernández, M
- 5: Roselli, L
- 6: Kelsey, M
- 7: Valencia, I D
- 8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed: websites: chapter: series: pages: 276 issue: year: 2000

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source: type: book

tags: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus fumigatus

abstract:

last_modified:

created:

group_id:

profile id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: El Pibí Ahumado mide aproximadamente 17 cm; es un atrapamosca distintivo por su coloración oscura, cresta corta y la costumbre de pararse con postura erecta en las copas de los árboles. Los adultos son por encima gris oscuro, más oscuro en el copete; dos barras alares poco llamativas de gris más claro, por debajo gris un poco más pálido. Maxila negra, mandíbula anaranjada o amarillenta. Los juveniles son más cafecino en general con barras alares angostas de color ocre (Asociación Bogotana de Ornitología, 2000). briefDescription:

references

link: http://maps.iucnredlist.org/map.html?id=22699804

keywords: Contopus fumigatus

editors

institution: address:

publisher: accessed: websites: chapter: series: issue: year: 2014

authors

authors: BirdLife International

authors: NatureServe

title: Bird Species Distribution Maps of the World. 2012. Contopus fumigatus. The IUCN Red List of Threatened

Species. Version 2014.3

source:

type: miscellany

tags: Contopus fumigatus

abstract:

identifiers

last_modified: created: group_id: profile_id: id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

replacementNameFor

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

sanctioned

microReference:

source:

datatype: identifier:

source:

datatype: identifier:

note: ruleConsidered:

laterHomonymOf

•

microReference: source: datatype: identifier: source:

datatype: identifier:

note:

ruleConsidered:

conservedAgainst

basedOn

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

basionym

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

spellingCorrentionOf

typeNameEntity

source: datatype: identifier: source:

datatype: identifier:
source:
datatype: identifier:
typificacion
typeOfType:
lectotypeMicroReferenceVoucher
lectotypePublicationVoucher
voucherReference
simple: microReference: year:
publishedIn
simple: source: datatype: identifier:
specialAuthorship
authors
year
simple:
canonicalAuthorship
authors
year
simple: (Orbigny & Lafresnaye, 1837)
canonicalName
infraspecificEpithet: specificEpithet: infragenericEpithet: linkType: ref: uninomial: simple: Contopus fumigatus rank: SPECIES simple: Contopus fumigatus (Orbigny & Lafresnaye, 1837)
attributes
nomenclaturalCode:

isAnamorphic:

id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location: thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rigthsHolder:

rights: license: title:

modified: 2016-02-17T21:28:42.548Z created: 2016-02-17T21:28:42.548Z

agent

homepage:

role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:

parentTaxon: Contopus higherClassification: infraspecificEpithet: specificEpithet:

taxonRank: SPECIES

subGenus: genus: Contopus family: Tyrannidae order: Passeriformes classHierarchy: Aves phylum: Chordata kingdom: Animalia recommended: classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject: additionalInformation:

0

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis.Diglossa lafresnavii.Diglossa sittoides.Dolichonyx oryzivorus.Dubusia taeniata.Egretta thula.Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis.Muscisaxicola alpinus.Mviarchus crinitus.Mviodvnastes luteiventris.Mviotheretes fumigatus.Mviotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax, nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon querinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto

alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis

institution: address:

publisher: Princeton University Press

accessed: websites: chapter: series: pages: 836 issue: year: 1986 0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans.Colibri thalassinus.Columba livia.Columbina talpacoti.Conirostrum rufum.Conirostrum sitticolor.Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis. Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba. Vanellus chilensis. Vireo olivaceus. Vultur gryphus. Zenaida auriculata. Zonotrichia capensis abstract:

last_modified:
created: group_id:
profile_id:
id:
geoPoint: location:
thumbnailURL:
description: source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created: mimeType:
dataType:
identifier:
reproductionUnstructured: Se reproduce en diferentes épocas del año, el nido es una taza construida con material vegetal (musgo, líquenes), soportada en una rama horizontal a menudo alta (Hilty & Brown, 1986).
molecularData
molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions
interactionsUnstructured:
feeding
0
dataObject: additionalInformation:
0
keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus

cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus. Glaucidium jardinii. Grallaria guitesis. Grallaria ruficapilla. Grallaria squamigera. Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon querinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis institution:

address:

accessed:

publisher: Princeton University Press

websites: chapter: series: pages: 836 issue: year: 1986 0: Hilty, Steven 1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa

humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis.Henicorhina leucophrys.Hirundo rustica.Icterus chrysater.Icterus icterus.licterus nigrogularis.Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis abstract:

last_modified: created: group id:

profile_id:

id:

keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus fumigatus

0: Stiles, F G

1: Bohórquez, C I

2: Cadena. C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed: websites: chapter: series: pages: 276 issue: year: 2000 0: Asociación Bogotana de Ornitología title: Aves de la Sabana de Bogotá: guía de campo source: type: book tags: Actitis macularius, Ampelion rubrocristatus, Anas discors, Anisognathus igniventris, Ardea alba, Arremon assimilis, Asio flammeus, Asio stygius, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bartramia longicauda, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Catharus fuscescens, Catharus ustulatus, Chaetocercus mulsant, Chrysomus icterocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Occozyus americanus, Coeligena bonapartei, Colibri coruscans, Columba livia, Conirostrum rufum, Conirostrum sitticolor, Contopus cooperi, Contopus fumigatus abstract: last_modified: created: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rights Holder: rights: Asociación Bogotana de Ornitología license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created:
mimeType: dataType:
identifier:
feedingUnstructured: Como las otras especies de atrapamoscas del género Contopus sp., la dieta del Pibí Ahumado se compone principalmente de insectos (Asociación Bogotana de Ornitología, 2000; Hilty & Brown, 1986).
environmentalEnvelope
environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:
dispersal
dispersalUnstructured: distance: structureDispersed: type: purpose:
behavior

0

dataObject: additionalInformation:

0

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster. Elaenia frantzii. Elanus leucurus. Empidonax alnorum. Empidonax traillii. Ensifera ensifera. Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis. Myiothlypis nigrocristata. Nomonyx dominicus. Nycticorax nycticorax. Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina, Oxypogon querinii, Oxyura jamaicensis, Pachyramphus versicolor, Pandion haliaetus, Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia. Setophaga ruticilla. Setophaga striata. Sicalis flaveola. Sicalis luteola. Spiza americana. Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis institution:

address:

publisher: Princeton University Press

accessed: websites: chapter: series: pages: 836 issue: year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris. Ardea alba. Ardea herodias. Arremon assimilis. Asio flammeus. Asio stygius. Asthenes flammulata. Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria quitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus.Merganetta armata.Metallura tyrianthina.Milyago chimachima.Mimus gilyus.Mniotilta yaria.Molothrus bonariensis. Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis. Myiothlypis nigrocristata. Nomonyx dominicus. Nycticorax nycticorax. Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex, Phalacrocorax brasilianus, Pheucticus aureoventris, Pheucticus Iudovicianus, Phimosus infuscatus, Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis abstract:

last_modified:

created: group id:

profile id:

id:

4

isbn: 978 987 9132272

keywords: Bartramia longicauda, Buteo swainsoni, Calidris melanotos, Catamenia analis, Coccyzus melacoryphus, Columbina talpacoti, Contopus cinereus, Contopus fumigatus, Crotophaga ani, Crotophaga major, Dendrocygna viduata, Egretta caerulea, Hirundo rustica, Icterus icterus, Jacana jacana, Machetornis

rixosa, Milvago chimachima, Nycticorax nycticorax, Pandion haliaetus, Pardirallus maculatus, Patagioenas fasciata, Phaetusa simplex, Pheucticus aureoventris, Phimosus infuscatus, Pipraeidea melanonota, Progne tapera, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pygochelidon cyanoleuca, Sayornis nigricans 0: Editores, Vásques Mazzini

institution: address: publisher: accessed: websites: chapter: series: pages: 432 issue: year: 2010 0: Narosky, T

1: Yzurieta. D

title: Aves de Argentina y Uruguay-Birds of Argentina and Uruguay: guía de identificación edición total- a field guide

total edition source: type: book

tags: Bartramia longicauda,Buteo swainsoni,Calidris melanotos,Catamenia analis,Coccyzus melacoryphus,Columbina talpacoti,Contopus cinereus,Contopus fumigatus,Crotophaga ani,Crotophaga major,Dendrocygna viduata,Egretta caerulea,Hirundo rustica,Icterus icterus,Jacana jacana,Machetornis rixosa,Milvago chimachima,Nycticorax nycticorax,Pandion haliaetus,Pardirallus maculatus,Patagioenas fasciata,Phaetusa simplex,Pheucticus aureoventris,Phimosus infuscatus,Pipraeidea melanonota,Progne tapera,Pseudocolopteryx acutipennis,Pseudoscops clamator,Pygochelidon cyanoleuca,Sayornis nigricans abstract: Narosky, T., \& Yzurieta, D. (2010). Aves de Argentina y Uruguay-Birds of Argentina and Uruguay: guía de identificación edición total- a field guide total edition. (V. M. Editores, Ed.) (p. 432). Buenos Aires, Argentina. last modified:

created: group_id: profile_id: id:

2

keywords: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus fumigatus

0: Stiles, F G

1: Bohórquez, C I

2: Cadena, C D

3: De la Zerda, S

4: Hernández, M

5: Roselli, L

6: Kelsey, M

7: Valencia, I D

8: Knapp, D

institution:

edition: Primera Ed

address:

publisher: Asociación Bogotana de Ornitología; Corporación Autónoma Regional de Cundinamarca

accessed:	
websites:	
chapter:	
series:	
pages: 276	
issue:	
year: 2000	

0: Asociación Bogotana de Ornitología

title: Aves de la Sabana de Bogotá: guía de campo

source: type: book

tags: Actitis macularius,Ampelion rubrocristatus,Anas discors,Anisognathus igniventris,Ardea alba,Arremon assimilis,Asio flammeus,Asio stygius,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bartramia longicauda,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Catharus fuscescens,Catharus ustulatus,Chaetocercus mulsant,Chrysomus icterocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Coccyzus americanus,Coeligena bonapartei,Colibri coruscans,Columba livia,Conirostrum rufum,Conirostrum sitticolor,Contopus cooperi,Contopus

fumigatus abstract: last modified:

created:

group_id: profile_id: id:

geoPoint: location:

thumbnailURL: description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

behaviorUnstructured: Es una especie confiada, pasiva (Narosky & Yzurieta, 2010) y solitaria excepto en temporada reproductiva (Hilty & Brown, 1986); se posa erecto y alerta en ramas secas, usualmente en alturas medias y raramente en el dosel; más raro aún en la copa de los árboles. Hace vuelos medios a largos y usualmente regresa a la misma percha (Hilty & Brown, 1986).

El reclamo usual es un pip pip pip pip brillante y vivaz, a menudo persistente desde la percha prominente. El canto al amanecer es un juiiirit o pirrpiirITT enfático (Hilty & Brown, 1986). También otros reclamos como un alto y penetrante uiiiiiu y un pit pit que puede repetir dos y tres veces seguidas (Asociación Bogotana de Ornitología, 2000; Narosky & Yzurieta, 2010).

ancillaryData

dataObject:

additionalInformation:

geoPoint: location:

thumbnailURL: description:

source:

bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

endemicAtomized

distribution

ancillaryData

dataObject: additionalInformation:

reference

keywords: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis, Catamenia analis, Catamenia inornata, Cathartes aura, Catharus minimus, Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis.Coccycua pumila.Coccyzus americanus.Coccyzus melacoryphus.Coeligena bonapartei.Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera, Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria guitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca

cinnamomeiventris,Ochthoeca diadema,Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus,Parkesia noveboracensis,Patagioenas fasciata,Petrochelidon pyrrhonota,Phaetusa simplex,Phalacrocorax brasilianus,Pheucticus aureoventris,Pheucticus ludovicianus,Phimosus infuscatus,Phrygilus unicolor,Phyllomyias uropygialis,Piaya cayana,Picoides fumigatus,Pipraeidea melanonota,Pipreola riefferii,Piranga olivacea,Piranga rubra,Pitangus sulphuratus,Pluvialis dominica,Podilymbus podiceps,Porphyrio martinicus,Porzana carolina,Progne tapera,Protonotaria citrea,Pseudocolopteryx acutipennis,Pseudoscops clamator,Pterophanes cyanopterus,Pygochelidon cyanoleuca,Pyrocephalus rubinus,Pyrrhomyias cinnamomeus,Quiscalus lugubris,Rallus semiplumbeus,Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea,Setophaga castanea,Setophaga cerulea,Setophaga petechia,Setophaga ruticilla,Setophaga striata,Sicalis flaveola,Sicalis luteola,Spiza americana,Sporagra spinescens,Sturnella magna,Synallaxis subpudica,Systellura longirostris,Tangara vitriolina,Thraupis episcopus,Thraupis palmarum,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon,Troglodytes solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto alba,Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaida auriculata,Zonotrichia capensis

editors

institution: address:

publisher: Princeton University Press

accessed: websites: chapter: series: pages: 836 issue: year: 1986

authors

authors: Hilty, Steven authors: Brown, William

title: A Guide to the Birds of Colombia

source: type: book

tags: Accipiter striatus, Actitis macularius, Adelomyia melanogenys, Aglaiocercus kingii, Amblycercus holosericeus, Ampelion rubrocristatus, Anas andium, Anas clypeata, Anas discors, Andigena nigrirostris, Anisognathus igniventris, Ardea alba, Ardea herodias, Arremon assimilis, Asio flammeus, Asio stygius, Asthenes flammulata, Asthenes fuliginosa, Astragalinus psaltria, Atlapetes pallidinucha, Atlapetes schistaceus, Bubulcus ibis, Buteo platypterus, Buteo swainsoni, Butorides striata, Butorides virescens, Cairina moschata, Calidris bairdii, Calidris melanotos, Cardellina canadensis.Catamenia analis.Catamenia inornata.Cathartes aura.Catharus minimus.Catharus ustulatus, Chaetocercus mulsant, Chaetura pelagica, Chalcostigma heteropogon, Chordeiles minor, Chrysomus icterocephalus, Cinclodes albidiventris, Cinclus leucocephalus, Cinnycerthia unirufa, Cistothorus apolinari, Cistothorus platensis, Coccycua pumila, Coccyzus americanus, Coccyzus melacoryphus, Coeligena bonapartei, Colibri coruscans, Colibri thalassinus, Columba livia, Columbina talpacoti, Conirostrum rufum, Conirostrum sitticolor, Contopus cinereus, Contopus fumigatus, Contopus sordidulus, Contopus virens, Coragyps atratus, Crotophaga ani, Crotophaga major, Dendrocygna autumnalis, Dendrocygna viduata, Diglossa albilatera, Diglossa cyanea, Diglossa humeralis, Diglossa lafresnayii, Diglossa sittoides, Dolichonyx oryzivorus, Dubusia taeniata, Egretta thula, Elaenia flavogaster, Elaenia frantzii, Elanus leucurus, Empidonax alnorum, Empidonax traillii, Ensifera ensifera. Eriocnemis cupreoventris, Eriocnemis vestita, Eupsittula pertinax, Falco columbarius, Falco peregrinus, Falco sparverius, Forpus conspicillatus, Fulica americana, Gallinago nobilis, Gallinula galeata, Geothlypis philadelphia, Geranoaetus melanoleucus, Glaucidium jardinii, Grallaria guitesis, Grallaria ruficapilla, Grallaria squamigera, Haplospiza rustica, Heliangelus amethysticollis, Heliangelus exortis, Hellmayrea gularis, Hemispingus superciliaris, Hemispingus verticalis, Henicorhina leucophrys, Hirundo rustica, Icterus chrysater, Icterus icterus, Icterus nigrogularis, Ixobrychus exilis, Jacana jacana, Lafresnaya lafresnayi, Leiothlypis peregrina, Leptasthenura andicola, Lesbia nuna, Lesbia victoriae, Machetornis rixosa, Margarornis squamiger, Mecocerculus leucophrys, Megascops choliba, Melanerpes

rubricapillus, Merganetta armata, Metallura tyrianthina, Milvago chimachima, Mimus gilvus, Mniotilta varia, Molothrus bonariensis, Muscisaxicola alpinus, Myiarchus crinitus, Myiodynastes Iuteiventris, Myiotheretes fumigatus, Myiotheretes striaticollis, Myiothlypis nigrocristata, Nomonyx dominicus, Nycticorax nycticorax, Ochthoeca cinnamomeiventris, Ochthoeca diadema, Ochthoeca fumicolor, Ochthoeca rufipectoralis, Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus, Parkesia noveboracensis, Patagioenas fasciata, Petrochelidon pyrrhonota, Phaetusa simplex.Phalacrocorax brasilianus.Pheucticus aureoventris.Pheucticus Iudovicianus.Phimosus infuscatus.Phrygilus unicolor, Phyllomyias uropygialis, Piaya cayana, Picoides fumigatus, Pipraeidea melanonota, Pipreola riefferii, Piranga olivacea, Piranga rubra, Pitangus sulphuratus, Pluvialis dominica, Podilymbus podiceps, Porphyrio martinicus, Porzana carolina, Progne tapera, Protonotaria citrea, Pseudocolopteryx acutipennis, Pseudoscops clamator, Pterophanes cyanopterus, Pygochelidon cyanoleuca, Pyrocephalus rubinus, Pyrrhomyias cinnamomeus, Quiscalus lugubris, Rallus semiplumbeus, Ramphocelus dimidiatus, Ramphomicron microrhynchum, Riparia riparia, Rupornis magnirostris, Sayornis nigricans, Serpophaga cinerea, Setophaga castanea, Setophaga cerulea, Setophaga petechia, Setophaga ruticilla, Setophaga striata, Sicalis flaveola, Sicalis luteola, Spiza americana, Sporagra spinescens, Sturnella magna, Synallaxis subpudica, Systellura longirostris, Tangara vitriolina, Thraupis episcopus, Thraupis palmarum, Tringa flavipes, Tringa melanoleuca, Tringa solitaria, Troglodytes aedon, Troglodytes solstitialis, Turdus fuscater, Turdus ignobilis, Tyrannus melancholicus, Tyrannus savana, Tyrannus tyrannus, Tyto alba, Vanellus chilensis, Vireo olivaceus, Vultur gryphus, Zenaida auriculata, Zonotrichia capensis abstract:

identifiers

last_modified: created:

group_id: profile_id:

id:

geoPoint: location: thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

agent

mimeType:

dataType:

identifier:

distributionUnstructured: El Pibí Ahumado se distribuye desde Costa Rica, occidente de Panamá, norte y sur de Venezuela por los Andes hasta el noroccidente de Argentina. En Colombia desde la Serranía de Perijá y los Andes (Hilty & Brown, 1986).

Se distribuye entre los 300 msnm a los 3000 msnm (principalmente entre los 1500 msnm), pero comúnmente hasta los 500 msnm en la vertiente Pacífica (Hilty & Brown, 1986).

distributionAtomized

stateProvince: Cauca

ancillaryData

locality: municipality: country: Colombia

temporalCoverage

endDate: startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality: municipality: county: stateProvince: country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks: measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit: measurementAccuracy: measurementType: IUCN measurementID:

territory

territoryUnstructured: areaOfOccupancy: extentOfOccurrence:

populationBiology

Λ

dataObject:

additionalInformation:

0

link: http://maps.iucnredlist.org/map.html?id=22699804 keywords: Contopus fumigatus
institution: address:
publisher:
accessed:
websites:
chapter:
series:
issue: year: 2014
0: BirdLife International
1: NatureServe
title: Bird Species Distribution Maps of the World. 2012. Contopus fumigatus. The IUCN Red List of Threatened
Species. Version 2014.3
source:
type: miscellany
tags: Contopus fumigatus
abstract:
last modified:
created:
group_id:
profile_id:
id:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Asociación Bogotana de Ornitología
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
populationBiologyUnstructured: El Pibí Ahumado, tiene un amplio rango de distribución, la tendencia de la población
es estable y se encuentra en categoría de amenaza de Preocupación Menor (BirdLife International and
NatureServe, 2014).
legislation
legislationUnstructured:
directThreats
directThreatsUnstructured:
ancillaryData
dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:

COL	irce:
500	ись.

bibliographicCitation:

rightsHolder:

rights: license: modified: created: mimeType:

dataType: identifier:

measurementOrFact

relatedTo:

measurementRemarks: measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: El Pibí Ahumado mide aproximadamente 17 cm; es un atrapamosca distintivo por su coloración oscura, cresta corta y la costumbre de pararse con postura erecta en las copas de los árboles (Asociación Bogotana de Ornitología, 2000); su plumaje gris pizarra uniforme, algo más pálido por debajo, especialmente en el abdomen y con mandíbula inferior amarillenta (Hilty & Brown, 1986). Es una especie confiada y pasiva; habita en árboles secos (Narosky & Yzurieta, 2010); bordes de bosques primarios o secundarios, bosques abiertos de alisos o Eucaliptos (Asociación Bogotana de Ornitología, 2000). Tiene un amplio rango de distribución, la tendencia de la población es estable y se encuentra en categoría de amenaza de Preocupación Menor (BirdLife International and NatureServe, 2014).

associatedParty

role:

personnelldentifier:

personnelDirectory: homepage:

email: enataliagomezm@gmail.com

phone: postalCode: country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint: location:

thumbnailURL:

mediaURL: http://avibase.bsc-eoc.org/species.jsp?lang=ES&avibaseid=ECF2FBB2B62909A8

description:

subject

source: http://avibase.bsc-eoc.org

audience

bibliographicCitation: rightsHolder:

rights: license: modified: created:

agent

mimeType: dataType: identifier:

_id

_bsontype: ObjectID id: VÄç8ð lg#ç8à