

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e04ac283c45700544e3cb7

Fecha de creación: 2016-03-09T16:09:44.491Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Esta especie habita en bosques húmedos.

[commonNamesAtomized](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:  
dataFormat:  
identifier:  
usedBy:

usedIn

distributionUnstructured:  
endDate:  
startDate:

## ancillaryData

type:  
language: Español  
name: Avispa negra  
creation\_date: 2016-03-09T16:09:44.491Z

usesManagementAndConservation

## ancillaryData

managementAndConservationUnstructured:

## managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
contentType:  
dataFormat:  
identifier:

fullDescriptionUnstructured: Prominencia frente a la fovea pronotal formando un borde afilado; especie principalmente negra con patas, tibia anterior y flagelo rojizos. Placa dorsal del mesepisterno mucho más estrecha y la sutura forma un ángulo con la parte anterior por debajo de los 60°. Escutelo con una línea central muy notoria, algunas veces superficial; mesoescuto con un margen continuo desde atrás hasta la parte anterior de la tégula; coloración oscura y superficie corporal brillante.

briefDescription:

## references

keywords: Agelaia angulata angulata,Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia

fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca trilli, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistrata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raii raii, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea

## editors

institution:  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
pages: 585  
issue:  
year: 1978

## authors

authors: Richards, O. W.  
title: The social wasps of the Americas excluding the Vespinae  
source:  
type: book

tags: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca trilli, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus weyrauchi, Polybia bistrata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raii raii, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia

fuscatus,Protopolybia rugulosa,Protopolybia sedula,Pseudopolybia compressa,Pseudopolybia vespiceps,Synoeca septentrionalis,Synoeca surinama,Synoeca virginea  
abstract:

identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:  
microReference:  
year:  
  
publishedIn  
  
simple:  
source:  
datatype:  
identifier:  
  
specialAuthorship

[authors](#)

[year](#)

simple:  
  
canonicalAuthorship

[authors](#)

[year](#)

simple:  
simple:  
  
canonicalName  
  
infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: Agelaia angulata angulata  
rank: SUBSPECIES  
simple: Agelaia angulata subsp. angulata  
  
attributes  
  
nomenclaturalCode:  
isAnamorphic:  
id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:  
additionalInformation:

[reference](#)

geoPoint:  
location:  
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: NUB Generator [autonym]

[audience](#)

bibliographicCitation:  
rightsHolder:  
rights:  
license:  
title:  
modified: 2016-03-09T16:09:24.490Z  
created: 2016-03-09T16:09:24.490Z

[agent](#)

homepage:  
role:  
email:  
phone:  
address:  
position:  
organisation:  
lastName:  
firstName:  
mimeType:  
dataType:  
identifier:  
parentTaxon: Agelaia angulata  
higherClassification:  
infraspecificEpithet:  
specificEpithet:  
taxonRank: SUBSPECIES  
subGenus:  
genus: Agelaia  
family: Vespidae  
order: Hymenoptera  
classHierarchy: Insecta  
phylum: Arthropoda  
kingdom: Animalia  
recommended:  
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
reproductionUnstructured:

0

dataObject:  
additionalInformation:

## [reference](#)

geoPoint:  
location:  
thumbnailURL:

## [mediaURL](#)

description:

## [subject](#)

## [audience](#)

rightsHolder:  
rights:  
license:  
modified:  
created:

## [agent](#)

mimeType:  
dataType:  
identifier:  
relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedBy: TraitBank  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue: Sexual  
measurementType: Tipo de Reproducción  
measurementID:



molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataFormat:

identifier:

interactionsUnstructured: Su alta tasa de abundancia en las zonas donde habita ha determinado que existan muchas especies imitadoras.

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataFormat:

identifier:

feedingUnstructured: Su fuente principal de proteína es la carroña y se les ha observado tomando presas de

telarañas; cuando están recolectando alimento de un animal muerto, agreden cualquier otro insecto que se acerque, una vez cortan un trozo de carne del tamaño de su tórax, lo amasan y luego vuelan hacia su nido.

0

0

strategy: Depredadores

type: Heterótrofos - Omnívoros

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

behaviorUnstructured: Es una subespecie sociable y de hábitos carroñeros. Sus nidos son estructuras localizadas bajo tierra que poseen particulares panales cilíndricos.

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

## measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

## endemicAtomized

## distribution

## ancillaryData

dataObject:  
additionalInformation:

## reference

geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

mimeType:  
dataType:  
identifier:

distributionUnstructured: Se distribuye en en el continente sudamericano en los paises de Venezuela, Guyana, Guyana Francesa, Surinam, Brasil, Perú, Ecuador, Bolivia y Colombia. En el país se ha registrado en las regiones

de la Amazonia, Andina, Pacífica y Caribe. Esta especie alcanza distribuciones altitudinales de hasta 2350 msnm.

distributionAtomized

stateProvince: Amazonas

ancillaryData

locality:  
municipality:  
country: Colombia

temporalCoverage

endDate:  
startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured:  
areaOfOccupancy:  
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

measurementOrFact

relatedTo:

measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

abstract: Esta subespecie perteneciente a la familia Vespidae se distribuye en 11 departamentos del país, en as regiones de la Amazonia, Andina Pacífica y Caribe. Es una especie de avispa muy sociable y de hábitos carroñeros. Presenta una coloración oscura y su superficie corporal es brillante.

## associatedParty

role:  
personnelIdentifier:  
personnelDirectory:  
homepage:  
email: enataliagomezm@gmail.com  
phone:  
postalCode:  
country: Colombia  
state: Bogotá Distrito Capital  
city: Bogotá D.C.  
address:  
organisation:  
position:  
lastName: Gómez Melendro  
firstName: Edith Natalia

## ancillaryData

dataObject:  
additionalInformation: Información de tipos: El tipo es una hembra y está depositado en el Universitetes Zoologiske Museum, Copenage, Dinamarca (UZMC).  
Registros biológicos: Colección Entomológica, Instituto de Ciencias Naturales, Universidad Nacional de Colombia. Bogotá, Colombia <http://www.biovirtual.unal.edu.co/colecciones/datos/animalia/consultasMammalia.jsp?clase=4>  
Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".

## reference

geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:  
rightsHolder:  
rights:  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

agent

mimeType:  
dataType:  
identifier:

\_id  
\_bsontype: ObjectID  
id: VàJÄfÄW DãËp