

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d464cc3c16479905cba883

Fecha de creación: 2016-02-29T15:33:34.449Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Bosque húmedo tropical, bosque secundario, cultivos (Hilty y Brown, 2001).

Sotobosques, márgenes, bosquetes secundarios y claros de bosque especialmente en las tierras bajas de ambas estribaciones de los Andes, tanto al oriente como al occidente de Ecuador (Ridgely y Greenfield 2001); también en jardines y parques urbanos y semiurbanos (Olmedo, 2001).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Francés

name: Grive à dos olive

creation_date: 2016-02-29T15:33:34.449Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

fullDescriptionUnstructured: 18 a 19 cm de longitud. Su plumaje por encima es color oliva pardusco uniforme, en algunos individuos casi bermejos; mejillas y anillo ocular usualmente prominente; partes bajas del cuerpo de color blanco, pecho y lados anteriores profusamente punteados de negruzco (Hilty y Brown, 2001).

briefDescription:

[references](#)

taxonRecordName

[ancillaryData](#)

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

replacementNameFor

microReference:

source:

datatype:

identifier:

source:

datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: (Nuttall, 1840)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:

ref:
uninomial:
simple: Catharus ustulatus
rank: SPECIES

simple: Catharus ustulatus (Nuttall, 1840)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

ancillaryData

synonymStatus:
synonymName
anyTwo:

anyOne

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:

identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:
datatype:
identifier:

[combinationAuthorship](#)

basionymAuthorship

simple:

authorship

simple:
simple: Nuttall, 1840

epithet

infraspecificEpithet:
specificEpithet:
infragenericEpithet:

genus

linkType:
ref:
uninomial:
simple: Turdus ustulatus
rank: SPECIES
simple: Turdus ustulatus Nuttall, 1840
nomenclaturalCode:
isAnamorphic:
id:

[hierarchy](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:
rights:
license:
title:
modified: 2016-02-29T15:33:14.448Z
created: 2016-02-29T15:33:14.448Z

agent

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Catharus
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Catharus
family: Turdidae
order: Passeriformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:

dataType:

identifier:

reproductionUnstructured: Se reproduce de abril a agosto en Norte América (Collar 2005). El nido es una taza pequeña o abultada de pastos, tallos, musgo, corteza y ramas, rellena con hojas, raicillas, hongos, líquenes y/o musgos, colocado usualmente en el sotobosque, particularmente en matorral deciduo o árboles jóvenes de coníferas. La puesta es de 1-5 huevos, principalmente de 3-4; el periodo de incubación dura entre 10-14 días y los polluelos abandonan el nido a los 10-14 días (Collar 2005).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

feedingUnstructured: Se alimenta de invertebrados y fruta; invertebrados principalmente en primavera y verano boreales, y frutas en otoño e invierno (Collar 2005).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Timido y usualmente visto cerca de cubierta densa; a veces con bandadas mixtas.

Principalmente en nivel medio o más bajo, pero arriba del suelo (Hilty y Brown, 2001).

El canto de *C. ustulatus* consiste en frases cortas, musicales y líquidas: quii-bïrr-rre, quii-bïrr-rre-rrerri-rrerri...

(Olmedo, 2001). Canta débilmente a fin de marzo y abril (Hilty y Brown, 2001).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:

dataType:

identifier:

distributionUnstructured: Argentina, Aruba, Bahamas, Belice, Bermuda, Bolivia, Brasil, Canadá, Colombia, Costa Rica, Cuba, República Dominicana, Ecuador, El Salvador, Guatemala, Haití, Honduras, México, Nicaragua, Panamá, Perú, Estados Unidos, Venezuela (BirdLife 2010), hasta 2700 msnm (Hilty y Brown, 2001). Se distribuye en todo el país (Hilty y Brown 2001).

[distributionAtomized](#)

stateProvince: Córdoba

[ancillaryData](#)

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

populationBiologyUnstructured: Esta especie tiene un rango de distribución amplio, y por lo tanto no se aproxima a los umbrales de la categoría de amenaza vulnerable bajo el criterio de tamaño del rango (BirdLife 2010).

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: 18-19 cm. Pico oscuro con punta amarillenta, patas rosadas, anillo ocular pálido conspicuo. Pardo oliva a gris oliva en el dorso, lores y mejillas color crema. Vientre color blanco cremoso, con puntos oscuros en la garganta y pecho; flancos más oscuros. El Zorzal Carigrís (*C. minimus*), también migratorio, es similar pero tiene la cara más grisácea y un anillo ocular menos prominente (por tanto, luce menos “anteojudo” que *C. ustulatus*) (Olmedo, 2001).

[associatedParty](#)

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomez@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto “Definición y conformación de la PIRAA, a partir del fortalecimiento del nodo subnacional de Colombia: Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIATAC”, como parte del desarrollo del programa BIOCAN. A Thomas McNish por la fotografía de esta especie.
Imagen: BIR-3942 Catharus ustulatus.jpg

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder: Nancy Carolina Rojas Riaño
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

mimeType:
dataType:
identifier:

_id
_bsontype: ObjectID
id: VÔdl< G™\0°0