

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56c7af0cf0106c67230e73fd

Fecha de creación: undefined

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

0

keywords: Accipiter striatus,Actitis macularius,Adelomyia melanogenys,Aglaiocercus kingii,Amblycercus holosericeus,Ampelion rubrocristatus,Anas andium,Anas clypeata,Anas discors,Andigena nigrirostris,Anisognathus igniventris,Ardea alba,Ardea herodias,Arremon assimilis,Asio flammeus,Asio stygius,Asthenes flammulata,Asthenes fuliginosa,Astragalinus psaltria,Atlapetes pallidinucha,Atlapetes schistaceus,Bubulcus ibis,Buteo platypterus,Buteo swainsoni,Butorides striata,Butorides virescens,Cairina moschata,Calidris bairdii,Calidris melanotos,Cardellina canadensis,Catamenia analis,Catamenia inornata,Cathartes aura,Catharus minimus,Catharus ustulatus,Chaetocercus mulsant,Chaetura pelagica,Chalcostigma heteropogon,Chordeiles minor,Chrysomus icterocephalus,Cinclodes albidiventris,Cinclus leucocephalus,Cinnycerthia unirufa,Cistothorus apolinari,Cistothorus platensis,Coccyzua pumila,Coccyzus americanus,Coccyzus melacoryphus,Coeligena bonapartei,Colibri coruscans,Colibri thalassinus,Columba livia,Columbina talpacoti,Conirostrum rufum,Conirostrum sitticolor,Contopus cinereus,Contopus fumigatus,Contopus sordidulus,Contopus virens,Coragyps atratus,Crotophaga ani,Crotophaga major,Dendrocygna autumnalis,Dendrocygna viduata,Diglossa albilatera,Diglossa cyanea,Diglossa humeralis,Diglossa lafresnayii,Diglossa sittoides,Dolichonyx oryzivorus,Dubusia taeniata,Egretta caerulea,Egretta thula,Elaenia flavogaster,Elaenia frantzii,Elanus leucurus,Empidonax alnorum,Empidonax traillii,Ensifera ensifera,Eriocnemis cupreiventris,Eriocnemis vestita,Eupsittula pertinax,Falco columbarius,Falco peregrinus,Falco sparverius,Forpus conspicillatus,Fulica americana,Gallinago nobilis,Gallinula galeata,Gallinula melanops,Geothlypis philadelphia,Geranoaetus melanoleucus,Glaucidium jardinii,Grallaria quitesis,Grallaria ruficapilla,Grallaria squamigera,Haplospiza rustica,Heliangelus amethysticollis,Heliangelus exortis,Hellmayrea gularis,Hemispingus superciliaris,Hemispingus verticalis,Henicorhina leucophrys,Hirundo rustica,Icterus chrysater,Icterus icterus,Icterus nigrogularis,Ixobrychus exilis,Jacana jacana,Lafresnaya lafresnayi,Leiothlypis peregrina,Leptasthenura andicola,Lesbia nuna,Lesbia victoriae,Leucophaeus atricilla,Machetornis rixosa,Margarornis squamiger,Mecocerculus leucophrys,Megascops choliba,Melanerpes rubricapillus,Merganetta armata,Metallura tyrianthina,Milvago chimachima,Mimus gilvus,Mniotilta varia,Molothrus bonariensis,Muscisaxicola alpinus,Myiarchus crinitus,Myiodynastes luteiventris,Myiotheretes fumigatus,Myiotheretes striaticollis,Myiothlypis nigrocristata,Myornis senilis,Nomonyx dominicus,Nycticorax nycticorax,Ochthoeca cinnamomeiventris,Ochthoeca diadema,Ochthoeca fumicolor,Ochthoeca rufipectoralis,Orochelidon murina,Oxypogon guerinii,Oxyura jamaicensis,Pachyramphus versicolor,Pandion haliaetus,Pardirallus maculatus,Parkesia noveboracensis,Patagioenas fasciata,Petrochelidon pyrrhonota,Phaetusa simplex,Phalacrocorax brasilianus,Pheucticus aureoventris,Pheucticus ludovicianus,Phimosus infuscatus,Phrygilus unicolor,Phyllomyias uropygialis,Piaya cayana,Picoides fumigatus,Pipraeidea melanonota,Pipreola riefferii,Piranga olivacea,Piranga rubra,Pitangus sulphuratus,Pluvialis dominica,Podilymbus podiceps,Porphyrion martinicus,Porzana carolina,Progne tapera,Protonotaria citrea,Pseudocolopteryx acutipennis,Pseudoscops clamator,Pterophanes cyanopterus,Pygochelidon cyanoleuca,Pyrocephalus rubinus,Pyrrhomyias cinnamomeus,Quiscalus lugubris,Rallus semiplumbeus,Ramphocelus dimidiatus,Ramphomicron microrhynchum,Riparia riparia,Rupornis magnirostris,Sayornis nigricans,Serpophaga cinerea,Setophaga castanea,Setophaga cerulea,Setophaga petechia,Setophaga ruticilla,Setophaga striata,Sicalis flaveola,Sicalis luteola,Spiza americana,Sporagra spinescens,Sturnella magna,Synallaxis subpudica,Systellura longirostris,Tangara vitriolina,Thraupis episcopus,Thraupis palmarum,Tringa flavipes,Tringa melanoleuca,Tringa solitaria,Troglodytes aedon,Troglodytes solstitialis,Turdus fuscater,Turdus ignobilis,Tyrannus melancholicus,Tyrannus savana,Tyrannus tyrannus,Tyto alba,Vanellus chilensis,Vireo olivaceus,Vultur gryphus,Zenaida auriculata,Zonotrichia capensis

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigrirostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Cistothorus platensis*, *Coccyzua pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta caerulea*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus superciliosus*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Leucophaeus atricilla*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Myornis senilis*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxypogon guerinii*, *Oxyura jamaicensis*, *Pachyramphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Playa cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

abstract:

last\_modified:

created:

group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
habitatUnstructured: Bastante común en pequeños bosques abiertos, bordes de bosque o en el matorral denso siempre cerca de corrientes de agua. Ocasionalmente a corta distancia del interior del bosque húmedo (Hilty & Brown, 1986).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:  
usedIn  
distributionUnstructured:  
endDate:  
startDate:

[ancillaryData](#)

type:  
language: Alemán  
name: Graumanteltyrann  
usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:  
[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0  
dataObject:  
additionalInformation:  
0

keywords: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus holosericeus*, *Ampelion rubrocristatus*, *Anas andium*, *Anas clypeata*, *Anas discors*, *Andigena nigrirostris*, *Anisognathus igniventris*, *Ardea alba*, *Ardea herodias*, *Arremon assimilis*, *Asio flammeus*, *Asio stygius*, *Asthenes flammulata*, *Asthenes fuliginosa*, *Astragalinus psaltria*, *Atlapetes pallidinucha*, *Atlapetes schistaceus*, *Bubulcus ibis*, *Buteo platypterus*, *Buteo swainsoni*, *Butorides striata*, *Butorides virescens*, *Cairina moschata*, *Calidris bairdii*, *Calidris melanotos*, *Cardellina canadensis*, *Catamenia analis*, *Catamenia inornata*, *Cathartes aura*, *Catharus minimus*, *Catharus ustulatus*, *Chaetocercus mulsant*, *Chaetura pelagica*, *Chalcostigma heteropogon*, *Chordeiles minor*, *Chrysomus icterocephalus*, *Cinclodes albidiventris*, *Cinclus leucocephalus*, *Cinnycerthia unirufa*, *Cistothorus apolinari*, *Cistothorus platensis*, *Coccyzua pumila*, *Coccyzus americanus*, *Coccyzus melacoryphus*, *Coeligena bonapartei*, *Colibri coruscans*, *Colibri thalassinus*, *Columba livia*, *Columbina talpacoti*, *Conirostrum rufum*, *Conirostrum sitticolor*, *Contopus cinereus*, *Contopus fumigatus*, *Contopus sordidulus*, *Contopus virens*, *Coragyps atratus*, *Crotophaga ani*, *Crotophaga major*, *Dendrocygna autumnalis*, *Dendrocygna viduata*, *Diglossa albilatera*, *Diglossa cyanea*, *Diglossa humeralis*, *Diglossa lafresnayii*, *Diglossa sittoides*, *Dolichonyx oryzivorus*, *Dubusia taeniata*, *Egretta caerulea*, *Egretta thula*, *Elaenia flavogaster*, *Elaenia frantzii*, *Elanus leucurus*, *Empidonax alnorum*, *Empidonax traillii*, *Ensifera ensifera*, *Eriocnemis cupreiventris*, *Eriocnemis vestita*, *Eupsittula pertinax*, *Falco columbarius*, *Falco peregrinus*, *Falco sparverius*, *Forpus conspicillatus*, *Fulica americana*, *Gallinago nobilis*, *Gallinula galeata*, *Gallinula melanops*, *Geothlypis philadelphia*, *Geranoaetus melanoleucus*, *Glaucidium jardinii*, *Grallaria quitesis*, *Grallaria ruficapilla*, *Grallaria squamigera*, *Haplospiza rustica*, *Heliangelus amethysticollis*, *Heliangelus exortis*, *Hellmayrea gularis*, *Hemispingus supercilialis*, *Hemispingus verticalis*, *Henicorhina leucophrys*, *Hirundo rustica*, *Icterus chrysater*, *Icterus icterus*, *Icterus nigrogularis*, *Ixobrychus exilis*, *Jacana jacana*, *Lafresnaya lafresnayi*, *Leiothlypis peregrina*, *Leptasthenura andicola*, *Lesbia nuna*, *Lesbia victoriae*, *Leucophaeus atricilla*, *Machetornis rixosa*, *Margarornis squamiger*, *Mecocerculus leucophrys*, *Megascops choliba*, *Melanerpes rubricapillus*, *Merganetta armata*, *Metallura tyrianthina*, *Milvago chimachima*, *Mimus gilvus*, *Mniotilta varia*, *Molothrus bonariensis*, *Muscisaxicola alpinus*, *Myiarchus crinitus*, *Myiodynastes luteiventris*, *Myiotheretes fumigatus*, *Myiotheretes striaticollis*, *Myiothlypis nigrocristata*, *Myornis senilis*, *Nomonyx dominicus*, *Nycticorax nycticorax*, *Ochthoeca cinnamomeiventris*, *Ochthoeca diadema*, *Ochthoeca fumicolor*, *Ochthoeca rufipectoralis*, *Orochelidon murina*, *Oxypogon guerinii*, *Oxyura jamaicensis*, *Pachyramphus versicolor*, *Pandion haliaetus*, *Pardirallus maculatus*, *Parkesia noveboracensis*, *Patagioenas fasciata*, *Petrochelidon pyrrhonota*, *Phaetusa simplex*, *Phalacrocorax brasilianus*, *Pheucticus aureoventris*, *Pheucticus ludovicianus*, *Phimosus infuscatus*, *Phrygilus unicolor*, *Phyllomyias uropygialis*, *Piaya cayana*, *Picoides fumigatus*, *Pipraeidea melanonota*, *Pipreola riefferii*, *Piranga olivacea*, *Piranga rubra*, *Pitangus sulphuratus*, *Pluvialis dominica*, *Podilymbus podiceps*, *Porphyrio martinicus*, *Porzana carolina*, *Progne tapera*, *Protonotaria citrea*, *Pseudocolopteryx acutipennis*, *Pseudoscops clamator*, *Pterophanes cyanopterus*, *Pygochelidon cyanoleuca*, *Pyrocephalus rubinus*, *Pyrrhomyias cinnamomeus*, *Quiscalus lugubris*, *Rallus semiplumbeus*, *Ramphocelus dimidiatus*, *Ramphomicron microrhynchum*, *Riparia riparia*, *Rupornis magnirostris*, *Sayornis nigricans*, *Serpophaga cinerea*, *Setophaga castanea*, *Setophaga cerulea*, *Setophaga petechia*, *Setophaga ruticilla*, *Setophaga striata*, *Sicalis flaveola*, *Sicalis luteola*, *Spiza americana*, *Sporagra spinescens*, *Sturnella magna*, *Synallaxis subpudica*, *Systellura longirostris*, *Tangara vitriolina*, *Thraupis episcopus*, *Thraupis palmarum*, *Tringa flavipes*, *Tringa melanoleuca*, *Tringa solitaria*, *Troglodytes aedon*, *Troglodytes solstitialis*, *Turdus fuscater*, *Turdus ignobilis*, *Tyrannus melancholicus*, *Tyrannus savana*, *Tyrannus tyrannus*, *Tyto alba*, *Vanellus chilensis*, *Vireo olivaceus*, *Vultur gryphus*, *Zenaida auriculata*, *Zonotrichia capensis*

institution:

address:

publisher: Princeton University Press

accessed:

websites:

chapter:

series:

pages: 836

issue:

year: 1986

0: Hilty, Steven

1: Brown, William

title: A Guide to the Birds of Colombia

source:

type: book

tags: *Accipiter striatus*, *Actitis macularius*, *Adelomyia melanogenys*, *Agelaiocercus kingii*, *Amblycercus*

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: Mide 12.2 cm. Inconfundible. Rechoncho; pico corto y delgado. La mayoría del plumaje, corona, garganta, dorso, alas y cola, negro pizarra; presenta cejas blancas (falsos ojos). El pecho y vientre castaño rojizo. No presenta barras alares (Hilty & Brown, 1986).

briefDescription:

## references

link: <http://www.birdlife.org/datazone/species/factsheet/22699926>

keywords: *Ochthoeca cinnamomeiventris*

## editors

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

## authors

authors: BirdLife International

title: Species factsheet: *Ochthoeca cinnamomeiventris*

source:

type: miscellany

tags: *Ochthoeca cinnamomeiventris*

abstract:

## identifiers

last\_modified:

created:

group\_id:

profile\_id:

id:

taxonRecordName

## ancillaryData

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:  
note:  
ruleConsidered:  
  
replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

conservedAgainst

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:

identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:

publishedIn

simple:  
source:  
datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:



simple: (Lafresnaye, 1843)

canonicalName

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

linkType:

ref:

uninomial:

simple: Ochthoeca cinnamomeiventris

rank: SPECIES

simple: Ochthoeca cinnamomeiventris (Lafresnaye, 1843)

attributes

nomenclaturalCode:

isAnamorphic:

id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-02-23T01:09:40.278Z

created: 2016-02-23T01:09:40.278Z

[agent](#)

homepage:

role:

email:

phone:

address:

position:  
organisation:  
lastName:  
firstName:  
mimeType:  
dataType:  
identifier:  
parentTaxon: Ochthoeca  
higherClassification:  
infraspecificEpithet:  
specificEpithet:  
taxonRank: SPECIES  
subGenus:  
genus: Ochthoeca  
family: Tyrannidae  
order: Passeriformes  
classHierarchy: Aves  
phylum: Chordata  
kingdom: Animalia  
recommended:  
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:  
additionalInformation:

0

link: <http://www.hbw.com/node/57402>  
keywords: Ochthoeca cinnamomeiventris  
0: del Hoyo, J  
1: Elliott, A  
2: Sargatal, J  
3: Christie, D A  
4: de Juana, E

institution:  
address:  
publisher: Lynx Edicions  
accessed:  
websites:

chapter:  
series:  
issue:  
year: 2004  
0: Farnsworth, A  
1: Langham, G  
title: Slaty-backed Chat-tyrant (Ochthoeca cinnamomeiventris)  
source: Handbook of the Birds of the World Alive  
type: book\_section  
tags: Ochthoeca cinnamomeiventris  
abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

reproductionUnstructured: En Colombia se han registrado huevos en febrero (Huila), juveniles de febrero a agosto y aves con gónadas agrandadas en octubre (Boyacá). El nido es una taza elaborada con musgo, colocado entre las raíces en la grietas, acantilados o de vez en cuando en arbustos (Farnsworth & Langham, 2004).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

0

link: <http://www.hbw.com/node/57402>

keywords: Ochthoeca cinnamomeiventris

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions  
accessed:  
websites:  
chapter:  
series:  
issue:  
year: 2004  
0: Farnsworth, A  
1: Langham, G  
title: Slaty-backed Chat-tyrant (*Ochthoeca cinnamomeiventris*)  
source: Handbook of the Birds of the World Alive  
type: book\_section  
tags: *Ochthoeca cinnamomeiventris*  
abstract:  
last\_modified:  
created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
feedingUnstructured: Consume principalmente insectos; no acompaña bandadas mixtas. Realiza salidas aéreas por lo general a corta distancia, rebusca y con frecuencia regresa a la misma percha (Farnsworth & Langham, 2004).  
environmentalEnvelope  
environmentalEnvelopeUnstructured:  
ecologicalSignificance  
ecologicalSignificanceUnstructured:  
dispersal  
dispersalUnstructured:  
distance:  
structureDispersed:  
type:  
purpose:  
behavior  
0  
dataObject:  
additionalInformation:

0

link: <http://www.hbw.com/node/57402>

keywords: *Ochthoeca cinnamomeiventris*

0: del Hoyo, J

1: Elliott, A

2: Sargatal, J

3: Christie, D A

4: de Juana, E

institution:

address:

publisher: Lynx Edicions

accessed:

websites:

chapter:

series:

issue:

year: 2004

0: Farnsworth, A

1: Langham, G

title: Slaty-backed Chat-tyrant (*Ochthoeca cinnamomeiventris*)

source: Handbook of the Birds of the World Alive

type: book\_section

tags: *Ochthoeca cinnamomeiventris*

abstract:

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Prefiere perchas verticales y bajas al borde de vegetación densa en el sotobosque, por lo general cerca de arroyos; discreto (a causa de plumaje oscuro), pero la voz a menudo delata su presencia (Farnsworth & Langham, 2004).

Realiza llamados persistentes y fuertes a menudo perceptibles cerca de corrientes de agua; su vocalización más común es un fuerte dzzweee-yew, a veces también un silbido alto chew; el canto al amanecer consiste en la misma nota seguida de 3-4 notas tseet (Farnsworth & Langham, 2004).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

## measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

## endemicAtomized

## distribution

## ancillaryData

dataObject:  
additionalInformation:

## reference

link: <http://maps.iucnredlist.org/map.html?id=22699926>  
keywords: Ochthoeca cinnamomeiventris

## editors

institution:  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
issue:  
year: 2014

## authors

authors: BirdLife International and NatureServe  
title: Bird Species Distribution Maps of the World. 2012. Ochthoeca cinnamomeiventris. The IUCN Red List of Threatened Species. Version 2014.3  
source:  
type: miscellany

tags: Ochthoeca cinnamomeiventris

abstract:

## identifiers

last\_modified:

created:

group\_id:

profile\_id:

id:

geoPoint:

location:

thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:

rightsHolder:

rights: Asociación Bogotana de Ornitología

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

## agent

contentType:

dataType:

identifier:

distributionUnstructured: La especie se distribuye principalmente en Sudamérica desde el norte de Venezuela hasta el norte de Bolivia. En Colombia se registra a lo largo de las tres cordilleras (BirdLife International and NatureServe, 2014; Hilty & Brown, 1986).

La especie se encuentra principalmente entre los 1600 y los 3000 msnm (una vez a los 900 msnm en la vertiente Pacífica) (Hilty & Brown, 1986).

## distributionAtomized

stateProvince: Tolima

## ancillaryData

locality:

municipality:

country: Colombia

## temporalCoverage

endDate:

startDate:

## distributionScope

type:

## threatStatus

## ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

## measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

0

dataObject:

additionalInformation:

0

link: <http://www.birdlife.org/datazone/species/factsheet/22699926>

keywords: Ochthoeca cinnamomeiventris

institution:

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 2014

0: BirdLife International

title: Species factsheet: Ochthoeca cinnamomeiventris

source:

type: miscellany

tags: Ochthoeca cinnamomeiventris

abstract:

last\_modified:



created:  
group\_id:  
profile\_id:  
id:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Asociación Bogotana de Ornitología  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
populationBiologyUnstructured: El tamaño de la población mundial no se ha cuantificado, pero esta especie se ha descrito como "bastante común" (BirdLife International, 2014).

## legislation

legislationUnstructured:

## directThreats

directThreatsUnstructured:

## ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

## measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

abstract: El Pitajo Torrentero es una especie de tiránido presente en Sudamérica distribuyéndose desde Venezuela hasta Bolivia. Habita en bosques abiertos, bordes de bosque o en el matorral denso siempre cerca de corrientes de agua (Hilty & Brown, 1986). Consume principalmente insectos y no acompaña bandadas mixtas; prefiere perchas verticales y bajas al borde de vegetación densa en el sotobosque, por lo general cerca de arroyos. Realiza llamados persistentes y fuertes a menudo perceptibles cerca de corrientes de agua (Farnsworth & Langham, 2004).

## associatedParty

role:  
personnelIdentifier:  
personnelDirectory:  
homepage:  
email: enataliagomez@gmail.com  
phone:  
postalCode:  
country: Colombia  
state: Bogotá Distrito Capital  
city: Bogotá D.C.  
address:  
organisation:  
position:  
lastName: Gómez Melendro  
firstName: Edith Natalia

## ancillaryData

dataObject:  
additionalInformation:

## reference

geoPoint:  
location:  
thumbnailURL:  
mediaURL: <https://www.youtube.com/watch?v=OYxQIClk540>  
description:

## subject

source: <https://www.youtube.com>

## audience

bibliographicCitation:  
rightsHolder:  
rights:  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

contentType:  
dataType:  
identifier:  
\_id  
\_bsontype: ObjectId  
id: VÇ~À

đ lg#ç?Đ