

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d477053c16479905cba887
Fecha de creación: 2016-02-29T16:51:20.208Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

habitatUnstructured: Selvas húmedas de tierras firmes y várzea, plantaciones, bosque subtropical, bosque seco tropical y subtropical (Hilty y Brown 2001).

Habita selvas húmedas de tierra firme y várzea, rastrojos en las orillas de arroyos y manglares, también en potreros secos con bordes de rastrojo, y selva de galería (Álvarez y Iannacone 2007).

commonNamesAtomized

ancillaryData

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

ancillaryData

type:

language: Inglés

name: Dark-billed Cuckoo

creation_date: 2016-02-29T16:51:20.208Z

usesManagementAndConservation

ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

fullDescriptionUnstructured: Cuclillo de 25 cm de longitud. El pico es negro, presenta anillo ocular desnudo grisáceo. Por encima su cuerpo es color café grisáceo, la coronilla y la nuca son grisáceas; la máscara es negra; la parte inferior del cuerpo es color ante. Tienen estrecha banda de color gris pálido que va desde debajo de las mejillas hacia abajo por los lados del cuello; la cola es graduada y de color bronce por encima, negruzca debajo con ápices blancos, grandes y redondos (Hilty y Brown 2001).

briefDescription:

[references](#)

taxonRecordName

[ancillaryData](#)

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

replacementNameFor

microReference:

source:

datatype:

identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

[spellingCorrentionOf](#)

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: Vieillot, 1817

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:

simple: Coccyzus melacoryphus
rank: SPECIES
simple: Coccyzus melacoryphus Vieillot, 1817

attributes

nomenclaturalCode:
isAnamorphic:
id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-02-29T16:51:00.205Z
created: 2016-02-29T16:51:00.205Z

[agent](#)

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Coccyzus
higherClassification:

infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Coccyzus
family: Cuculidae
order: Cuculiformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:

reproductionUnstructured: Machos en condición reproductiva han sido registrados en octubre. El nido construido por esta especie tiene forma de plato, es hecho de palitos o tallos gruesos de enredaderas en la base y nervaduras de hojas en el interior, en donde puede tener también líquenes y musgos. Los huevos son de color azul pálido (de la Peña 1983).

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured: Insectívoro (Schulenberg 2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:
structureDispersed:
type:
purpose:

behavior

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
behaviorUnstructured: Migratoria austral (Salaman et al. 2009). Su canto es cu-cu-cu-cu-cu-culp-culp-culop o con

las últimas 3-4 notas ausentes, también un cascabeleo seco dddrrrr (Hilty y Brown 2001).

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataFormat:
identifier:
distributionUnstructured: Argentina, Brasil, Chile, Colombia, Ecuador, Guyana Francesa, Guyana, Paraguay, Perú, Surinam, Trinidad y Tobago, Uruguay (Hilty y Brown 2001), hasta 2100 msnm (Hilty y Brown 2001). En todo el país (Hilty y Brown 2001).

distributionAtomized

stateProvince: Córdoba

ancillaryData

locality:
municipality:
country: Colombia
temporalCoverage
endDate:
startDate:
distributionScope
type:

threatStatus

ancillaryData

threatStatusUnstructured:
threatStatusAtomized
apendiceCITES:
locality:
municipality:
county:
stateProvince:
country: Global
0: IUCN
measurementValue: Preocupación menor (LC)
relatedTo:
measurementRemarks:
measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementType: IUCN
measurementID:

territory

territoryUnstructured:
areaOfOccupancy:
extentOfOccurrence:

populationBiology

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
populationBiologyUnstructured: Sus poblaciones a nivel mundial no han sido cuantificadas pero esta especie ha sido descrita como bastante común (Stotz et al. 1996).

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:

measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:
abstract:

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaría General de la CAN y el Instituto Sinchi para el proyecto "Definición y conformación de la PIRAA, a partir del fortalecimiento del nodo subnacional de Colombia: Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIATAC", como parte del desarrollo del programa BIOCAN.
Recursos multimedia: Sonidos, fotos y videos de esta especie disponibles en The Internet Bird Collection y en Aves de la Pampa.

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder: Nancy Carolina Rojas Riaño
rights:
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:
created:

agent

contentType:
dataType:
identifier:

_id
_bsonType: ObjectID
id: VÔwSÁdy•E"‡