Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e45ff283c45700544e3e33

Fecha de creación: 2016-03-12T18:29:07.975Z

creation\_date: 2016-03-12T18:29:07.975Z usesManagementAndConservation

### ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

0

link: http://www.biodiversitylibrary.org/part/135991

keywords: Battus belus varus,Battus crassus crassus,Battus laodamas copanae,Battus laodamas laodamas,Battus lycidas,Eurytides dolicaon deileon,Eurytides serville acritus,Heraclides anchisiades idaeus,Heraclides androgeus androgeus,Heraclides paeon thrason,Parides erithalion erithalion,Parides eurimedes arriphus,Pterourus birchallii

birchallii institution: address: publisher: accessed: websites: chapter: series:

pages: 411--752

issue: volume: 13 year: 1906

0: Rothschild, Lionel Walter, Jordan, Karl title: A Revision of the American Papilios

source: Novitates Zoologicae

type: article

tags: Battus belus varus,Battus crassus crassus,Battus laodamas copanae,Battus laodamas laodamas,Battus lycidas,Eurytides dolicaon deileon,Eurytides serville acritus,Heraclides anchisiades idaeus,Heraclides androgeus androgeus,Heraclides paeon thrason,Parides erithalion erithalion,Parides eurimedes arriphus,Pterourus birchallii

birchallii abstract: last\_modified: created: group\_id: profile\_id:

id: geoPoint:

location: thumbnailURL: description: source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created: mimeType: dataType: identifier:

fullDescriptionUnstructured: Las alas anteriores de esta subespecie poseen una mancha beige en la punta de las celdas, ya sea por encima y por debajo, o sólo a por debajo, a menudo tiene vestigios, en la parte inferior del disco. El macho generalmente tiene beige de M1 hacia adelante, no hay punto empezando M1 o solo detrás del mismo, rara hay un punto distinto entre M1-M2, la mancha se encuentra a veces en las subespecies que tienen una celda de puntos en la parte superior, las manchas beige son mayores en mujeres, las filas a menudo se extienden hacia atrás hasta SM2, allí empiezan barras submarginales distales de los parches en muchos individuos(Rothschild & Jordan 1906).

Las alas posteriores son algo variables en forma, siendo proporcionalmente menor en algunos ejemplares que en otros, las dos filas de manchas rojas en la parte inferior son a veces muy distantes y pequeñas, estos individuos se reconocen por la posición más proximal de la línea anterior de puntos, por el color pálido de los puntos de R2-M3 de la fila distal, y por los lugares marginales de color beige, al no ser marcados con rojizo; la celda frecuentemente tiene un punto apical rojizo, especialmente en las mujeres (Rothschild & Jordan 1906). briefDescription:

### references

keywords: Battus belus varus,Battus crassus crassus,Battus crassus lepidus,Battus laodamas laodamas,Battus lycidas,Battus polydamas polydamas,Catasticta cinerea laurentina,Eurytides dolicaon deileon,Eurytides orabilis isocharis,Eurytides serville acritus,Eurytides serville columbus,Heraclides anchisiades idaeus,Heraclides androgeus androgeus,Heraclides paeon thrason,Heraclides thoas nealces,Mimoides pausanias pausanias,Papilio polyxenes americus,Parides erithalion browni,Parides erithalion erithalion,Parides lysander orinocoensis,Protographium dioxippus dioxippus,Pterourus birchallii birchallii,Pterourus menatius

#### editors

institution: address:

publisher: Carlec Ltda

accessed: websites: chapter: series: pages: 112 issue:

volume: Tomo I year: 2002

#### authors

authors: Le Crom, J F authors: Constantino, L M authors: Salazar, J A authors: Llorente, J

title: Mariposas de Colombia. Tomo I: Papilionidae

source: type: book tags: Battus belus varus,Battus crassus crassus,Battus crassus lepidus,Battus laodamas laodamas,Battus lycidas,Battus polydamas polydamas,Catasticta cinerea laurentina,Eurytides dolicaon deileon,Eurytides orabilis isocharis,Eurytides serville acritus,Eurytides serville columbus,Heraclides anchisiades idaeus,Heraclides androgeus androgeus,Heraclides paeon thrason,Heraclides thoas nealces,Mimoides pausanias pausanias,Papilio polyxenes americus,Parides erithalion browni,Parides erithalion erithalion,Parides lysander orinocoensis,Protographium dioxippus dioxippus,Pterourus birchallii birchallii,Pterourus menatius abstract:

# identifiers

last\_modified: created: group\_id: profile\_id: id:

taxonRecordName

## ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source:
datatype:
identifier:
source:
datatype:
identifier:
note:

ruleConsidered:

replacementNameFor

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

sanctioned

microReference:

source: datatype: identifier: source: datatype: identifier: note:

# ruleConsidered: laterHomonymOf microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: conservedAgainst basedOn microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: basionym microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered: spellingCorrentionOf typeNameEntity source: datatype: identifier: source: datatype: identifier: source: datatype: identifier: typificacion typeOfType: **lectotypeMicroReferenceVoucher**

# **lectotypePublicationVoucher**

# voucherReference

simple:

microReference:

year:

publishedIn

simple: source: datatype: identifier:

specialAuthorship

### authors

### year

simple:

canonicalAuthorship

#### authors

## year

simple:

# canonicalName

infraspecificEpithet: specificEpithet: infragenericEpithet:

linkType: ref:

uninomial:

simple: Heraclides anchisiades idaeus

rank:

simple: Heraclides anchisiades idaeus

#### attributes

nomenclaturalCode:

isAnamorphic:

id.

# synonymsAtomized

# ancillaryData

synonymStatus:

synonymName

anyTwo:

# anyOne

providerLink: microReference:

# publishedIn

source: datatype: identifier:

### reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

# publishedIn

source: datatype: identifier:

### reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

### publishedIn

source: datatype: identifier:

#### reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:

# publishedIn

source: datatype: identifier:

#### reletedName

source: datatype: identifier: note:

ruleConsidered: microReference:
publishedIn
source: datatype: identifier:
reletedName
source: datatype: identifier: note: ruleConsidered: microReference:
publishedIn
source: datatype: identifier:
reletedName
source: datatype: identifier: note: ruleConsidered:
typeNameEntity
source: datatype: identifier: source: datatype: identifier: source: datatype: identifier: source: datatype: identifier:
typeVoucherEntity
typeOfType: simple: microReference: year: source: datatype: identifier:
combinationAuthorship
basionymAuthorship
simple:
authorship

simple: simple:
epithet
infraspecificEpithet: specificEpithet: infragenericEpithet:
genus
linkType: ref: uninomial: simple: Papilio idaeus (Fabricius, 1793) rank: simple: nomenclaturalCode: isAnamorphic: id:
hierarchy
ancillaryData
dataObject: additionalInformation:
reference
geoPoint: location: thumbnailURL:
mediaURL
description:
subject
source:
audience
bibliographicCitation: rigthsHolder: rights: license: title: modified: 2016-03-12T18:28:47.961Z created: 2016-03-12T18:28:47.961Z
agent
homepage: role: email: phone: address: position:

organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:

parentTaxon: Papilio anchisiades

higherClassification: infraspecificEpithet: idaeus specificEpithet: anchisiades taxonRank: SUBSPECIES

subGenus:

genus: Heraclides family: Papilionidae order: Lepidoptera classHierarchy: Insecta phylum: Arthropoda kingdom: Animalia recommended: classification:

#### commonNameAtomized

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

0

dataObject: additionalInformation:

0

keywords: Aphrissa statira statira,Ascia monuste orseis,Battus crassus crassus,Battus crassus lepidus,Battus laodamas copanae,Battus laodamas laodamas,Battus polydamas polydamas,Colias dimera,Eurytides orabilis isocharis,Eurytides serville acritus,Eurytides serville columbus,Heraclides anchisiades idaeus,Heraclides androgeus androgeus,Heraclides paeon thrason,Heraclides thoas nealces,Mimoides pausanias pausanias,Parides erithalion erithalion,Protesilaus protesilaus,Protographium dioxippus dioxippus,Pterourus birchallii birchallii,Pterourus

cacicus,Pterourus menatius
institution:
address:
publisher: Monografias tercer milenio accessed:
websites:
chapter:
series:
pages: 536
issue:
volume: 8
year: 2008
0: Beccaloni G. W., Viloria A. L., Hall S. K. \& Robinson G.S.
title: Catálogo de las plantas huésped de las mariposas neotropicales
source:
type: book
tags: Aphrissa statira statira, Ascia monuste orseis, Battus crassus crassus, Battus crassus lepidus, Battus laodamas
copanae,Battus laodamas laodamas,Battus polydamas polydamas,Colias dimera,Eurytides orabilis isocharis,Eurytides serville acritus,Eurytides serville columbus,Heraclides anchisiades idaeus,Heraclides androgeus
androgeus, Heraclides paeon thrason, Heraclides thoas nealces, Mimoides pausanias pausanias, Parides erithalion
erithalion, Protesilaus protesilaus, Protographium dioxippus dioxippus, Pterourus birchallii, Pterourus
cacicus,Pterourus menatius
abstract:
last_modified:
created:
group_id:
profile_id:
id:
geoPoint: Iocation:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType: identifier:
interactionsUnstructured: Las plantas hospederas de esta especie son pertenecientes a la familia Rutaceae, Amyris
pinnata, Casimiroa edulis, Casimiroa tetrameria, Citrus sp, Citrus spp, Citrus aurantiifolia, Citrus limón, Citrus
sinensis, Esenbeckia berlandieri, Zanthoxylum sp, Zanthoxylum ekmanii, Zanthoxylum melanostictum, Zanthoxylum
setulosum (Beccaloni et. al 2008).
feeding
feedingUnstructured:
environmentalEnvelope
environmentalEnvelopeUnstructured:
GHVII OHHIGHAIEHVEIODEOHISH UCIUIEU.

ecologicalSignificance

ecological Significance Unstructured:

# dispersal dispersalUnstructured: distance: structureDispersed: type: purpose: behavior behaviorUnstructured: ancillaryData dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier: measurementOrFact relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID: annualCycle annual Cycle Unstructured:habitat habitatUnstructured: endemicAtomized

# ancillaryData

distribution

dataObject:

additionalInformation:

### reference

keywords: Battus belus varus,Battus crassus crassus,Battus crassus lepidus,Battus laodamas laodamas,Battus lycidas,Battus polydamas polydamas,Catasticta cinerea laurentina,Eurytides dolicaon deileon,Eurytides orabilis isocharis,Eurytides serville acritus,Eurytides serville columbus,Heraclides anchisiades idaeus,Heraclides androgeus androgeus,Heraclides paeon thrason,Heraclides thoas nealces,Mimoides pausanias pausanias,Papilio polyxenes americus,Parides erithalion browni,Parides erithalion erithalion,Parides lysander orinocoensis,Protographium dioxippus dioxippus,Pterourus birchallii birchallii,Pterourus menatius

#### editors

institution: address:

publisher: Carlec Ltda

accessed: websites: chapter: series: pages: 112 issue:

volume: Tomo I year: 2002

#### authors

authors: Le Crom, J F authors: Constantino, L M authors: Salazar, J A authors: Llorente, J

title: Mariposas de Colombia. Tomo I: Papilionidae

source: type: book

tags: Battus belus varus,Battus crassus crassus,Battus crassus lepidus,Battus laodamas laodamas,Battus lycidas,Battus polydamas polydamas,Catasticta cinerea laurentina,Eurytides dolicaon deileon,Eurytides orabilis isocharis,Eurytides serville acritus,Eurytides serville columbus,Heraclides anchisiades idaeus,Heraclides androgeus androgeus,Heraclides paeon thrason,Heraclides thoas nealces,Mimoides pausanias pausanias,Papilio polyxenes americus,Parides erithalion browni,Parides erithalion erithalion,Parides lysander orinocoensis,Protographium dioxippus dioxippus,Pterourus birchallii birchallii,Pterourus menatius abstract:

#### identifiers

last\_modified: created:

group\_id:
profile\_id:

id:

geoPoint: location:

thumbnailURL:

#### mediaURL

description:

# subject

source:

#### audience

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified created:

# agent

mimeType: dataType: identifier:

distributionUnstructured: Se distribuye desde el Sur de Estados Unidos, hasta Colombia y norte de Venezuela. Esta subespecie se encuentra en Colombia en la costa pacífica, el río Cauca, alto Magdalena, Magdalena medio, Serranía de San Lucas, y Sierra Nevada de Santa Marta (Le Crom et. al 2002).

### distributionAtomized

stateProvince: Cauca

## ancillaryData

locality: municipality: country: Colombia

temporalCoverage

endDate: startDate:

distributionScope

type:

## threatStatus

territory

territoryUnstructured: areaOfOccupancy: extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint: location:

thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

#### measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Esta mariposa se puede encontrar desde el sur de los Estados Unidos hasta Colombia. Algunas plantas de la familia de las Rutaceas son sus hosopederas. Presentan alas posteriores de formas variables con dos filas de manchas rojas en la parte inferior, y alas anteriores con una mancha beige en las puntas.

# associatedParty

role:

personnelldentifier: personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone: postalCode: country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

# ancillaryData

dataObject:

additionalInformation: Registros biológicos"¢ &Vv—7G&÷2 Vâ Æ 6öÆV66Ÿ6â 66†Ö–GBÔ×VÖÒÂ 6öÆöÖ&– ¢ " d, Ò C B (Corregimiento La Pedrera, Amazonas), 1.974.

Créditos específicos: Los créditos de las imágenes corresponden a la Unidad de Producción Audiovisual (UPA) del Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. En la revisión científica y taxonómica se contó con el apoyo del profesor investigador M. Gonzalo Andrade-C. del Instituto de Ciencias Naturales de la Universidad Nacional de Colombia.

#### reference

geoPo location thumb	
medi	aURL
descri	ption:
subje	ect
source	e:
audie	ence
rights!	e: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) ed:
agen	t
mime dataT identif	ype:
_id	
	ntype: ObjectID _òfÄW Dãã0