Reporte de Fichas para el usuario: edith gómez Ficha ID: 56d4d5c13c16479905cba8d1 Fecha de creación: 2016-02-29T23:35:31.929Z annualCycles annualCycleUnstructured: habitats 0 dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Clara Maria Pérez Gallego license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: habitatUnstructured: Terrestre y arbóreo. Es común en claros, parques, jardines y montes claros, ocasionalmente selva húmeda o bordes. commonNamesAtomized ancillaryData usedBy: usedIn distributionUnstructured: endDate: startDate: ancillaryData type: language: Inglés name: Black-billed Thrush creation date: 2016-02-29T23:35:31.929Z usesManagementAndConservation

# ancillaryData

managementAndConservationUnstructured:

managementAndConservationAtomized

identificationKeys

### fullDescription

fullDescriptionUnstructured: briefDescription:

### references

### taxonRecordName

# ancillaryData

providerSpecificData

anyTwo: providerLink:

publicationStatus

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

### replacementNameFor

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

sanctioned

microReference:

source: datatype: identifier: source: datatype: identifier: note:

ruleConsidered:

### laterHomonymOf

microReference:

source: datatype: identifier: source:

datatype: identifier: note: ruleConsidered:
conservedAgainst
basedOn
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
basionym
microReference: source: datatype: identifier: source: datatype: identifier: note: ruleConsidered:
spellingCorrentionOf
typeNameEntity
source: datatype: identifier:

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

 ${\bf lectotype MicroReference Voucher}$ 

lectotypePublicationVoucher

## voucherReference

simple:

microReference:

year:

# publishedIn simple: source: datatype: identifier: specialAuthorship authors year simple: canonicalAuthorship authors year simple: simple: Sclater, 1857 canonicalName infraspecificEpithet: specificEpithet: infragenericEpithet: linkType: ref: uninomial: simple: Turdus ignobilis rank: SPECIES simple: Turdus ignobilis Sclater, 1857 attributes nomenclaturalCode: isAnamorphic: synonymsAtomized hierarchy ancillaryData dataObject: additionalInformation: reference geoPoint:

location: thumbnailURL:

mediaURL

### description:

### subject

source: The Catalogue of Life, 3rd January 2011

### audience

bibliographicCitation:

rigthsHolder:

rights: license: title:

modified: 2016-02-29T23:35:11.923Z created: 2016-02-29T23:35:11.923Z

### agent

homepage:

role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:

parentTaxon: Turdus higherClassification: infraspecificEpithet: specificEpithet:

taxonRank: SPECIES

subGenus: genus: Turdus family: Turdidae order: Passeriformes classHierarchy: Aves phylum: Chordata kingdom: Animalia recommended: classification:

invasiveness

invasivenessUnstructured:

reproduction

#### 0

dataObject:

additionalInformation:

geoPoint: location:

thumbnailURL: description:

source: bibliographicCitation: rightsHolder: rights: Clara Maria Pérez Gallego license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: reproductionUnstructured: El nido es una taza burda, usualmente con barro a baja altura en arbustos, arboles o tocón; 2 huevos azules fuertemente marcados de café.
molecularData
molecularDataUnstructured:
migratory
migratoryUnstructured:
lifeForm
lifeFormUnstructured:
lifeCycle
lifeCycleUnstructured:
interactions
interactionsUnstructured:
feeding
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: Clara Maria Pérez Gallego license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: feedingUnstructured: Principalmente frugívoro, pero pueden cazar otra variedad de animales.
environmentalEnvelope
environmentalEnvelopeUnstructured:
ecologicalSignificance
ecologicalSignificanceUnstructured:

dispersal
dispersalUnstructured: distance: structureDispersed: type: purpose:
behavior
0
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights Clara Maria Pérez Gallego license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified: created: mimeType: dataType: identifier: behaviorUnstructured: Diurna, común y conspicua en áreas habitadas por el hombre. Individuos o parejas saltan po el suelo en prados o áreas abiertas. Canto débil y poco variado, a menudo introducido por una nota uirt y con frases yur-yur-ui frecuentes.
ancillaryData
dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source: bibliographicCitation: rightsHolder: rights: license: modified: created: mimeType: dataType: identifier:
measurementOrFact
relatedTo: measurementRemarks: measurementMethod: measurementDeterminedDate: measurementUnit:

measurementValue: measurementType: measurementID:

### endemicAtomized

### distribution

### ancillaryData

dataObject:

additionalInformation:

#### reference

geoPoint: location: thumbnailURL:

#### mediaURL

description:

### subject

source:

#### audience

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified: created:

#### agent

mimeType: dataType: identifier:

distributionUnstructured: Desde Colombia a Surinam, N de Bolivia y Amazonia brasileño. En el país se distribuye en el Valle del río Cauca, valles secos o deforestados de la vertiente del Pacífico, valle del Magdalena, E de los Andes hasta S Amazonas. En las regiones Andina, Pacífica y Amazónica hasta los 2800 msnm.

### distributionAtomized

stateProvince: Tolima

### ancillaryData

locality: municipality: country: Colombia temporalCoverage

endDate: startDate:

# distributionScope type: threatStatus ancillaryData threatStatusUnstructured: threatStatusAtomized apendiceCITES: locality: municipality: county: stateProvince: country: Global 0: IUCN measurementValue: Preocupación menor (LC) relatedTo: measurementRemarks: measurementMethod: measurementDeterminedBy measurementDeterminedDate: measurementUnit: measurementAccuracy: measurementType: IUCN measurementID: territory territoryUnstructured: areaOfOccupancy: extentOfOccurrence: populationBiology populationBiologyUnstructured: legislation legislationUnstructured: directThreats directThreatsUnstructured: ancillaryData dataObject: additionalInformation: geoPoint: location: thumbnailURL: description: source:

bibliographicCitation:

rightsHolder: rights:

license: modified: created: mimeType: dataType: identifier:

#### measurementOrFact

relatedTo:

measurementRemarks: measurementMethod:

measurementDeterminedDate:

measurementUnit: measurementAccuracy: measurementValue: measurementType: measurementID:

### associatedParty

abstract:

personnelldentifier: personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone: postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address: organisation: position:

lastName: Gómez Melendro firstName: Edith Natalia

### ancillaryData

dataObject:

additionalInformation: Registros biológicos: En la Central Hidroeléctrica Miel I se registró en los bosques de El Tigre, La Mula, La Clara, Campiña y Cariaño, en los rastrojos de Mirador, Encanto y Reten Norcasia y en las áreas

intervenidas de Cantera y Terraplen.

Imagen: Turdus\_ignobilis.jpg

### reference

geoPoint: location: thumbnailURL:

### mediaURL

description:

### subject

source:

# audience

bibliographicCitation: rightsHolder: Clara Maria Pérez Gallego

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0) modified:

created:

# agent

mimeType: dataType: identifier:

\_id

\_bsontype: ObjectID id: VÔÕÁ< G™\⁰•