

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d5ee693c16479905cba902

Fecha de creación: 2016-03-01T19:32:59.311Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

habitatUnstructured: Común en regiones secas a húmedas, en bosque ralo, bosque secundario alto, bordes de bosques várzea y terra firme (Hilty & Brown 1986). En La Guajira habita en el bosque deciduo tropical, de galería y secundario (Cerrejón 2009-2010).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Alemán

name: Tropeneule

creation_date: 2016-03-01T19:32:59.311Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

fullDescriptionUnstructured: Se han descrito 9 subespecies:

Megascops choliba choliba (Vieillot, 1817)

Megascops choliba crucigerus (Spix, 1824)

Megascops choliba decussatus (Lichtenstein, 1823)

Megascops choliba duida (Chapman, 1929)

Megascops choliba luctisonus (Bangs & T. E. Penard, 1921)

Megascops choliba margaritae (Cory, 1915)

Megascops choliba surutus (L. Kelso, 1941)

Megascops choliba uruguiensis (Hekstra, 1982)

Megascops choliba wetmorei (Brodkorb, 1937)

briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:

datatype:

identifier:

note:

ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:

ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:

datatype:

identifier:

source:

datatype:

identifier:

source:

datatype:

identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:

microReference:

year:

publishedIn

simple:

source:

datatype:

identifier:

specialAuthorship

authors

year

simple:

canonicalAuthorship

authors

year

simple:

simple: (Vieillot, 1817)

canonicalName

infraspecificEpithet:

specificEpithet:

infragenericEpithet:
linkType:
ref:
uninomial:
simple: Megascops choliba
rank: SPECIES
simple: Megascops choliba (Vieillot, 1817)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

ancillaryData

synonymStatus:
synonymName
anyTwo:

anyOne

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:

datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:
source:

datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOfType:
simple:
microReference:
year:
source:
datatype:
identifier:

combinationAuthorship

basionymAuthorship

simple:

authorship

simple:
simple: (Vieillot, 1817)

epithet

infraspecificEpithet:
specificEpithet:
infragenericEpithet:

genus

linkType:
ref:
uninomial:
simple: Otus choliba
rank: SPECIES
simple: Otus choliba (Vieillot, 1817)
nomenclaturalCode:
isAnamorphic:
id:

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-01T19:32:39.309Z

created: 2016-03-01T19:32:39.309Z

agent

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataFormat:

identifier:

parentTaxon: Megascops

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Megascops

family: Strigidae

order: Strigiformes

classHierarchy: Aves

phylum: Chordata

kingdom: Animalia

recommended:

classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

feedingUnstructured: En La Guajira se observó que los individuos forrajea en el dosel y su dieta es carnívora (Cerrejón 2009-2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
behaviorUnstructured: Esta especie tiene hábitos enteramente nocturnos, es común encontrar parejas; descansa sobre ramas u orificios a moderada altura (Hilty & Brown 1986).

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:

dataType:

identifier:

distributionUnstructured: En bases de datos se encontró que:

Esta especie se distribuye en: Bolivia, Brasil, Costa Rica, Argentina, Ecuador, Panamá, Paraguay, Venezuela (Collections MNH SI 2012); Colombia, Costa Rica, Honduras, Venezuela (Arctos 2012)

Las siguientes subespecies se distribuyen en:

Megascops choliba crucigerus Colombia (Collections MNH SI 2012)

Megascops choliba luctisonus Colombia (Collections MNH SI 2012)

Megascops choliba margaritae Colombia (Collections MNH SI 2012)

Megascops choliba wetmorei Bolivia (Collections MNH SI 2012).

En bases de datos y monitoreos de fauna se encontró que:

Esta especie se distribuye en: La Guajira (Cerrejón 2009-2010); Boyacá (Arctos 2012)

Las siguientes subespecies se distribuyen en Colombia:

Megascops choliba crucigerus Amazonas, Arauca, Bogotá, Boyacá, Cundinamarca, Guaviare, Huila, Meta, Santander, Tolima (ICN 2004); Antioquia, Bolívar, Cundinamarca, Norte de Santander, Tolima, Valle del Cauca (Collections MNH SI 2012)

Megascops choliba luctisonus Antioquia, Cauca (ICN 2004); Boyacá, Chocó (Collections MNH SI 2012)

Megascops choliba margaritae Bolívar, Magdalena (ICN 2004); La Guajira, Magdalena (Collections MNH SI 2012).

En todas las regiones naturales del país hasta los 3000 m (Hilty & Brown 1986).

[distributionAtomized](#)

stateProvince: Tolima

[ancillaryData](#)

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

[threatStatus](#)

[ancillaryData](#)

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

[measurementDeterminedBy](#)

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

mimeType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract: Mide 23 cm. Tiene ojos amarillos. El plumaje dorsal de color grisáceo a pardo canela con estrías estrechas parduscas y moteado ante; con una línea extensa y punteada de blanco en escapulares y cobertoras alares; las rémiges con barras pardo canela y pardusco; superciliares y disco facial blanco ante con conspicuo reborde negro; partes ventrales blanquecinas a canela con algunas estrías negras prominentes cruzadas con líneas finas parduscas (Hilty & Brown 1986).

associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation: Registros biológicos: En el portal de Global Biodiversity Information Facility se reporta que para Colombia hay 359 registros. GBIF

Metadatos: GBIF

Créditos específicos: Ficha elaborada con el apoyo de Carbones del Cerrejón Limited.

Otros recursos en Internet: Arctos, Avibase, AVoCet, Colecciones en Línea, EoL, iSpecies, IUCN Red List, SI

NMNH,

xeno-canto

reference

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:
rightsHolder:
rights: Equipo Coordinador del SiB Colombia
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:

agent

contentType:
dataType:
identifier:

_id
_bsontype: ObjectID
id: VÕî< G™\0'