

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d4d09c3c16479905cba8cf

Fecha de creación: 2016-02-29T23:13:34.188Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Clara Maria Pérez Gallego

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

habitatUnstructured: Esta especie se encuentra en Selva húmeda. Esta especie es aparentemente errante entre áreas quemadas en selvas, macizos, bordes de selva y monte secundario.

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Alemán

name: Schieferpfäffchen

creation\_date: 2016-02-29T23:13:34.188Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Clara Maria Pérez Gallego  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
fullDescriptionUnstructured: Esta especie tiene un largo total promedio 11.4 cm. El macho pico amarillo, principalmente gris pizarra, más negro en alas y cola, pequeñas manchas blancas en el cuello. La hembra por encima café oliva y por debajo más pardusca y pálida.  
briefDescription:

references

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

conservedAgainst

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:  
datatype:

identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion  
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:  
  
publishedIn

simple:  
source:  
datatype:  
identifier:  
  
specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: (Lawrence, 1862)

canonicalName

infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:

ref:  
uninomial:  
simple: Sporophila schistacea  
rank: SPECIES  
simple: Sporophila schistacea (Lawrence, 1862)

attributes

nomenclaturalCode:

isAnamorphic:

id:

[synonymsAtomized](#)

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-02-29T23:13:14.188Z

created: 2016-02-29T23:13:14.188Z

[agent](#)

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataType:

identifier:

parentTaxon: Sporophila

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Sporophila  
family: Emberizidae  
order: Passeriformes  
classHierarchy: Aves  
phylum: Chordata  
kingdom: Animalia  
recommended:  
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Clara Maria Pérez Gallego  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:

reproductionUnstructured: Esta especie hace nido en forma de taza delgada de raíces y crin de caballo, 2 - 4 m en árbol pequeño o palma espinosa; 3 huevos de color blanco grisáceo punteado y manchado más oscuro.

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:  
additionalInformation:

geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Clara Maria Pérez Gallego  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
feedingUnstructured: Esta especie se alimenta de semillas y frutas.

#### environmentalEnvelope

environmentalEnvelopeUnstructured:

#### ecologicalSignificance

ecologicalSignificanceUnstructured:

#### dispersal

dispersalUnstructured:

distance:  
structureDispersed:  
type:  
purpose:

#### behavior

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Clara Maria Pérez Gallego  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
behaviorUnstructured: Esta especie es diurna y en partes altas de arboles. Esta especie son individuos dispersos o parejas, con menos frecuencia pequeños grupos. A menudo se posa en lo alto de árboles. Esta especie emite canto agudo y rápido zit, zit zi z z z z z z zi zi zi.

#### ancillaryData

dataObject:

additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

## measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

## endemicAtomized

## distribution

## ancillaryData

dataObject:  
additionalInformation:

## reference

geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:  
rightsHolder:  
rights: Clara Maria Pérez Gallego  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:



## agent

contentType:

dataType:

identifier:

distributionUnstructured: Esta especie se distribuye desde el S México hasta W Ecuador, N Bolivia, Guayanas y NE Brasil. En el país no tiene una distribución definida, hay registros en la costa Pacífica, en el valle del Cauca, en la Cordillera Central y E Andes. Esta especie se encuentra en la Región Pacífica y Andina hasta los 2000 msnm.

## distributionAtomized

temporalCoverage

endDate:

startDate:

distributionScope

type:

## threatStatus

## ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

## measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

abstract:

[associatedParty](#)

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomezm@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

[ancillaryData](#)

dataObject:

additionalInformation: Registros biológicos: En la Central Hidroeléctrica Miel I se capturo esta especie en red en área intervenida de Terraplén y se registro en los bosques de Campiña y Cariaño.

Imagen: Sporophila\_schistacea.jpg

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:  
rightsHolder: Clara Maria Pérez Gallego  
rights:  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

agent

mimeType:  
dataType:  
identifier:

\_id  
\_bsontype: ObjectID  
id: VÓÐæ< G™\°Œø