

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56d6f74a3c16479905cba917

Fecha de creación: 2016-03-02T14:23:08.878Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Residente común en dehesas y sabanas húmedas desde el páramo bajo y ecotono con límites de vegetación arbórea hacia abajo. Algo local pero actualmente en expansión con la deforestación. (Hilty & Brown 1986). En La Guajira habita en pastizales (Cerrejón 2009-2010).

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Inglés

name: Eastern Meadowlark

creation_date: 2016-03-02T14:23:08.878Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

fullDescriptionUnstructured: Se reconocen 16 subespecies:

Sturnella magna alticola Nelson, 1900

Sturnella magna argutula Bangs, 1899

Sturnella magna auropectoralis G. B. Saunders, 1934

Sturnella magna griscomi Van Tyne & Trautman, 1941

Sturnella magna hippocrepis (Wagler, 1832)

Sturnella magna hoopesi Stone, 1897

Sturnella magna inexpectata Ridgway, 1888

Sturnella magna lilianae Oberholser, 1930

Sturnella magna magna (Linnaeus, 1758)

Sturnella magna meridionalis P. L. Sclater, 1861

Sturnella magna mexicana P. L. Sclater, 1861

Sturnella magna paralius Bangs, 1901

Sturnella magna praticola Chubb, 1921

Sturnella magna quinta Dickerman, 1989

Sturnella magna saundersi Dickerman & A. R. Phillips, 1970

Sturnella magna subulata Griscom, 1934

briefDescription:

[references](#)

taxonRecordName

[ancillaryData](#)

providerSpecificData

anyTwo:

providerLink:

publicationStatus

microReference:

source:

datatype:

identifier:

source:
datatype:
identifier:
note:
ruleConsidered:

replacementNameFor

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

[conservedAgainst](#)

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:

source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:

canonicalAuthorship

authors

year

simple:
simple: (Linnaeus, 1758)

canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninomial:
simple: Sturnella magna
rank: SPECIES
simple: Sturnella magna (Linnaeus, 1758)

attributes

nomenclaturalCode:
isAnamorphic:
id:

synonymsAtomized

ancillaryData

synonymStatus:
synonymName
anyTwo:

anyOne

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:

note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOfTypes:
simple:
microReference:
year:
source:
datatype:
identifier:

[combinationAuthorship](#)

basionymAuthorship

simple:

authorship

simple:
simple: Oberholser, 1930

epithet

infraspecificEpithet:
specificEpithet:
infragenericEpithet:

genus

linkType:
ref:
uninomial:
simple: Sturnella lilianae
rank: SPECIES
simple: Sturnella lilianae Oberholser, 1930
nomenclaturalCode:
isAnamorphic:
id:

[hierarchy](#)

[ancillaryData](#)

dataObject:
additionalInformation:

[reference](#)

geoPoint:
location:
thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:
rightsHolder:
rights:
license:
title:
modified: 2016-03-02T14:22:48.875Z
created: 2016-03-02T14:22:48.875Z

[agent](#)

homepage:
role:
email:
phone:
address:
position:
organisation:
lastName:
firstName:
mimeType:
dataType:
identifier:
parentTaxon: Sturnella
higherClassification:
infraspecificEpithet:
specificEpithet:
taxonRank: SPECIES
subGenus:
genus: Sturnella
family: Icteridae
order: Passeriformes
classHierarchy: Aves
phylum: Chordata
kingdom: Animalia
recommended:
classification:

invasiveness

invasivenessUnstructured:

reproduction

reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

feedingUnstructured: En La Guajira se observó que los individuos forrajean en el suelo y se alimentan de insectos (Cerrejón 2009-2010).

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Principalmente terrestre en dehesas. Camina por el suelo y se agazapa para evitar detección. Canta principalmente desde una percha elevada y conspicua como un poste o un pequeño arbusto. En vuelo alterna unos pocos aletazos rápidos con cortos planeos (Hilty & Brown 1986).

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint:

location:

thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

mimeType:

dataType:

identifier:

distributionUnstructured: En bases de datos se encontró que: Esta especie se distribuye en: Colombia, Costa Rica, Estados Unidos, Honduras, México (Arctos 2012)

Las siguientes subespecies se distribuyen en:

Sturnella magna alticola Costa Rica, México (Arctos 2012)

Sturnella magna argutula Estados Unidos (Arctos 2012)

Sturnella magna hoopesi Estados Unidos (Arctos 2012)

Sturnella magna lilianae Estados Unidos; México (Arctos 2012)

Sturnella magna magna Canadá, Estados Unidos (Arctos 2012)

Sturnella magna meridionalis Colombia (Arctos 2012)

En bases de datos, literatura y monitoreos de fauna se encontró que: Esta especie se distribuye en: La Guajira (Cerrejón 2009-2010); Boyacá, Santander (Arctos 2012)

Las siguientes subespecies se distribuyen en Colombia:

Sturnella magna meridionalis Bogotá, Boyacá, Cundinamarca, Santander (ICN 2004); Boyacá, Cundinamarca, Huila, Norte de Santander, Santander (Collections MNH SI 2012); Boyacá, Cundinamarca, Santander (Arctos 2012)

Sturnella magna paralius Cesar, Magdalena, (ICN 2004)

Sturnella magna praticola Boyacá, Casanare, Meta (ICN 2004)

Se encuentra en las regiones Andina, Caribe y Orinoquía hasta los 3500 m (Hilty & Brown 1986).

distributionAtomized

stateProvince: La Guajira

ancillaryData

locality:

municipality:

country: Colombia

temporalCoverage

endDate:

startDate:

distributionScope

type:

threatStatus

ancillaryData

threatStatusUnstructured:

threatStatusAtomized

apendiceCITES:

locality:

municipality:

county:

stateProvince:

country: Global

0: IUCN

measurementValue: Preocupación menor (LC)

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedBy

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementType: IUCN

measurementID:

territory

territoryUnstructured:

areaOfOccupancy:

extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:
measurementID:

abstract: Mide 24 cm. Rechoncho con cola corta y cuadrada. Pico largo y agudo a ras con la frente. Por encima fuertemente estriado negruzco y ante; coronilla estriada negra y blanca; ventralmente es de color amarillo brillante con una amplia V negra en el pecho. Al volar destella rectrices laterales blancas; a menudo colea nerviosamente al caminar (Hilty & Brown 1986).

associatedParty

role:
personnelIdentifier:
personnelDirectory:
homepage:
email: enataliagomezm@gmail.com
phone:
postalCode:
country: Colombia
state: Bogotá Distrito Capital
city: Bogotá D.C.
address:
organisation:
position:
lastName: Gómez Melendro
firstName: Edith Natalia

ancillaryData

dataObject:
additionalInformation: Créditos específicos: Ficha elaborada con el apoyo de Carbones del Cerrejón Limited.
Otros recursos en internet: Arctos, Avibase, AVoCet, BirdLife International, Colecciones en Línea, EoL, iSpecies, IUCN Red List, SI NMNH, xeno-canto

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights: Equipo Coordinador del SiB Colombia

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:

dataType:

identifier:

_id

_bsontype: ObjectId

id: VÖ÷J< G™\0'p