

Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e1b39a83c45700544e3d2e

Fecha de creación: 2016-03-10T17:49:21.002Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataObject:

identifier:

habitatUnstructured: Es una especie muy común y habita lagunas y pantanos en zonas abiertas como pastizales y otras áreas alteradas.

[commonNamesAtomized](#)

[ancillaryData](#)

usedBy:

usedIn

distributionUnstructured:

endDate:

startDate:

[ancillaryData](#)

type:

language: Inglés

name: Red-faced Dragonlet

creation_date: 2016-03-10T17:49:21.002Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

fullDescriptionUnstructured:
briefDescription:

references

keywords: Erythrodiplax fusca,Erythrodiplax kimminsi,Erythrodiplax unimaculata

editors

institution:
address:
publisher: The Ohio State University (Columbus)
accessed:
websites:
chapter:
series:
pages: 587--588
issue:
volume: 36
year: 1943

authors

authors: Walker, E. M.
title: A revision of the libelluline genus Erythrodiplax (Odonata)
source: Annals of the Entomological Society of America
type: article
tags: Erythrodiplax fusca,Erythrodiplax kimminsi,Erythrodiplax unimaculata
abstract:

identifiers

last_modified:
created:
group_id:
profile_id:
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:
providerLink:

publicationStatus

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:

ruleConsidered:
replacementNameFor
microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

sanctioned

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

laterHomonymOf

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

conservedAgainst

basedOn

microReference:
source:
datatype:
identifier:
source:
datatype:
identifier:
note:
ruleConsidered:

basionym

microReference:
source:
datatype:
identifier:
source:

datatype:
identifier:
note:
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:
datatype:
identifier:
source:
datatype:
identifier:
source:
datatype:
identifier:

typificacion
typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:
microReference:
year:

publishedIn

simple:
source:
datatype:
identifier:

specialAuthorship

authors

year

simple:
canonicalAuthorship

authors

year

simple:
simple: Rambur, 1842
canonicalName

infraspecificEpithet:
specificEpithet:
infragenericEpithet:
linkType:
ref:
uninominal:
simple: Erythrodiplax fusca
rank: SPECIES
simple: Erythrodiplax fusca Rambur, 1842

attributes

nomenclaturalCode:
isAnamorphic:
id:

[synonymsAtomized](#)

[ancillaryData](#)

synonymStatus:
synonymName
anyTwo:

[anyOne](#)

providerLink:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:
microReference:

publishedIn

source:
datatype:
identifier:

reletedName

source:
datatype:
identifier:
note:
ruleConsidered:

typeNameEntity

source:
datatype:
identifier:

source:
datatype:
identifier:
source:
datatype:
identifier:

typeVoucherEntity

typeOf type:
simple:
microReference:
year:
source:
datatype:
identifier:

combinationAuthorship

basionymAuthorship

simple:

authorship

simple:
simple: Rambur, 1842

epithet

infraspecificEpithet:
specificEpithet:
infragenericEpithet:

genus

linkType:
ref:
uninomial:
simple: Erythrodiplax incompta
rank: SPECIES
simple: Erythrodiplax incompta Rambur, 1842
nomenclaturalCode:
isAnamorphic:
id:

hierarchy

ancillaryData

dataObject:
additionalInformation:

reference

geoPoint:
location:
thumbnailURL:

mediaURL

description:

subject

source: The Catalogue of Life, 3rd January 2011

audience

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-10T17:49:00.996Z

created: 2016-03-10T17:49:00.996Z

agent

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

mimeType:

dataType:

identifier:

parentTaxon: Erythrodiplax

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Erythrodiplax

family: Libellulidae

order: Odonata

classHierarchy: Insecta

phylum: Arthropoda

kingdom: Animalia

recommended:

classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
reproductionUnstructured: La cópula dura alrededor de un minuto y ocurre sobre la vegetación. La hembra oviposita sola, pero el macho vuela cerca a ella vigilándola. La hembra pone los huevos tocando el agua con la punta del abdomen repetidamente.

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights: Nancy Carolina Rojas Riaño
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)
modified:
created:
mimeType:
dataType:
identifier:
feedingUnstructured:

0

0

strategy: Depredadores

type:

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Los machos de esta especie presentan territorialidad. Se posan muy cerca del agua (15 a 20 cm), si está soleado son muy activos y pueden perseguirse con frecuencia, si se nubla reposan.

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:

measurementRemarks:

measurementMethod:

measurementDeterminedDate:

measurementUnit:

measurementAccuracy:

measurementValue:

measurementType:

measurementID:

[endemicAtomized](#)

[distribution](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:

dataType:

identifier:

distributionUnstructured: Se ha registrado en el Continente Americano en los Estados Unidos, México, Guatemala, Belice, El Salvador, Honduras, Nicaragua, Costa Rica, Panamá, Colombia, Ecuador, Perú, Bolivia, Venezuela, Trinidad y Tobago, Guyana, Guyana Francesa, Surinam, Brasil, Paraguay, Uruguay y Argentina. En Colombia se han reportado poblaciones de este insecto en todas las regiones naturales del país.

[distributionAtomized](#)

stateProvince: Tolima

[ancillaryData](#)

locality:
municipality:
country: Colombia

temporalCoverage

endDate:
startDate:

distributionScope

type:

[threatStatus](#)

territory

territoryUnstructured:
areaOfOccupancy:
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:
additionalInformation:
geoPoint:
location:
thumbnailURL:
description:
source:
bibliographicCitation:
rightsHolder:
rights:
license:
modified:
created:
mimeType:
dataType:
identifier:

measurementOrFact

relatedTo:
measurementRemarks:
measurementMethod:
measurementDeterminedDate:
measurementUnit:
measurementAccuracy:
measurementValue:
measurementType:

measurementID:

abstract: Tamaño de la hembra 2.3 a 3.2 cm y del macho 2.5 a 3.4 cm. El macho tiene la cabeza café oscuro, frente rojizo. Tórax dorsalmente negro, lateralmente azul grisáceo. Alas anteriores con una pequeña mancha café oscuro en la base del ala, alas posteriores también con una mancha café oscura de mayor extensión. Mayor parte del abdomen azul grisáceo, últimos tres segmentos negros. Cercos blancuzcos. La hembra tiene ojos café oscuro, frente y cíleo café claro. Tórax café. Alas con manchas similares a las del macho pero café amarillento. Abdomen café claro.

associatedParty

role:

personnelIdentifier:

personnelDirectory:

homepage:

email: enataliagomez@gmail.com

phone:

postalCode:

country: Colombia

state: Bogotá Distrito Capital

city: Bogotá D.C.

address:

organisation:

position:

lastName: Gómez Melendro

firstName: Edith Natalia

ancillaryData

dataObject:

additionalInformation:

reference

geoPoint:

location:

thumbnailURL:

mediaURL: <http://www.biovirtual.unal.edu.co/coleccion/datos/animalia/consultasMammalia.jsp?clase=4>

description:

subject

source: <http://www.biovirtual.unal.edu.co>

audience

bibliographicCitation:

rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

mimeType:

dataType:

identifier:

_id

_bsontype: ObjectId
id: Vá³šƒÄW DãÒà