

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e1fd9183c45700544e3d52

Fecha de creación: 2016-03-10T23:04:55.487Z

annualCycles

annualCycleUnstructured:

habitats

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

habitatUnstructured: Esta especie habita bosques húmedos de Sudamérica.

[commonNamesAtomized](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source:

[audience](#)

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

[agent](#)

contentType:  
dataFormat:  
identifier:  
usedBy:

usedIn

distributionUnstructured:  
endDate:  
startDate:

## ancillaryData

type:  
language: Español  
name: Avispa  
creation\_date: 2016-03-10T23:04:55.487Z

usesManagementAndConservation

## ancillaryData

managementAndConservationUnstructured:

## managementAndConservationAtomized

identificationKeys

fullDescription

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
contentType:  
dataFormat:  
identifier:

fullDescriptionUnstructured: Cuerpo de la hembra negro, dientes mandibulares rojizos. Occipucio débilmente marginado. Tégula café con mancha circular amarilla. Clípeo finamente granuloso con algunas punteaduras ventralmente. Espacio malar 1/3 la longitud entre las fosetas antenales. Quilla pronotal moderadamente fuerte, más alta a los lados. Alas hialinas fuertemente infuscadas, venación café, longitud 8 mm.

briefDescription:

## references

keywords: Agelaia angulata angulata,Agelaia angulicollis,Agelaia brevistigma,Agelaia cajennensis,Agelaia fulvofasciata,Agelaia hamiltoni,Agelaia testacea,Angiopolybia obidensis,Angiopolybia pallens,Angiopolybia

paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Brachymenes wagnerianus, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona guerini, Epipona tatua, Eumenes versicolor filiformis, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Omicron aequale nigrum, Omicron conclamatum, Omicron criticum pulchripictum, Omicron foxi, Omicron globicollis, Omicron graculum atripes, Omicron lustratum, Omicron nanum incarum, Omicron paranympus, Omicron ypsilon, Pachymenes ghilianii olivaceus, Pachymenes orellanae orellanae, Pachymenes orellanae vardyi, Parachartergus amazonensis, Parachartergus colobopteris, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus smithii, Parachartergus weyrauchi, Pararhaphidoglossa clypeolaris, Pararhaphidoglossa confluenta confluenta, Pararhaphidoglossa cressoniana, Pararhaphidoglossa proxima, Polybia bistrata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raii raii, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Santamenes peregrinus, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea, Synoecoides depressus, Zethus brasiliensis fuscatus, Zethus mexicanus lugubris, Zethus mexicanus mexicanus

## editors

institution: Universidad Nacional de Colombia

address:

publisher:

accessed:

websites:

chapter:

series:

issue:

year: 1997

## authors

authors: Sarmiento, C. E.

title: Vespídeos de Colombia (Hymenoptera: Vespidae)

source:

type: thesis

tags: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Brachymenes wagnerianus, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona guerini, Epipona tatua, Eumenes versicolor filiformis, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus alfkeni, Mischocyttarus alfkenii excrucians, Mischocyttarus alfkenii trinitatis, Mischocyttarus carbonarius carbonarius, Mischocyttarus flavicans, Mischocyttarus foveatus, Mischocyttarus injucundus tingomariae, Mischocyttarus insolitus, Mischocyttarus labiatus, Mischocyttarus latissimus, Mischocyttarus metoecus, Mischocyttarus petiolatus, Mischocyttarus pseudomimeticus, Mischocyttarus punctatus, Mischocyttarus tomentosus, Mischocyttarus vaqueroi, Omicron aequale nigrum, Omicron conclamatum, Omicron criticum pulchripictum, Omicron foxi, Omicron globicollis, Omicron graculum atripes, Omicron lustratum, Omicron nanum incarum, Omicron paranympus, Omicron ypsilon, Pachymenes ghilianii olivaceus, Pachymenes orellanae

orellanae, Pachymenes orellanae vardyi, Parachartergus amazonensis, Parachartergus colobopterus, Parachartergus fasciipennis, Parachartergus fraternus, Parachartergus fulgidipennis, Parachartergus griseus, Parachartergus richardsi, Parachartergus smithii, Parachartergus weyrauchi, Pararhaphidoglossa clypeolaris, Pararhaphidoglossa confluenta confluenta, Pararhaphidoglossa cressoniana, Pararhaphidoglossa proxima, Polybia bistrata, Polybia catillifex, Polybia eberhardae, Polybia emaciata, Polybia fastidiosuscula, Polybia jurinei, Polybia liliacea, Polybia micans, Polybia occidentalis, Polybia platycephala sylvestris, Polybia quadricincta, Polybia raii raii, Polybia rejecta, Polybia scrobalis pronotalis, Polybia scrobalis scrobalis, Polybia sericea, Polybia striata, Polybia velutina, Protopolybia acutiscutis, Protopolybia amarella, Protopolybia chartergoides chartergoides, Protopolybia exigua binominata, Protopolybia exigua exigua, Protopolybia fuscatus, Protopolybia rugulosa, Protopolybia sedula, Pseudopolybia compressa, Pseudopolybia vespiceps, Santamenes peregrinus, Synoeca septentrionalis, Synoeca surinama, Synoeca virginea, Synoecoides depressus, Zethus brasiliensis fuscatus, Zethus mexicanus lugubris, Zethus mexicanus mexicanus

abstract:

identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

taxonRecordName

ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:

datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
  
laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

conservedAgainst

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:

identifier:

typificacion

typeOfType:

[lectotypeMicroReferenceVoucher](#)

[lectotypePublicationVoucher](#)

[voucherReference](#)

simple:

microReference:

year:

publishedIn

simple:

source:

datatype:

identifier:

specialAuthorship

[authors](#)

[year](#)

simple:

canonicalAuthorship

[authors](#)

[year](#)

simple:

simple: Zikan, 1949

canonicalName

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

linkType:

ref:

uninomial:

simple: Mischocyttarus adolphi

rank: SPECIES

simple: Mischocyttarus adolphi Zikan, 1949

attributes

nomenclaturalCode:

isAnamorphic:

id:

[synonymsAtomized](#)

ancillaryData

synonymStatus:  
synonymName  
anyTwo:

anyOne

providerLink:  
microReference:  
  
publishedIn  
  
source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:  
microReference:

publishedIn

source:  
datatype:  
identifier:

reletedName

source:  
datatype:  
identifier:  
note:  
ruleConsidered:

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typeVoucherEntity

typeOfType:  
simple:  
microReference:  
year:  
source:  
datatype:  
identifier:

[combinationAuthorship](#)



basionymAuthorship

simple:

authorship

simple:

simple: Zikan, 1935

epithet

infraspecificEpithet:

specificEpithet:

infragenericEpithet:

genus

linkType:

ref:

uninomial:

simple: *Mischocyttarus dukei*

rank: SPECIES

simple: *Mischocyttarus dukei* Zikan, 1935

nomenclaturalCode:

isAnamorphic:

id:

[hierarchy](#)

[ancillaryData](#)

dataObject:

additionalInformation:

[reference](#)

geoPoint:

location:

thumbnailURL:

[mediaURL](#)

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-10T23:04:35.474Z

created: 2016-03-10T23:04:35.474Z

[agent](#)

homepage:

role:

email:  
phone:  
address:  
position:  
organisation:  
lastName:  
firstName:  
mimeType:  
dataType:  
identifier:  
parentTaxon: Mischocyttarus  
higherClassification:  
infraspecificEpithet:  
specificEpithet:  
taxonRank: SPECIES  
subGenus:  
genus: Mischocyttarus  
family: Vespidae  
order: Hymenoptera  
classHierarchy: Insecta  
phylum: Arthropoda  
kingdom: Animalia  
recommended:  
classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
reproductionUnstructured:

0

dataObject:  
additionalInformation:

[reference](#)

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

audience

rightsHolder:  
rights:  
license:  
modified:  
created:

agent

contentType:  
dataType:  
identifier:  
relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedBy: TraitBank  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue: Sexual  
measurementType: Tipo de Reproducción  
measurementID:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:  
additionalInformation:  
geoPoint:  
location:

thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
feedingUnstructured:

0

0

strategy: Depredadores  
type: Heterótrofos - Omnívoros

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:  
structureDispersed:  
type:  
purpose:

behavior

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:

behaviorUnstructured: Las avispas de este género caminan a lo largo de los hilos de las telarañas, presumiblemente gracias a una especial adaptación en los lóbulos tarsales de las patas medias y posteriores. Después de esto las

avispas forrajeras retornan al nido y allí mastican la comida compartiéndola con otras avispas de la colonia, las larvas son alimentadas por regurgitación. El néctar recolectado de una amplia variedad de flores también es llevado al nido y guardado en sus celdas para luego ser consumido por los adultos.

ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

endemicAtomized

distribution

ancillaryData

dataObject:  
additionalInformation:

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

agent

mimeType:  
dataType:  
identifier:  
distributionUnstructured: Estos insectos se distribuye en Colombia y Brasil. En Colombia esta especie se conoce para la región de la Amazónica.

distributionAtomized

stateProvince: Caquetá

ancillaryData

locality:  
municipality:  
country: Colombia  
temporalCoverage  
endDate:  
startDate:

distributionScope  
type:

threatStatus

territory  
territoryUnstructured:  
areaOfOccupancy:  
extentOfOccurrence:  
populationBiology  
populationBiologyUnstructured:  
legislation  
legislationUnstructured:  
directThreats  
directThreatsUnstructured:  
ancillaryData  
dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:

bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

#### measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:  
abstract:

#### associatedParty

role:  
personnelIdentifier:  
personnelDirectory:  
homepage:  
email: enataliagomezm@gmail.com  
phone:  
postalCode:  
country: Colombia  
state: Bogotá Distrito Capital  
city: Bogotá D.C.  
address:  
organisation:  
position:  
lastName: Gómez Melendro  
firstName: Edith Natalia

#### ancillaryData

dataObject:  
additionalInformation: Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto "Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC".  
Información de tipos: El holotipo se encuentra en el Departamento de Entomología de Instituto Oswaldo Cruz (FIOC), Sao Paulo, Brasil.

#### reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

subject

source:

audience

bibliographicCitation:

rightsHolder:

rights:

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

agent

contentType:

dataType:

identifier:

\_id

\_bsontype: ObjectId

id: Váy'fÄW DãÕ