

## Reporte de Fichas para el usuario: edith gómez

Ficha ID: 56e0e72683c45700544e3cf6

Fecha de creación: 2016-03-10T03:17:00.615Z

annualCycles

annualCycleUnstructured:

habitats

habitatUnstructured:

[commonNamesAtomized](#)

creation\_date: 2016-03-10T03:17:00.615Z

usesManagementAndConservation

[ancillaryData](#)

managementAndConservationUnstructured:

[managementAndConservationAtomized](#)

identificationKeys

fullDescription

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

mimeType:

dataType:

identifier:

fullDescriptionUnstructured: Visto de perfil el escutelo es prominente con dos superficies y proyectándose sobre el metanoto y el segmento abdominal I es corto y estrecho pero no en forma de copa. Escutelo corto y ancho que se proyecta sobre el metanoto; amplias manchas amarillas del propodeo.

briefDescription:

[references](#)

keywords: Agelaia angulata angulata, Agelaia angulicollis, Agelaia brevistigma, Agelaia cajennensis, Agelaia fulvofasciata, Agelaia hamiltoni, Agelaia testacea, Angiopolybia obidensis, Angiopolybia pallens, Angiopolybia paraensis, Angiopolybia zischkai, Apoica thoracica, Asteloeca traili, Brachygastra albula, Brachygastra augusti, Brachygastra lecheguana, Brachygastra propodealis, Brachygastra scutellaris, Chartergellus amazonicus, Chartergellus punctatior, Chartergus chartarius, Chartergus globiventris, Chartergus metanotalis, Clypearia sulcata, Clypearia weyrauchi, Epipona tatua, Leipomeles dorsata, Metapolybia acincta, Metapolybia docilis, Metapolybia nigra, Metapolybia suffusa, Mischocyttarus adolphi, Mischocyttarus

alfkeni, *Mischocyttarus alfkenii excrucians*, *Mischocyttarus alfkenii trinitatis*, *Mischocyttarus carbonarius carbonarius*, *Mischocyttarus flavicans*, *Mischocyttarus foveatus*, *Mischocyttarus injucundus injucundus*, *Mischocyttarus insolitus*, *Mischocyttarus labiatus*, *Mischocyttarus latissimus*, *Mischocyttarus metoecus*, *Mischocyttarus petiolatus*, *Mischocyttarus pseudomimeticus*, *Mischocyttarus punctatus*, *Mischocyttarus tomentosus*, *Mischocyttarus vaqueroi*, *Parachartergus amazonensis*, *Parachartergus colobopterus*, *Parachartergus fasciipennis*, *Parachartergus fraternus*, *Parachartergus fulgidipennis*, *Parachartergus griseus*, *Parachartergus richardsi*, *Parachartergus weyrauchi*, *Polybia bistriata*, *Polybia catillifex*, *Polybia eberhardae*, *Polybia emaciata*, *Polybia fastidiosuscula*, *Polybia jurinei*, *Polybia liliacea*, *Polybia micans*, *Polybia occidentalis*, *Polybia platycephala platycephala*, *Polybia quadricincta*, *Polybia raii*, *Polybia rejecta*, *Polybia scrobalis pronotalis*, *Polybia scrobalis scrobalis*, *Polybia sericea*, *Polybia striata*, *Polybia velutina*, *Protopolybia acutiscutis*, *Protopolybia amarella*, *Protopolybia chartergoides chartergoides*, *Protopolybia exigua binominata*, *Protopolybia exigua exigua*, *Protopolybia fuscatus*, *Protopolybia rugulosa*, *Protopolybia sedula*, *Pseudopolybia compressa*, *Pseudopolybia vespiceps*, *Synoeca septentrionalis*, *Synoeca surinama*, *Synoeca virginea*

## editors

institution:  
address:  
publisher:  
accessed:  
websites:  
chapter:  
series:  
pages: 585  
issue:  
year: 1978

## authors

authors: Richards, O. W.  
title: The social wasps of the Americas excluding the Vespinae  
source:  
type: book

tags: *Agelaia angulata angulata*, *Agelaia angulicollis*, *Agelaia brevistigma*, *Agelaia cajennensis*, *Agelaia fulvofasciata*, *Agelaia hamiltoni*, *Agelaia testacea*, *Angiopolybia obidensis*, *Angiopolybia pallens*, *Angiopolybia paraensis*, *Angiopolybia zischkai*, *Apoica thoracica*, *Asteloeca trilli*, *Brachygastra albula*, *Brachygastra augusti*, *Brachygastra lecheguana*, *Brachygastra propodealis*, *Brachygastra scutellaris*, *Chartergellus amazonicus*, *Chartergellus punctator*, *Chartergus chartarius*, *Chartergus globiventris*, *Chartergus metanotalis*, *Clypearia sulcata*, *Clypearia weyrauchi*, *Epipona tatua*, *Leipomeles dorsata*, *Metapolybia acincta*, *Metapolybia docilis*, *Metapolybia nigra*, *Metapolybia suffusa*, *Mischocyttarus adolphi*, *Mischocyttarus alfkeni*, *Mischocyttarus alfkenii excrucians*, *Mischocyttarus alfkenii trinitatis*, *Mischocyttarus carbonarius carbonarius*, *Mischocyttarus flavicans*, *Mischocyttarus foveatus*, *Mischocyttarus injucundus injucundus*, *Mischocyttarus insolitus*, *Mischocyttarus labiatus*, *Mischocyttarus latissimus*, *Mischocyttarus metoecus*, *Mischocyttarus petiolatus*, *Mischocyttarus pseudomimeticus*, *Mischocyttarus punctatus*, *Mischocyttarus tomentosus*, *Mischocyttarus vaqueroi*, *Parachartergus amazonensis*, *Parachartergus colobopterus*, *Parachartergus fasciipennis*, *Parachartergus fraternus*, *Parachartergus fulgidipennis*, *Parachartergus griseus*, *Parachartergus richardsi*, *Parachartergus weyrauchi*, *Polybia bistriata*, *Polybia catillifex*, *Polybia eberhardae*, *Polybia emaciata*, *Polybia fastidiosuscula*, *Polybia jurinei*, *Polybia liliacea*, *Polybia micans*, *Polybia occidentalis*, *Polybia platycephala platycephala*, *Polybia quadricincta*, *Polybia raii*, *Polybia rejecta*, *Polybia scrobalis pronotalis*, *Polybia scrobalis scrobalis*, *Polybia sericea*, *Polybia striata*, *Polybia velutina*, *Protopolybia acutiscutis*, *Protopolybia amarella*, *Protopolybia chartergoides chartergoides*, *Protopolybia exigua binominata*, *Protopolybia exigua exigua*, *Protopolybia fuscatus*, *Protopolybia rugulosa*, *Protopolybia sedula*, *Pseudopolybia compressa*, *Pseudopolybia vespiceps*, *Synoeca septentrionalis*, *Synoeca surinama*, *Synoeca virginea*

abstract:

## identifiers

last\_modified:  
created:  
group\_id:  
profile\_id:  
id:

taxonRecordName

## ancillaryData

providerSpecificData

anyTwo:  
providerLink:

publicationStatus

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

replacementNameFor

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

sanctioned

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

laterHomonymOf

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:

identifier:  
note:  
ruleConsidered:

conservedAgainst

basedOn

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

basionym

microReference:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
note:  
ruleConsidered:

spellingCorrentionOf

typeNameEntity

source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:  
source:  
datatype:  
identifier:

typificacion

typeOfType:

lectotypeMicroReferenceVoucher

lectotypePublicationVoucher

voucherReference

simple:  
microReference:  
year:

publishedIn

simple:  
source:  
datatype:  
identifier:

specialAuthorship

authors

year

simple:  
canonicalAuthorship

authors

year

simple:  
simple: Bequard, 1942  
canonicalName  
infraspecificEpithet:  
specificEpithet:  
infragenericEpithet:  
linkType:  
ref:  
uninomial:  
simple: Brachygastra propodealis  
rank: SPECIES  
simple: Brachygastra propodealis Bequard, 1942

attributes

nomenclaturalCode:  
isAnamorphic:  
id:

synonymsAtomized

hierarchy

ancillaryData

dataObject:  
additionalInformation:

reference

geoPoint:  
location:  
thumbnailURL:

mediaURL

description:

[subject](#)

source: The Catalogue of Life, 3rd January 2011

[audience](#)

bibliographicCitation:

rightsHolder:

rights:

license:

title:

modified: 2016-03-10T03:16:40.614Z

created: 2016-03-10T03:16:40.614Z

[agent](#)

homepage:

role:

email:

phone:

address:

position:

organisation:

lastName:

firstName:

contentType:

dataType:

identifier:

parentTaxon: Brachygastra

higherClassification:

infraspecificEpithet:

specificEpithet:

taxonRank: SPECIES

subGenus:

genus: Brachygastra

family: Vespidae

order: Hymenoptera

classHierarchy: Insecta

phylum: Arthropoda

kingdom: Animalia

recommended:

classification:

invasiveness

invasivenessUnstructured:

reproduction

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
reproductionUnstructured:

molecularData

molecularDataUnstructured:

migratory

migratoryUnstructured:

lifeForm

lifeFormUnstructured:

lifeCycle

lifeCycleUnstructured:

interactions

interactionsUnstructured:

feeding

0

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:  
mimeType:  
dataType:  
identifier:  
feedingUnstructured:

0

0

strategy: Depredadores  
type: Heterótrofos - Omnívoros

environmentalEnvelope

environmentalEnvelopeUnstructured:

ecologicalSignificance

ecologicalSignificanceUnstructured:

dispersal

dispersalUnstructured:

distance:

structureDispersed:

type:

purpose:

behavior

0

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights: Nancy Carolina Rojas Riaño

license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)

modified:

created:

contentType:

dataType:

identifier:

behaviorUnstructured: Esta especie exhibe comportamientos eusociales y depredación.

ancillaryData

dataObject:

additionalInformation:

geoPoint:

location:

thumbnailURL:

description:

source:

bibliographicCitation:

rightsHolder:

rights:

license:

modified:

created:

contentType:

dataType:

identifier:

measurementOrFact

relatedTo:



measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

## endemicAtomized

## distribution

## ancillaryData

dataObject:  
additionalInformation:

## reference

geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:  
rightsHolder:  
rights: Nancy Carolina Rojas Riaño  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

contentType:  
dataType:  
identifier:  
distributionUnstructured: Se distribuye en Colombia, Ecuador, Perú y Bolivia. En el país se han registrado en la región de la Orinoquía y en la Amazonia.

## distributionAtomized

stateProvince: Caquetá

## ancillaryData

locality:  
municipality:  
country: Colombia

temporalCoverage

endDate:  
startDate:

distributionScope

type:

threatStatus

territory

territoryUnstructured:  
areaOfOccupancy:  
extentOfOccurrence:

populationBiology

populationBiologyUnstructured:

legislation

legislationUnstructured:

directThreats

directThreatsUnstructured:

ancillaryData

dataObject:  
additionalInformation:  
geoPoint:  
location:  
thumbnailURL:  
description:  
source:  
bibliographicCitation:  
rightsHolder:  
rights:  
license:  
modified:  
created:  
mimeType:  
dataType:  
identifier:

measurementOrFact

relatedTo:  
measurementRemarks:  
measurementMethod:  
measurementDeterminedDate:  
measurementUnit:  
measurementAccuracy:  
measurementValue:  
measurementType:  
measurementID:

abstract: Estas avispa almacenan grandes cantidades de néctar en sus nidos el cual es tóxico en algunas épocas del año ya que recurren a plantas alucinógenas para su extracción; visita también algodones en busca de néctar y presas.

## associatedParty

role:  
personnelIdentifier:  
personnelDirectory:  
homepage:  
email:  
phone:  
postalCode:  
country: Colombia  
state: Bogotá Distrito Capital  
city: Bogotá D.C.  
address:  
organisation:  
position:  
lastName: Gómez Melendro  
firstName: Edith Natalia

## ancillaryData

dataObject:  
additionalInformation: Información de tipos: Holotipo: hembra coleccionado en Perú y depositado en el Museum of Comparative Zoology, University of Harvard Cambridge, EEUU (MCZ).  
Créditos específicos: Al Instituto Amazónico de Investigaciones Científicas Sinchi. Al Sistema de Información Ambiental Territorial de la Amazonia Colombiana SIAT-AC. La elaboración de esta ficha fue posible gracias al apoyo del gobierno de Finlandia a través del convenio entre la Secretaria General de la CAN y el Instituto Sinchi para el proyecto “Sistema Regional Amazónico de Información Ambiental con énfasis en Biodiversidad de los países miembros de la CAN, (programa BIOCAN), componente SIAT-AC”.

## reference

geoPoint:  
location:  
thumbnailURL:

## mediaURL

description:

## subject

source:

## audience

bibliographicCitation:  
rightsHolder:  
rights:  
license: Atribución - No Comercial - Compartir igual (CC BY-NC-SA 4.0)  
modified:  
created:

## agent

contentType:  
dataType:  
identifier:

\_id

\_bsontype: ObjectId  
id: Vạc&fÄW Dãi